

**NON-RELATIVISTIC EFFECTIVE FIELD THEORIES:
RENORMALIZATION GROUP AND RENORMALONS**

ANTONIO PINEDA

(IFAE, Universitat Autònoma de Barcelona)

**RENORMALIZATION GROUP IN
NON-RELATIVISTIC EFFECTIVE FIELD THEORIES**

Motivation

Motivation

Resummation of logarithms in Quantum Field Theories (a long tale)

Motivation

Resummation of logarithms in Quantum Field Theories (a long tale)

Sudakov logarithms

Resummation of Logarithms in Deep Inelastic scattering

Resummation of logarithms in HQET

Electroweak logarithms

and so on ...

Motivation

Resummation of logarithms in Quantum Field Theories (a long tale)

Sudakov logarithms

Resummation of Logarithms in Deep Inelastic scattering

Resummation of logarithms in HQET

Electroweak logarithms

and so on ...

BUT!!!

WHAT ABOUT THE FIRST QUANTUM-FIELD-THEORY LOG?

THE LAMB SHIFT

$$\delta E \sim m\alpha^4 + m\alpha^5 \ln \alpha + (???)m\alpha^6 \ln^2 \alpha + \dots$$

Summing logs in non-relativistic systems

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} \sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots$$

$$+ \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots$$

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} \sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots$$

$$+ \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots$$

Relevant for...

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\begin{aligned} \frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} &\sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots \\ &\quad + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots \end{aligned}$$

Relevant for...

Determination of the bottom and charm masses

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\begin{aligned} \frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} &\sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots \\ &\quad + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots \end{aligned}$$

Relevant for...

Determination of the bottom and charm masses

Spectroscopy: Hyperfine splitting of heavy quarkonium: η_b , B_c , η_c , ...

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\begin{aligned} \frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} &\sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots \\ &\quad + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots \end{aligned}$$

Relevant for...

Determination of the bottom and charm masses

Spectroscopy: Hyperfine splitting of heavy quarkonium: η_b , B_c , η_c , ...

Decays of heavy quarkonium

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$

$$\Gamma(V_Q(nS) \rightarrow e^+e^-) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\Gamma(P_Q(nS) \rightarrow \gamma\gamma) \sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots)$$

$$\begin{aligned} \frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} &\sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots \\ &\quad + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots \end{aligned}$$

Relevant for...

Determination of the bottom and charm masses

Spectroscopy: Hyperfine splitting of heavy quarkonium: η_b , B_c , η_c , ...

Decays of heavy quarkonium

Heavy quarkonium sum rules

Summing logs in non-relativistic systems

Large logs understood as ratios of scales: $\ln(mv/m) \sim \ln \alpha$, $\ln(mv^2/(mv)) \sim \ln \alpha$.

Resummation of logs: $(\alpha \ln)^n$.

$$\begin{aligned}\delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &\quad + m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \\ \Gamma(V_Q(nS) \rightarrow e^+e^-) &\sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots) \\ \Gamma(P_Q(nS) \rightarrow \gamma\gamma) &\sim m\alpha^3(1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots) \\ \frac{\Gamma(V_Q(nS) \rightarrow e^+e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} &\sim 1 + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots \\ &\quad + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots\end{aligned}$$

Relevant for...

Determination of the bottom and charm masses

Spectroscopy: Hyperfine splitting of heavy quarkonium: η_b , B_c , η_c , ...

Decays of heavy quarkonium

Heavy quarkonium sum rules

$t\bar{t}$ production near threshold: m_t , α_s , Higgs-top coupling

Renormalization group in NRQCD (LL) (Soft running)

Aim: to obtain the running of the NRQCD matching coefficients: $(\alpha_s \ln \frac{m}{\nu})^n$

Relevant for:

- pNRQCD in the perturbative regime
- pNRQCD in the nonperturbative regime
- "Standard" NRQCD

$$\begin{aligned} \mathcal{L}_{NRQCD} = & \bar{\Psi} i \gamma^0 D_0 \Psi + \bar{\Psi} \left\{ \frac{\mathbf{D}^2}{2m} + c_F g \frac{\boldsymbol{\Sigma} \cdot \mathbf{B}}{2m} + c_D g \frac{\gamma^0 (\mathbf{D} \cdot \mathbf{E} - \mathbf{E} \cdot \mathbf{D})}{8m^2} \right. \\ & \left. + i c_S g \frac{\gamma^0 \boldsymbol{\Sigma} \cdot (\mathbf{D} \times \mathbf{E} - \mathbf{E} \times \mathbf{D})}{8m^2} + \frac{\mathbf{D}^4}{8m^3} \right\} \Psi \\ & - \frac{1}{4} c_1 F_{\mu\nu} F^{\mu\nu} + \frac{c_2}{m^2} g F_{\mu\nu} D^2 g F^{\mu\nu} + \frac{c_3}{m^2} g^3 f_{ABC} F_{\mu\nu}^A F_{\mu\alpha}^B F_{\nu\alpha}^C \end{aligned}$$

$$\begin{aligned} \delta \mathcal{L}_{NRQCD} = & \frac{d_{ss}}{m_1 m_2} \psi_1^\dagger \psi_1 \chi_2^\dagger \chi_2 + \frac{d_{sv}}{m_1 m_2} \psi_1^\dagger \boldsymbol{\sigma} \psi_1 \chi_2^\dagger \boldsymbol{\sigma} \chi_2 \\ & + \frac{d_{vs}}{m_1 m_2} \psi_1^\dagger \Gamma^a \psi_1 \chi_2^\dagger \Gamma^a \chi_2 + \frac{d_{vv}}{m_1 m_2} \psi_1^\dagger \Gamma^a \boldsymbol{\sigma} \psi_1 \chi_2^\dagger \Gamma^a \boldsymbol{\sigma} \chi_2 . \end{aligned}$$

Typically, $c_i \sim 1 + \sum_n A_n \left(\alpha_s \ln \frac{m}{\nu} \right)^n$ $d_i \sim \alpha_s \left(1 + \sum_n B_n \left(\alpha_s \ln \frac{m}{\nu} \right)^n \right)$

$\nu_p \gg |\mathbf{p}|$: quark-antiquark relative three-momentum.


$\nu_s \gg |\mathbf{k}|$: gluon three-momentum, transfer momentum between the quark and antiquark.

$m \gg \nu_p \sim \nu_s$

Matching coefficients: $c(\nu_s), d(\nu_s, \nu_p)$

LL $\rightarrow c(\nu_s), d(\nu_s)$

Running ν_s LL: HQET; $1/m$ expansion, $\frac{i}{q^0 + i\epsilon}$



$$\nu_s \frac{d}{d\nu_s} c_D = \frac{\alpha_s}{4\pi} \left[\frac{4C_A}{3} c_D - \left(\frac{2C_A}{3} + \frac{32C_f}{3} \right) c_k^2 - \frac{10C_A}{3} c_F^2 + \frac{8T_{Fn_f}}{3} c_1^{hl} \right],$$

$$\nu_s \frac{d}{d\nu_s} d_{ss} = -2C_f \left(C_f - \frac{C_A}{2} \right) \alpha_s^2 c_k^2,$$

$$\nu_s \frac{d}{d\nu_s} d_{sv} = 0,$$

$$\nu_s \frac{d}{d\nu_s} d_{vs} = 4(C_f - C_A) \alpha_s^2 c_k^2 + \frac{3}{2} \alpha_s^2 C_A c_D,$$

$$\nu_s \frac{d}{d\nu_s} d_{vv} = -\frac{C_A}{2} \alpha_s^2 c_F^2.$$

We define $z = \left[\frac{\alpha_s(\nu_s)}{\alpha_s(m)} \right]^{\frac{1}{\beta_0}} \simeq 1 - 1/(2\pi)\alpha_s(\nu_s) \ln(\frac{\nu_s}{m})$, $\beta_0 = \frac{11}{3}C_A - \frac{4}{3}T_{Fn_f}$

$$c_F(\nu_s) = z^{-C_A},$$

$$c_S(\nu_s) = 2z^{-C_A} - 1,$$

$$c_D(\nu_s) = \frac{9C_A}{9C_A + 8T_{Fn_f}} \left\{ -\frac{5C_A + 4T_{Fn_f}}{4C_A + 4T_{Fn_f}} z^{-2C_A} + \frac{C_A + 16C_f - 8T_{Fn_f}}{2(C_A - 2T_{Fn_f})} \right. \\ \left. + \frac{-7C_A^2 + 32C_A C_f - 4C_A T_{Fn_f} + 32C_f T_{Fn_f}}{4(C_A + T_{Fn_f})(2T_{Fn_f} - C_A)} z^{4T_{Fn_f}/3 - 2C_A/3} \right. \\ \left. + \frac{8T_{Fn_f}}{9C_A} \left[z^{-2C_A} + \left(\frac{20}{13} + \frac{32C_f}{13C_A} \right) \left[1 - z^{\frac{-13C_A}{6}} \right] \right] \right\},$$

$$d_{ss}(\nu_s) = d_{ss}(m) + 4C_f \left(C_f - \frac{C_A}{2} \right) \frac{\pi}{\beta_0} \alpha_s(m) \left[z^{\beta_0} - 1 \right] ,$$

$$d_{sv}(\nu_s) = d_{sv}(m) ,$$

$$d_{vs}(\nu_s) = d_{vs}(m) - (C_f - C_A) \frac{8\pi}{\beta_0} \alpha_s(m) \left[z^{\beta_0} - 1 \right]$$

$$- \frac{27C_A^2}{9C_A + 8T_F n_f \beta_0} \frac{\pi}{\beta_0} \alpha_s(m) \left\{ - \frac{5C_A + 4T_F n_f}{4C_A + 4T_F n_f \beta_0 - 2C_A} \frac{\beta_0}{\beta_0} \left(z^{\beta_0 - 2C_A} - 1 \right) \right.$$

$$+ \frac{C_A + 16C_f - 8T_F n_f}{2(C_A - 2T_F n_f)} \left(z^{\beta_0} - 1 \right)$$

$$+ \frac{-7C_A^2 + 32C_A C_f - 4C_A T_F n_f + 32C_f T_F n_f}{4(C_A + T_F n_f)(2T_F n_f - C_A)}$$

$$\times \frac{3\beta_0}{3\beta_0 + 4T_F n_f - 2C_A} \left(z^{\beta_0 + 4T_F n_f/3 - 2C_A/3} - 1 \right)$$

$$+ \frac{8T_F n_f}{9C_A} \left[\frac{\beta_0}{\beta_0 - 2C_A} \left(z^{\beta_0 - 2C_A} - 1 \right) + \left(\frac{20}{13} + \frac{32C_f}{13C_A} \right) \right.$$

$$\left. \left. \times \left(\left[z^{\beta_0} - 1 \right] - \frac{6\beta_0}{6\beta_0 - 13C_A} \left[z^{\beta_0 - \frac{13C_A}{6}} - 1 \right] \right) \right] \right\} ,$$

$$d_{vv}(\nu_s) = d_{vv}(m) + \frac{C_A}{\beta_0 - 2C_A} \pi \alpha_s(m) \left\{ z^{\beta_0 - 2C_A} - 1 \right\} .$$

Bauer-Manohar; Pineda

One equation for the soft running.

Renormalization group in pNRQCD (LL) (Ultrasoft running)

Aim: to obtain the running of the pNRQCD matching coefficients: $(\alpha_s \ln)^n$,
 $\alpha_s (\alpha_s \ln)^n$

Relevant for:

- **Spectrum:** heavy quarkonium and QED.
- **Currents:** electromagnetic decays.
- **Currents:** Normalization of bottomonium sum rules.
- **Currents:** Normalization of $t\text{-}\bar{t}$ production near threshold.

$$L_{pNRQCD} = L'_{NRQCD} + \int \int d^3x_1 d^3x_2 \psi(x_1) \chi_c(x_2) V(x_1 - x_2) \psi^\dagger(x_1) \chi_c^\dagger(x_2)$$

L'_{NRQCD} , gluons multipole expanded (only ultrasoft gluons).

$$\begin{aligned} \mathcal{L}_{pNRQCD} = & \text{Tr}\{S^\dagger (i\partial_0 - V_s^{(0)}(\mathbf{x})) S + O^\dagger (iD_0 - V_o^{(0)}(\mathbf{x})) O\} \\ & + gV_A(\mathbf{x}) \text{Tr}\{O^\dagger \mathbf{x} \cdot \mathbf{E} S + S^\dagger \mathbf{x} \cdot \mathbf{E} O\} + g \frac{V_B(\mathbf{x})}{2} \text{Tr}\{O^\dagger \mathbf{x} \cdot \mathbf{E} O + O^\dagger O \mathbf{x} \cdot \mathbf{E}\} \\ & - \text{Tr}\{S^\dagger \left(\frac{\mathbf{p}^2}{m} + \sum_n \frac{V_s^{(n)}(\mathbf{x})}{m^n} \right) S - O^\dagger \left(\frac{\mathbf{p}^2}{m} + \sum_n \frac{V_o^{(n)}(\mathbf{x})}{m^n} \right) O\}, \end{aligned}$$

$$V_s^{(0)} \equiv -C_F \frac{\alpha_{V_s}}{r}$$

$$\frac{V_s^{(1)}}{m} \equiv -\frac{C_F C_A D_s^{(1)}}{2mr^2}.$$

$$\begin{aligned} \frac{V_s^{(2)}}{m^2} = & -\frac{C_F D_{1,s}^{(2)}}{2m^2} \left\{ \frac{1}{r}, \mathbf{p}^2 \right\} + \frac{C_F D_{2,s}^{(2)}}{2m^2} \frac{1}{r^3} \mathbf{L}^2 + \frac{\pi C_F D_{d,s}^{(2)}}{m^2} \delta^{(3)}(\mathbf{r}) \\ & + \frac{4\pi C_F D_{S^2,s}^{(2)}}{3m^2} \mathbf{S}^2 \delta^{(3)}(\mathbf{r}) + \frac{3C_F D_{LS,s}^{(2)}}{2m^2} \frac{1}{r^3} \mathbf{L} \cdot \mathbf{S} + \frac{C_F D_{S_{12},s}^{(2)}}{4m^2} \frac{1}{r^3} S_{12}(\hat{\mathbf{r}}), \end{aligned}$$

where $S_{12}(\hat{\mathbf{r}}) \equiv 3\hat{\mathbf{r}} \cdot \boldsymbol{\sigma}_1 \hat{\mathbf{r}} \cdot \boldsymbol{\sigma}_2 - \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2$ and $\mathbf{S} = \boldsymbol{\sigma}_1/2 + \boldsymbol{\sigma}_2/2$.

$\nu_p \gg |\mathbf{p}|$: quark-antiquark relative three-momentum.

$\nu_{us} \gg |\mathbf{k}|$: gluon three-momentum.







$|\mathbf{p}| \gg \nu_{us} \gg \mathbf{p}^2/m$

Matching coefficients: $\tilde{V}(d(\nu_p, \nu_s, m), c(\nu_s, m), \nu_s, \nu_{us}, r) = \tilde{V}(\nu_p, m, \nu_{us}, r) \equiv \tilde{V}(\nu_p, \nu_{us})$.

$\nu_s \frac{d}{d\nu_s} \tilde{V} = 0$; $\nu_s = 1/r$

LL: $\nu_p \frac{d}{d\nu_p} \tilde{V} = 0$

ν_{us} . The computation can be formally organized through the **multipole expansion**.



Corrections to the Green Function

$$G_s(E) = P_s \frac{1}{H - H_I - E} P_s = G_s^{(0)} + \delta G_s$$

From the potential:

$$\delta G_s \sim \frac{1}{H_s - E} \delta V \frac{1}{H_s - E}$$

From ultrasoft gluons:

$$\begin{aligned} \delta G_s &\sim \frac{1}{H_s - E} \int \frac{d^3\mathbf{k}}{(2\pi)^{D-1}} \mathbf{r} \frac{k}{k + H_o - E} \mathbf{r} \frac{1}{H_s - E} \\ &\sim \frac{1}{H_s - E} \mathbf{r} (H_o - E)^3 \left\{ \frac{1}{\epsilon} + \gamma + \ln \frac{(H_o - E)^2}{\nu^2} + C \right\} \mathbf{r} \frac{1}{H_s - E} \end{aligned}$$

$$\nu_{us} \frac{d}{d\nu_{us}} \alpha_{V_s} = \frac{2\alpha_s}{3\pi} V_A^2 \left(\left(\frac{C_A}{2} - C_f \right) \alpha_{V_o} + C_f \alpha_{V_s} \right)^3,$$

$$\nu_{us} \frac{d}{d\nu_{us}} \alpha_{V_o} = \frac{2\alpha_s}{3\pi} V_A^2 \left(\left(\frac{C_A}{2} - C_f \right) \alpha_{V_o} + C_f \alpha_{V_s} \right)^3,$$

$$\nu_{us} \frac{d}{d\nu_{us}} \alpha_s = -\beta_0 \frac{\alpha_s^2}{2\pi},$$

$$\nu_{us} \frac{d}{d\nu_{us}} V_A = 0,$$

$$\nu_{us} \frac{d}{d\nu_{us}} V_B = 0.$$

$$\nu_{us} \frac{d}{d\nu_{us}} C_A D_s^{(1)} = \frac{16\alpha_s}{3\pi} V_A^2 c_k \left[\left(\frac{C_A}{2} - C_f \right) \alpha_{V_o} + C_f \alpha_{V_s} \right] \left[2C_f \alpha_{V_s} + \left(\frac{C_A}{2} - C_f \right) \alpha_{V_o} \right],$$

$$\nu_{us} \frac{d}{d\nu_{us}} D_{d,s}^{(2)} = \frac{16\alpha_s}{3\pi} V_A^2 c_k^2 \left(\frac{C_A}{2} - C_f \right) \alpha_{V_o},$$

$$\nu_{us} \frac{d}{d\nu_{us}} D_{1,s}^{(2)} = \frac{8\alpha_s}{3\pi} V_A^2 c_k^2 \left[\left(\frac{C_A}{2} - C_f \right) \alpha_{V_o} + C_f \alpha_{V_s} \right],$$

and zero for the other matching coefficients (in particular for the spin-dependent potentials).

Soto-Pineda; Pineda

RG equations within an strict expansion in α

$$\nu_{us} \frac{d}{d\nu_{us}} \alpha_{V_s} = \frac{2\alpha_s(\nu_{us})}{3\pi} \left(\frac{C_A}{2}\right)^3 \alpha_s^3(r^{-1}),$$

$$\nu_{us} \frac{d}{d\nu_{us}} \alpha_{V_o} = 0,$$

$$\nu_{us} \frac{d}{d\nu_{us}} C_A D_s^{(1)} = \frac{16\alpha_s(\nu_{us}) C_A}{3\pi} \left(C_f + \frac{C_A}{2}\right) \alpha_s^2(r^{-1}),$$

$$\nu_{us} \frac{d}{d\nu_{us}} D_{1,s}^{(2)} = \frac{8\alpha_s(\nu_{us}) C_A}{3\pi} \alpha_s(r^{-1}),$$

$$\nu_{us} \frac{d}{d\nu_{us}} D_{d,s}^{(2)} = \frac{16\alpha_s(\nu_{us})}{3\pi} \left(\frac{C_A}{2} - C_f\right) \alpha_s(r^{-1}).$$

Initial conditions ($\nu_{us} = 1/r$):

$$\alpha_{V_s}(r^{-1}) = \alpha_s(r^{-1}) \left\{ 1 + (a_1 + 2\gamma_E\beta_0) \frac{\alpha_s(r^{-1})}{4\pi} + \left[\gamma_E (4a_1\beta_0 + 2\beta_1) + \left(\frac{\pi^2}{3} + 4\gamma_E^2 \right) \beta_0^2 + a_2 \right] \frac{\alpha_s^2(r^{-1})}{16\pi^2} \right\},$$

$$D_s^{(1)}(r^{-1}) = \alpha_s^2(r^{-1}),$$

$$D_{1,s}^{(2)}(r^{-1}) = \alpha_s(r^{-1}),$$

$$D_{2,s}^{(2)}(r^{-1}) = \alpha_s(r^{-1}),$$

$$D_{d,s}^{(2)}(r^{-1}) = \alpha_s(r^{-1}) (2 + c_D(r^{-1}) - 2c_F^2(r^{-1})) + \frac{1}{\pi} \left[d_{vs}(r^{-1}) + 3d_{vv}(r^{-1}) + \frac{1}{C_f} (d_{ss}(r^{-1}) + 3d_{sv}(r^{-1})) \right],$$

$$D_{S^2,s}^{(2)}(r^{-1}) = \alpha_s(r^{-1}) c_F^2(r^{-1}) - \frac{3}{2\pi C_f} (d_{sv}(r^{-1}) + C_f d_{vv}(r^{-1})),$$

$$D_{LS,s}^{(2)}(r^{-1}) = \frac{\alpha_s(r^{-1})}{3} (c_S(r^{-1}) + 2c_F(r^{-1})),$$

$$D_{S_{12},s}^{(2)}(r^{-1}) = \alpha_s(r^{-1}) c_F^2(r^{-1}),$$

$$\alpha_{V_o}(r^{-1}) = \alpha_s(r^{-1}),$$

$$V_A(r^{-1}) = 1,$$

The RG improved potentials for the singlet read:

$$\alpha_{V_s}(\nu_{us}) = \alpha_{V_s}(r^{-1}) + \frac{C_A^3}{6\beta_0} \alpha_s^3(r^{-1}) \log \left(\frac{\alpha_s(r^{-1})}{\alpha_s(\nu_{us})} \right),$$

$$D_s^{(1)}(\nu_{us}) = D_s^{(1)}(r^{-1}) + \frac{16}{3\beta_0} \left(\frac{C_A}{2} + C_f \right) \alpha_s^2(r^{-1}) \log \left(\frac{\alpha_s(r^{-1})}{\alpha_s(\nu_{us})} \right),$$

$$D_{1,s}^{(2)}(\nu_{us}) = D_{1,s}^{(2)}(r^{-1}) + \frac{8C_A}{3\beta_0} \alpha_s(r^{-1}) \log \left(\frac{\alpha_s(r^{-1})}{\alpha_s(\nu_{us})} \right),$$

$$D_{2,s}^{(2)}(\nu_{us}) = D_{2,s}^{(2)}(r^{-1}),$$

$$D_{d,s}^{(2)}(\nu_{us}) = D_{d,s}^{(2)}(r^{-1}) + \frac{32}{3\beta_0} \left(\frac{C_A}{2} - C_f \right) \alpha_s(r^{-1}) \log \left(\frac{\alpha_s(r^{-1})}{\alpha_s(\nu_{us})} \right),$$

$$D_{S^2,s}^{(2)}(\nu_{us}) = D_{S^2,s}^{(2)}(r^{-1}),$$

$$D_{LS,s}^{(2)}(\nu_{us}) = D_{LS,s}^{(2)}(r^{-1}),$$

$$D_{S_{12},s}^{(2)}(\nu_{us}) = D_{S_{12},s}^{(2)}(r^{-1}).$$

Soto, Pineda; Pineda

One equation for the ultrasoft running.

OBSERVABLE: NNLL heavy quarkonium mass $O(m\alpha^{4+n} \ln^n \alpha)$ (Pineda; Hoang-Stewart)

$$\begin{aligned} \delta E_{n,l,j}^{\text{pot}}(\nu_{us}) = E_n \alpha_s^2 & \left\{ -\frac{2C_A}{3\beta_0} \left[\frac{C_A^2}{2} + 4C_A C_f \frac{1}{n(2l+1)} + 2C_f^2 \left(\frac{8}{n(2l+1)} - \frac{1}{n^2} \right) \right] \log \left(\frac{\alpha_s(\nu_{us})}{\alpha_s} \right) \right. \\ & + \frac{C_f^2 \delta_{l0}}{3n} \left(-\frac{16}{\beta_0} \left[C_f - \frac{C_A}{2} \right] \log \left(\frac{\alpha_s(\nu_{us})}{\alpha_s} \right) \right. \\ & \quad \left. \left. - \frac{3}{2} (1 + c_D - 2C_F^2) - \frac{3}{2\pi\alpha_s} \left[d_{vs} + 3d_{vv} + \frac{1}{C_f} (d_{ss} + 3d_{sv}) \right] \right) \right\} \\ & - \frac{4C_f^2 \delta_{l0} \delta_{s1}}{3n} \left\{ z^{-2C_A} - 1 + \frac{3C_A}{2\beta_0 - 2C_A} [z^{-\beta_0} - z^{-2C_A}] \right\} \\ & - \frac{(1 - \delta_{l0}) \delta_{s1}}{l(2l+1)(l+1)n} C_{j,l} \frac{C_f^2}{2} \left. \right\}, \end{aligned}$$

where $E_n = -mC_f^2\alpha_s^2/(4n^2)$, $\nu_s = 2a_n^{-1}$ where $2a_n^{-1} = \frac{mC_f\alpha_s(2a_n^{-1})}{n}$, and

$$C_{j,l} = \begin{cases} -\frac{(l+1)}{2l-1} \{4(2l-1)(z^{-C_A} - 1) + (z^{-2C_A} - 1)\} & , j = l-1 \\ -4(z^{-C_A} - 1) + (z^{-2C_A} - 1) & , j = l \\ \frac{l}{2l+3} \{4(2l+3)(z^{-C_A} - 1) - (z^{-2C_A} - 1)\} & , j = l+1. \end{cases}$$

Check with $O(m\alpha^5 \ln \alpha)$ known logs: Brambilla, Vairo, Soto, Pineda; Kniehl, Penin; Hoang, Manohar and Stewart.

Muonic Hydrogen mass at **NNLL**. Check with known logs by Pachucki.

Renormalization group in pNRQCD (NLL) (potential running)

ν_p enters into the game.

The running on ν_p can be obtained from pNRQCD (there is no running of ν_p from NRQCD to pNRQCD). It can be obtained by Quantum mechanics computations. Example: iteration of the potentials. Divergent integrals in $|\mathbf{p}|$ and r .

$$h_s = c_k \frac{\mathbf{p}^2}{m} - C_f \frac{\alpha_{V_s}}{r} - c_4 \frac{\mathbf{p}^4}{4m^3} - \frac{C_f C_A D_s^{(1)}}{2mr^2} - \frac{C_f D_{1,s}^{(2)}}{2m^2} \left\{ \frac{1}{r}, \mathbf{p}^2 \right\} + \frac{C_f D_{2,s}^{(2)}}{2m^2} \frac{1}{r^3} \mathbf{L}^2$$

$$+ \frac{\pi C_f D_{d,s}^{(2)}}{m^2} \delta^{(3)}(\mathbf{r}) + \frac{4\pi C_f D_{S^2,s}^{(2)}}{3m^2} \mathbf{S}^2 \delta^{(3)}(\mathbf{r}) + \frac{3C_f D_{LS,s}^{(2)}}{2m^2} \frac{1}{r^3} \mathbf{L} \cdot \mathbf{S} + \frac{C_f D_{S_{12},s}^{(2)}}{4m^2} \frac{1}{r^3} S_{12}(\hat{\mathbf{r}}),$$

where $C_f = (N_c^2 - 1)/(2N_c)$ and $c_k = c_4 = 1$ (we only use c_4 for tracking of the contribution due to this term). The propagator of the singlet is (formally)

$$\frac{1}{E - h_s}.$$

At leading order (within an strict expansion in α_s) the propagator of the singlet reads

$$\text{||||||||||||||||} = G_c(E) = \frac{1}{E - h_s^{(0)}} = \frac{1}{E - \mathbf{p}^2/m - C_f \alpha_s / r}.$$


If we were interested in computing the spectrum at $O(m\alpha_s^6)$, one should consider the iteration of subleading potentials (δh_s) in the propagator as

follows:

$$G_c(E)\delta h_s G_c(E)\cdots\delta h_s G_c(E).$$

In general, these contributions will produce logarithmic divergences due to potential loops. These divergences can be absorbed in the matching coefficients, $D_{d,s}^{(2)}$ and $D_{S^2,s}^{(2)}$, of the local potentials (proportional to the $\delta^{(3)}(\mathbf{r})$) providing with the renormalization group equations of these matching coefficients in terms of ν_p . Let us explain how it works in detail. Since the singular behavior of the potential loops appears for $\mathbf{p}^2/m \gg \alpha_s/r$, a perturbative expansion in α_s is licit in $G_c(E)$, which can be approximated by

$$\text{—————} = G_c^{(0)}(E) = \frac{1}{E - \mathbf{p}^2/m}.$$



$$\langle \mathbf{r} = 0 | \frac{1}{E - \mathbf{p}^2/m} C_f \frac{\alpha_{V_s}}{r} \frac{1}{E - \mathbf{p}^2/m} | \mathbf{r} = 0 \rangle$$

$$\sim \int \frac{d^d p'}{(2\pi)^d} \int \frac{d^d p}{(2\pi)^d} \frac{m}{\mathbf{p}'^2 - mE} C_f \frac{4\pi\alpha_{V_s}}{\mathbf{q}^2} \frac{m}{\mathbf{p}^2 - mE} \sim -C_f \frac{m^2 \alpha_{V_s}}{16\pi} \frac{1}{\epsilon},$$

where $D = 4 + 2\epsilon$ and $\mathbf{q} = \mathbf{p} - \mathbf{p}'$. This divergence is absorbed in $D_{d,s}^{(2)}$ contributing to its running at NLL order as follows

$$\nu_p \frac{d}{d\nu_p} D_{d,s}^{(2)}(\nu_p) \sim \alpha_{V_s}(\nu_p) D_{d,s}^{(2)2}(\nu_p) + \dots$$

$O(m\alpha^8 \ln^3 \alpha)$ correction to the Hydrogen atom spectrum. Manohar-Stewart;
Pineda

and one equation for the potential running

$|\mathbf{p}| \gg \nu_{us} \gg \mathbf{p}^2/m \rightarrow \nu_{us} = \nu_p^2/m$ (Luke, Manohar, Rothstein)

We can not lower ν_{us} further. Fight between two terms.

$$\frac{1}{\mathbf{p}^2/m + k}$$

$$\begin{aligned} \tilde{V}(c(1/r), d(\nu_p, 1/r), 1/r, \nu_p^2/m, r) &\simeq \tilde{V}(c(\nu_p), d(\nu_p, \nu_p), \nu_p, \nu_p^2/m, \nu_p) \\ &+ \ln(\nu_p r) r \frac{d}{dr} \tilde{V} \Big|_{1/r=\nu_p} + \dots \end{aligned}$$

One equation for the soft running,
one equation for the ultrasoft running,
and one equation for the potential running,
which rules them all and at the hard scale binds them.

Nonrelativistic Sum rules ($b\text{-}\bar{b}$, $c\text{-}\bar{c}$), $t\text{-}\bar{t}$ production near threshold

Determination of m_b , m_t , α_s , Higgs-top yukawa coupling, ...

$$J^\mu = \bar{Q}\gamma^\mu Q = B_1\psi^\dagger\boldsymbol{\sigma}\chi + \dots,$$

$$B_1 = 1 + a_1\alpha_s + a_2\alpha_s^2 + \dots$$

B_1 at NNLO: Hoang(QED); Beneke, Signer, Smirnov; Czarnecki, Melnikov

B_1, B_0 at NLL: Pineda; Hoang, Stewart

B_1/B_0 at NNLL: Penin, Pineda, Smirnov, Steinhauser

B_1, B_0 at NNLL (partial): Pineda, Signer

$$(q_\mu q_\nu - g_{\mu\nu})\Pi(q^2) = i \int d^4x e^{iqx} \langle \text{vac} | J_\mu(x) J_\nu(0) | \text{vac} \rangle$$

$$\Pi(q^2) \sim B_1^2 \langle \mathbf{r} = \mathbf{0} | \frac{1}{E - H} | \mathbf{r} = \mathbf{0} \rangle$$

$$G(0, 0, E) = \sum_{m=0}^{\infty} \frac{|\phi_{0m}(0)|^2}{E_{0m} - E + i\epsilon - i\Gamma_t} + \frac{1}{\pi} \int_0^{\infty} dE' \frac{|\phi_{0E'}(0)|^2}{E_{0E'} - E + i\epsilon - i\Gamma_t}$$

A NNLL renormalization group improved expression of $M(V_Q(nS))$ is also needed in order to obtain expressions for the $t\text{-}\bar{t}$ production near threshold with NNLL accuracy:

$M(V_Q(nS))$ at NNLL: Pineda; Hoang, Stewart

$M(V_Q(nS)) - M(P_Q(nS))$ at NNNLL: Kniehl, Penin, Pineda, Smirnov, Steinhauser

**Relation of the vacuum polarization with $\sigma_{t\bar{t}}$, non-relativistic sum rules
and $\Gamma(V_Q(nS) \rightarrow e^+e^-)$**


$$\Gamma(V \rightarrow e^+e^-) \sim \frac{1}{m^2} B_1^2 |\phi(\mathbf{0})|^2$$

$$\sigma_{t\bar{t}} \sim B_1(\nu)^2 \text{Im}G(0, 0, \sqrt{s}) + \dots$$


$$M_n \equiv \frac{12\pi^2 e_b^2}{n!} \left(\frac{d}{dq^2} \right)^n \Pi(q^2)|_{q^2=0} = \int_0^\infty \frac{ds}{s^{n+1}} R_{b\bar{b}}(s),$$

$$M_n = 48\pi e_b^2 N_c \int_{-\infty}^\infty \frac{dE}{(E + 2m_b)^{2n+3}} \left(B_1^2 - B_1 d_1 \frac{E}{3m_b} \right) \text{Im} G(0, 0, E)$$


Matching coefficient of the electromagnetic current at NLL




$$B_s \quad D_s^{(1)}$$




$$B_s \quad \alpha_{V_s} \quad D_{d,s}^{(2)}$$



$$B_s \quad \alpha_{V_s} \quad D_{1,s}^{(2)}$$



$$B_s \quad \alpha_{V_s} \quad D_{S^2,s}^{(2)}$$



$$B_s \quad \alpha_{V_s} \quad c_4 \quad \alpha_{V_s}$$

$$\nu_p \frac{d}{d\nu_p} B_s = -\frac{C_A C_f}{2} D_s^{(1)} - \frac{C_f^2}{4} \alpha_s \left\{ \alpha_s - \frac{4}{3} s(s+1) D_{S^2,s}^{(2)} - D_{d,s}^{(2)} + 4D_{1,s}^{(2)} \right\},$$

$$b_1(m) = 1 - 2C_f \frac{\alpha_s(m)}{\pi}, \quad b_0(m) = 1 + \left(\frac{\pi^2}{4} - 5 \right) \frac{C_f \alpha_s(m)}{2\pi}.$$

The solution reads (Pineda; Hoang-Stewart)

$$B_s(\nu_p) = b_s(m) + A_1 \frac{\alpha_s(m)}{w^{\beta_0}} \ln(w^{\beta_0}) + A_2 \alpha_s(m) [z^{\beta_0} - 1] + A_3 \alpha_s(m) [z^{\beta_0 - 2C_A} - 1] \\ + A_4 \alpha_s(m) [z^{\beta_0 - 13C_A/6} - 1] + A_5 \alpha_s(m) \ln(z^{\beta_0}),$$

where $\beta_0 = \frac{11}{3}C_A - \frac{4}{3}T_F n_f$, $z = \left[\frac{\alpha_s(\nu_p)}{\alpha_s(m)} \right]^{\frac{1}{\beta_0}}$ and $w = \left[\frac{\alpha_s(\nu_p^2/m)}{\alpha_s(\nu_p)} \right]^{\frac{1}{\beta_0}}$. The coefficients A_i read

$$A_1 = \frac{8\pi C_f}{3\beta_0^2} (C_A^2 + 2C_f^2 + 3C_f C_A),$$

$$A_2 = \frac{\pi C_f [3\beta_0(26C_A^2 + 19C_A C_f - 32C_f^2) - C_A(208C_A^2 + 651C_A C_f + 116C_f^2)]}{78\beta_0^2 C_A},$$

$$A_3 = -\frac{\pi C_f^2 [\beta_0(4s(s+1) - 3) + C_A(15 - 14s(s+1))]}{6(\beta_0 - 2C_A)^2},$$

$$A_4 = \frac{24\pi C_f^2 (3\beta_0 - 11C_A)(5C_A + 8C_f)}{13C_A(6\beta_0 - 13C_A)^2},$$

$$A_5 = \frac{-\pi C_f^2}{\beta_0^2 (6\beta_0 - 13C_A)(\beta_0 - 2C_A)} \left\{ C_A^2(-9C_A + 100C_f) \right. \\ \left. + \beta_0 C_A(-74C_f + C_A(42 - 13s(s+1))) + 6\beta_0^2(2C_f + C_A(-3 + s(s+1))) \right\}.$$

Leading (Czarnecki-Melnikov; Beneke-Signer-Smirnov) and subleading (Kniehl-Penin) logs correct.

Inclusive decays to leptons and photons at NLL (Pineda)

By setting $\nu_p \sim m\alpha_s$, $B_s(\nu_p)$ includes all the large logs at **NLL** order in any (inclusive enough) S-wave heavy-quarkonium production observable we can think of. For instance, the decays to e^+e^- and to **two photons** at **NLL** $O(\alpha^{1+n} \ln^n \alpha)$ order read

$$\begin{aligned} \Gamma(V_Q(nS) \rightarrow e^+e^-) &= 2 \left[\frac{\alpha_{em} Q}{M_{V_Q(nS)}} \right]^2 \left(\frac{m_Q C_f \alpha_s}{n} \right)^3 \{B_1(\nu_p)(1 + \delta\phi_n)\}^2 \\ &\simeq 2 \left[\frac{\alpha_{em} Q}{M_{V_Q(nS)}} \right]^2 \left(\frac{m_Q C_f \alpha_s}{n} \right)^3 \{1 + 2(B_1(\nu_p) - 1) + 2\delta\phi_n\} , \\ \Gamma(P_Q(nS) \rightarrow \gamma\gamma) &= 6 \left[\frac{\alpha_{em} Q^2}{M_{P_Q(nS)}} \right]^2 \left(\frac{m_Q C_f \alpha_s}{n} \right)^3 \{B_0(\nu_p)(1 + \delta\phi_n)\}^2 \\ &\simeq 6 \left[\frac{\alpha_{em} Q^2}{M_{P_Q(nS)}} \right]^2 \left(\frac{m_Q C_f \alpha_s}{n} \right)^3 \{1 + 2(B_0(\nu_p) - 1) + 2\delta\phi_n\} , \end{aligned}$$

where V and P stand for the vector and pseudoscalar heavy quarkonium, we have fixed $\nu_p = m_Q C_f \alpha_s / n$, $\alpha_s = \alpha_s(\nu_p)$, and $(\Psi_n(z) = \frac{d^n \ln \Gamma(z)}{dz^n})$ and $\Gamma(z)$ is the Euler Γ -function)

$$\delta\phi_n = \frac{\alpha_s}{\pi} \left[-C_A + \frac{\beta_0}{4} \left(\Psi_1(n+1) - 2n\Psi_2(n) + \frac{3}{2} + \gamma_E + \frac{2}{n} \right) \right] .$$

NONRELATIVISTIC EFFECTIVE FIELD THEORIES

AND

RENORMALONS

Question

$$M_{\Upsilon(1S)} = m_{\text{OS}}(1 + A_2\alpha_s^2 + A_3\alpha_s^3 + \dots)$$

What if $A_n \sim n!$? Bad convergence

Should we expect that?

Renormalons

They are a potential problem in effective field theories of QCD (OPE) where the matching coefficients can be computed in perturbation theory.

Examples:

OPE

NRQCD

HQET

pNRQCD

SCET

Renormalons appear as soon as we have **factorization** between different scales: They can deteriorate the convergence of the perturbative series in **QCD**.

Can one understand the **renormalon** within an **effective field theory/factorization** formalism?

Problems:

- 1) Fix the parameters of the **Standard Model**. Search for weakly sensitive to long distance physics observables. One wants to avoid spurious dependence on the renormalon.
- 2) **Meaningful** determination of non-perturbative parameters.

$$\mathcal{L} = \sum_n \frac{1}{m^n} c_n O_n$$

Matching coefficients suffer from renormalon ambiguities that cancel with the ones of the matrix elements in effective field theory calculations.

$$c(\nu) = \bar{c} + \sum_{n=0}^{\infty} c_n \alpha_s^{n+1}.$$

Its Borel transform would be

$$B[c](t) \equiv \sum_{n=0}^{\infty} c_n \frac{t^n}{n!},$$

and c is written in terms of its Borel transform as

$$c = \bar{c} + \int_0^{\infty} dt e^{-t/\alpha_s} B[c](t).$$

The ambiguities in the matching coefficient ($c_n \sim n!$) reflects in poles in the Borel transform. If we take the one closest to the origin,

$$\delta B[c](t) \sim \frac{1}{a-t},$$

where a is a positive number, it sets up the maximal accuracy with which one can obtain the matching coefficients from a perturbative calculation, which is (roughly) of the order of

$$\delta c \sim r_{n^*} \alpha_s^{n^*},$$

where $n^* \sim \frac{a}{\alpha_s}$. Moreover, the fact that a is positive means that, even after Borel resummation, c suffers from a non-perturbative ambiguity of order

$$\delta c \sim (\Lambda_{QCD})^{\frac{a\beta_0}{2\pi}} .$$




Figure 1: *Symbolic relation between observables through the determination of the matching coefficients of the effective field theory.*

Examples

$$M_B = m_{\text{OS}} + \bar{\Lambda} + \mathcal{O}(1/m_{\text{OS}})$$

M_B is renormalon free. **Problem.** Therefore m_{OS} suffers from renormalon ambiguities:

$$m_{\text{OS}} = m_{\overline{\text{MS}}}(1 + B_1\alpha_s + B_2\alpha_s^2 + \dots)$$

with $B_n \sim n!$. In other words

$$\delta_{np}^{(\text{pert.})} m_{\text{OS}} = \delta_{np}^{(\text{pert.})} m_{\overline{\text{MS}}}(1 + B_1\alpha_s + B_2\alpha_s^2 + \dots) \sim \Lambda_{QCD}!$$

On the other hand

$$M_{\Upsilon(1S)} = m_{\text{OS}}(1 + A_2\alpha_s^2 + A_3\alpha_s^3 + \dots) + \mathcal{O}\left(\frac{\Lambda_{QCD}^3}{(m_{\text{OS}}\alpha_s)^2}\right).$$

$M_{\Upsilon(1S)}$ is renormalon free. Therefore, the perturbative series suffers from renormalon ambiguities: $A_n \sim n!$

$$\delta_{np}^{(\text{pert.})} M_{\Upsilon(1S)} = \delta_{np}^{(\text{pert.})} m_{\text{OS}}(1 + A_2\alpha_s^2 + A_3\alpha_s^3 + \dots) \sim \Lambda_{QCD}.$$

Physics. Computations to **n-loops** produce small scales: me^{-n} . From the effective field theory point of view these scales should be in the effective field theory instead that in the matching coefficients.

Proposal: to subtract the renormalon from the matching coefficients.

OS mass

$$m_{\text{OS}} = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n \alpha_s^{n+1},$$

The behavior of the perturbative expansion at large orders is dictated by the closest singularity to the origin of its Borel transform ($u = \frac{\beta_0 t}{4\pi}$).

$$B[m_{\text{OS}}](t(u)) = N_m \nu \frac{1}{(1-2u)^{1+b}} (1 + c_1(1-2u) + c_2(1-2u)^2 + \dots) + (\text{analytic term}),$$

Next renormalon at $u = 1$.

$$r_n \stackrel{n \rightarrow \infty}{\sim} N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

$$b = \frac{\beta_1}{2\beta_0^2}, \quad c_1 = \frac{1}{4b\beta_0^3} \left(\frac{\beta_1^2}{\beta_0} - \beta_2 \right), \quad \dots$$

Determination of N_m

$$\begin{aligned} D_m(u) &= \sum_{n=0}^{\infty} D_m^{(n)} u^n = (1-2u)^{1+b} B[m_{\text{OS}}](t(u)) \\ &= N_m \nu (1 + c_1(1-2u) + c_2(1-2u)^2 + \dots) + (1-2u)^{1+b} (\text{analytic term}). \end{aligned}$$

$$N_m \nu = D_m(u = 1/2).$$

$$\begin{aligned} N_m &= 0.4244 + 0.1379 + 0.0127 = 0.5750 & (n_f = 3) \\ &= 0.4244 + 0.1275 + 0.0004 = 0.5523 & (n_f = 4) \\ &= 0.4244 + 0.1199 - 0.0208 = 0.5235 & (n_f = 5) \end{aligned}$$




Figure 2: Plots of the exact (r_n^{ex}) and asymptotic (r_n^{as}) value of $r_n(\nu)$ at different orders in perturbation theory as a function of $\nu/m_{\overline{\text{MS}}}$. The scale dependence of r_3^{ex} is known exactly. The constant term has been fixed using renormalon dominance.

$$r_n \stackrel{n \rightarrow \infty}{\sim} m_{\overline{\text{MS}}} \left(\frac{\beta_0}{2\pi} \right)^n n! N_m \sum_{s=0}^n \frac{\ln^s[\nu/m_{\overline{\text{MS}}}]}{s!},$$

$$r_n \stackrel{n \rightarrow \infty}{\equiv} N_m \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right).$$

Renormalon subtracted matching and power counting

Effective field theory with renormalon free parameters but preserving the power counting rules.

The renormalon is associated to the non-analytic behavior in $1-2u$. These terms also exist in the effective theory. **Procedure:** to explicitly subtract them from the matching coefficients (the mass).

$$B[m_{\text{RS}}] \equiv B[m_{\text{OS}}] - N_m \nu_f \frac{1}{(1-2u)^{1+b}} (1 + c_1(1-2u) + c_2(1-2u)^2 + \dots),$$

$$m_{\text{RS}}(\nu_f) = m_{\text{OS}} - \sum_{n=0}^{\infty} N_m \nu_f \left(\frac{\beta_0}{2\pi}\right)^n \alpha_s^{n+1}(\nu_f) \sum_{k=0}^{\infty} c_k \frac{\Gamma(n+1+b-k)}{\Gamma(1+b-k)}.$$

Expansion in $\alpha_s(\nu)$

$$m_{\text{RS}}(\nu_f) = m_{\overline{\text{MS}}} + \sum_{n=0}^{\infty} r_n^{\text{RS}} \alpha_s^{n+1},$$

where $r_n^{\text{RS}} = r_n^{\text{RS}}(m_{\overline{\text{MS}}}, \nu, \nu_f)$. They are the ones expected to be of natural size. We now do not lose accuracy if we first obtain m_{RS} and later on $m_{\overline{\text{MS}}}$.

Different scheme

$$B[m_{\text{RS}'}] \equiv B[m_{\text{RS}}] + N_m \nu_f (1 + c_1 + c_2 + \dots).$$

Check of convergence improvement

Masses	$O(\alpha_s)$	$O(\alpha_s^2)$	$O(\alpha_s^3)$	$O(\alpha_s^4)$	total
m_{OS}	401	199	144	147	5 102
m_{RS}	111	50	17	7	4 395
$m_{RS'}$	401	114	38	15	4.778
m_{PS}	210	80	42	---	4 542
$m_{1S}^{(static)}$	102	50	19	8	4 389
m_{RS}	256	95	40	21	4 622
$m_{RS'}$	401	157	74	41	4.882
m_{PS}	306	120	67	---	4.703
$m_{1S}^{(static)}$	251	94	41	22	4 619

Table 1: Contributions at various orders in α_s for different mass definitions for the bottom quark case, either with $\nu_f = 1/r = 2$ GeV (middle panel) or with $\nu_f = 1/r = 1$ GeV (lower panel). The results are displayed in MeV. For the $O(\alpha_s^4)$ results, the estimate from Table ?? has been used. The other parameters have been fixed to the values $m_{\overline{MS}}(m_{\overline{MS}}) = 4.21$ GeV, $\nu = m_{\overline{MS}}(m_{\overline{MS}})$ and $n_f = 4$.

$$m_{1S}^{(static)} \equiv m_{OS} + \frac{V(r)}{2} = m_{\overline{MS}} + \left(r_0 - \frac{C_f}{2r} \right) \alpha_s + \dots$$

HQET

$$\mathcal{L} = \bar{h} (iD_0 - \delta m_{RS}) h + O\left(\frac{1}{m_{RS}}\right),$$

where $\delta m_{RS} = m_{OS} - m_{RS}$ and similarly for the **NRQCD** Lagrangian.

Weakly sensitive to long distance physics observable

$$\langle M_B \rangle - \langle M_D \rangle = m_{b,RS} - m_{c,RS} + \lambda_1 \left(\frac{1}{2m_{b,RS}} - \frac{1}{2m_{c,RS}} \right) + O(1/m_{RS}^2).$$

pNRQCD. If $\Lambda_{QCD} \ll m\alpha_s$

$$V_{s,RS(RS')}^{(0)}(\nu_f) = V_s^{(0)} + 2\delta m_{RS(RS')},$$

Check of convergence improvement

Potentials	$O(\alpha_s)$	$O(\alpha_s^2)$	$O(\alpha_s^3)$	$O(\alpha_s^4)$	total
$V_s^{(0)}$	-910	-306	-302	-383	-1 902
$V_{s,RS}^{(0)}$	-205	3	-2	-3	-208
$V_{s,RS'}^{(0)}$	-910	-54	-14	-6	-984
$V_{s,PS}^{(0)}$	-446	-42	-25	---	-513
$V_{s,RS}^{(0)}$	-558	-63	-41	-26	-687
$V_{s,RS'}^{(0)}$	-910	-180	-95	-54	-1 239
$V_{s,PS}^{(0)}$	-678	-116	-75	---	-869

Table 2: Contributions at various orders in α_s for different singlet static potential definitions for some typical scales in the Υ system, either with $\nu_f = 2$ GeV (middle panel) or with $\nu_f = 1$ GeV (lower panel). The results are displayed in MeV. For the $O(\alpha_s^4)$ results, the estimate from Table ?? has been used. The other parameters have been fixed to the values $\nu = 1/r = 2.5$ GeV and $n_f = 4$.

pNRQCD Lagrangian

$$\begin{aligned} \mathcal{L}^{(0)} = & \text{Tr} \left\{ S^\dagger \left(i\partial_0 - \frac{\mathbf{p}^2}{m_{RS}} + \sum_n \frac{V_{s,RS}^{(n)}(\mathbf{x})}{m_{RS}^n} \right) S + O^\dagger \left(iD_0 - \frac{\mathbf{p}^2}{m_{RS}} + \sum_n \frac{V_{o,RS}^{(n)}(\mathbf{x})}{m_{RS}^n} \right) O \right\} \\ & + gV_A(\mathbf{x}) \text{Tr} \left\{ O^\dagger \mathbf{x} \cdot \mathbf{E} S + S^\dagger \mathbf{x} \cdot \mathbf{E} O \right\} + g \frac{V_B(\mathbf{x})}{2} \text{Tr} \left\{ O^\dagger \mathbf{x} \cdot \mathbf{E} O + O^\dagger O \mathbf{x} \cdot \mathbf{E} \right\}, \end{aligned}$$

Weakly sensitive to long distance physics observable

$$M_{nlj} = 2m_{RS} + \sum_{m=2}^{\infty} A_{nlj}^{m,RS}(\nu_{us}) \alpha_s^m + \delta M_{nlj}^{US}(\nu_{us}).$$

The static potential

$$V_s^{(0)}(r; \nu_{us}) = \sum_{n=0}^{\infty} V_{s,n}^{(0)} \alpha_s^{n+1},$$

$2m_{OS} + V_s^{(0)}$ (not $2m_{OS} + V_o^{(0)}$) can be understood as an observable up to $O(r^2 \Lambda_{QCD}^3, \Lambda_{QCD}^2/m)$ renormalon (and/or non-perturbative) contributions.

We can use our knowledge of the asymptotic behavior of m_{OS} .

$$V_{s,n}^{(0)} \stackrel{n \rightarrow \infty}{\equiv} N_V \nu \left(\frac{\beta_0}{2\pi} \right)^n \frac{\Gamma(n+1+b)}{\Gamma(1+b)} \left(1 + \frac{b}{(n+b)} c_1 + \frac{b(b-1)}{(n+b)(n+b-1)} c_2 + \dots \right)$$

$$2N_m + N_V = 0$$


$$\begin{aligned} D_V(u) &= \sum_{n=0}^{\infty} D_V^{(n)} u^n = (1-2u)^{1+b} B[V_s^{(0)}](t(u)) \\ &= N_V \nu (1 + c_1(1-2u) + c_2(1-2u)^2 + \dots) + (1-2u)^{1+b} (\text{analytic term}). \end{aligned}$$

Next (IR) renormalon at $u = 3/2$.

$$\begin{aligned} N_V &= -1.333 + 0.572 - 0.345 = -1.107 \quad (n_f = 3) \\ &= -1.333 + 0.585 - 0.329 = -1.077 \quad (n_f = 4) \\ &= -1.333 + 0.587 - 0.295 = -1.042 \quad (n_f = 5). \end{aligned}$$

$$2 \frac{2N_m + N_V}{2N_m - N_V} = \begin{cases} 0.038 & , n_f = 3 \\ 0.025 & , n_f = 4 \\ 0.005 & , n_f = 5. \end{cases}$$

Bottom $\overline{\text{MS}}$ quark mass determination



Dependence on the parameters for the **RS** scheme: $\nu = 2.5_{-1}^{+1.5}$ GeV,
 $\nu_f = 2 \pm 1$ GeV, $\alpha_s(M_z) = 0.118 \pm 0.003$ and $N_m = 0.552 \pm 0.0552$


$$m_{b,\text{RS}}(2 \text{ GeV}) = 4387_{+28}^{+2} (\nu)_{+7}^{-5} (\nu_f)_{+16}^{-16} (\alpha_s)_{+68}^{-68} (N_m) \text{ MeV};$$

$$m_{b,\overline{\text{MS}}}(m_{b,\overline{\text{MS}}}) = 4203_{+25}^{+2} (\nu)_{+6}^{-5} (\nu_f)_{+27}^{-28} (\alpha_s)_{+10}^{-10} (N_m) \text{ MeV}.$$

Convergence. In the **RS** scheme

$$M_{\Upsilon(1S)} = 8774 + 559 + 120 + 7 \text{ MeV}.$$

NNLO(st. pot.) $\sim +62$ MeV. NNLO(rel.) ~ -55 MeV.



For the **RS'** scheme, we obtain the result

$$m_{b,\text{RS}'}(2 \text{ GeV}) = 4\,782_{+31}^{-08}(\nu)_{+3}^{-7}(\nu_f)_{-12}^{+15}(\alpha_s)_{+28}^{-28}(N_m) \text{ MeV};$$

$$m_{b,\overline{\text{MS}}}(m_{b,\overline{\text{MS}}}) = 4\,214_{+28}^{-08}(\nu)_{+3}^{-6}(\nu_f)_{+25}^{-25}(\alpha_s)_{+9}^{-9}(N_m) \text{ MeV}.$$

Convergence. In the **RS'** scheme

$$M_{\Upsilon(1S)} = 9\,564 - 158 + 56 - 2 \text{ MeV}.$$

NNLO(st. pot.) $\sim +45 \text{ MeV}$. **NNLO(rel.)** $\sim -47 \text{ MeV}$.

The static singlet potential

The introduction of renormalons allows to obtain agreement between lattice simulations and perturbation theory.

$$E_s = 2m_{\text{OS}} + V_{s,\text{OS}} + \mathcal{O}(r^2)$$

$$E_s = 2m_{\text{RS}}(\nu_f) + V_{s,\text{RS}}(\nu_f) + \mathcal{O}(r^2)$$




Figure 3: Plot of $V_{\text{OS}}(r)$ at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the leading single ultrasoft log (solid line). For the scale of $\alpha_s(\nu)$ $\nu = \text{constant}$. $\nu_{\text{us}} = 2.5 r_0^{-1}$.




Figure 4: Plot of $V_{\text{Os}}(r)$ at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the RG expression for the ultrasoft logs (solid line). For the scale of $\alpha_s(\nu)$, we set $\nu = 1/r$. $\nu_{us} = 2.5 r_0^{-1}$.




Figure 5: Plot of $V_{OS}(r) - V_{OS}(r') + E_{latt.}(r')$ versus r at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the RG expression for the ultrasoft logs (solid line) compared with the lattice simulations of Necco and Sommer. For the scale of $\alpha_s(\nu)$, we set $\nu = 1/r$. $\nu_{us} = 2.5 r_0^{-1}$.




Figure 6: Plot of $V_{\text{RS}}(r) - V_{\text{RS}}(r') + E_{\text{latt.}}(r')$ versus r at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the leading single ultrasoft log (solid line) compared with the lattice simulations of Necco and Sommer. For the scale of $\alpha_s(\nu)$ $\nu = \text{constant}$. $\nu_{\text{us}} = 2.5 r_0^{-1}$.




Figure 7: Plot of $V_{\text{RS}}(r) - V_{\text{RS}}(r') + E_{\text{latt.}}(r')$ versus r at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the RG expression for the ultrasoft logs (solid line) compared with the lattice simulations of Necco and Sommer. For the scale of $\alpha_s(\nu)$, we set $\nu = 1/r$. $\nu_f = \nu_{\text{us}} = 2.5 r_0^{-1}$.




Figure 8: Plot of $V_{\text{RS}}(r) - V_{\text{RS}}(r') + E_{\text{latt.}}(r')$ versus r at tree (dashed line), one-loop (dash-dotted line), two-loops (dotted line) and three loops (estimate) plus the leading single ultrasoft log (solid line) compared with the lattice simulations of Necco and Sommer. For the scale of $\alpha_s(\nu)$ $\nu = \text{constant}$. $\nu_{\text{us}} = 2.5 r_0^{-1}$.




Figure 9: Plot of $V_{\text{Os}}(r) - V_{\text{Os}}(r') + E_{\text{latt.}}(r') + \sigma(r - r')$ versus r at three loops (estimate) with the leading ultrasoft log compared with the lattice simulations of Necco and Sommer. For the scale of $\alpha_s(\nu)$, we set $\nu = \text{constant}$. $\sigma = 1.35 r_0^{-2}$ and $\nu_{\text{us}} = 2.5 r_0^{-1}$.

**Constraint on the size of nonperturbative effects for heavy quarkonium.
No linear non-perturbative potential at short distances.**