

**Nonlinear dynamics**  
**of spin and charge**  
**in the spin-Calogero model**



by

***Fabio Franchini***



Together with:

***A.G. Abanov,***  
***M. Kulkarni***

**Phys. Rev. B 80, 165105 (2009)**  
**(arXiv: 0904.3762)**

**Nucl. Phys. B 825, 320 (2010)**  
**(arXiv: 0908.2652)**

# Universality in 1-D systems

- In 1-D: no Fermi Liquid, but **Luttinger Liquid**
- Collective nature of excitations: **hydrodynamics**
- Low-Energy approximation: **phonons**
- Linear dispersion relation (**Lorentz invariance**)

⇒ **Bosonization & CFT**

# Limits of Luttinger Liquid

- Non-relativistic systems  $\rightarrow$  non-linearities
- Non-equilibrium  $\rightarrow$  non-linearities
- Linear theory  $\rightarrow$  no solitons
- Linear spectrum implies **spin-charge separation**: particles decouple into spinons and holons, but curvature **couples** spin and charge dynamics
- ...

# Outline


- Motivation (done) & introduction
- Hydrodynamics from Bethe Ansatz  
(Free Fermions example)
- Spin Calogero-Sutherland model  
(and its gradient-less hydrodynamics)
- Applications: Spin-Charge dynamics  
Emptiness Formation Probability
- Connection to the Haldane-Shastry model
- Conclusions

# Bosonization

- Bosonization formula:  $\psi(x) = \frac{1}{\sqrt{2\pi}} : e^{i\sqrt{4\pi}\phi(x)} :$
- Free Fermions Hamiltonian:

$$\psi^\dagger(x)\partial^2\psi(x) = \frac{1}{3\sqrt{4\pi}}\partial^3\phi + i\partial\phi(\partial\phi)^2 + \frac{3}{\sqrt{4\pi}}(\partial\phi)^3$$

cubic potential  $\rightarrow$  **unstable vacuum**



$\Rightarrow$  need correct normal ordering prescription

# Luttinger Liquid

- Apply bosonization after **linearizing** the spectrum:

$$-\psi^\dagger \partial_x^2 \psi \simeq - \sum_{\alpha=L,R} \psi_\alpha^\dagger (\partial_x \pm ik_F)^2 \psi_\alpha + \dots$$

- In low-energy approx, standard prescriptions work
- Bosonization = **Linearized Hydrodynamics**

$$\rho(x) = \psi^\dagger(x)\psi(x) = \frac{1}{\sqrt{\pi}} \partial_x \phi(x)$$

$$[\partial_x \phi(x), \Pi(y)] = [\rho(x), v(y)] - i\hbar \delta'(x-y)$$

# Non-linear effects

- Realization of 1-D systems
  - experimentally relevant
  - (Quantum quenches; Non-equilibrium dynamics ...)
- Several theoretical approaches toward non-linear effects (**Universality?**)
- Solitons?
- So far, not much effort toward spin-charge dynamics

# Hydrodynamics construction

Integrable system

$$\tau(k)$$

Free Fermions

$$\tau(k) = \frac{1}{2\pi}$$

Particle density:

$$\rho = \int_{k_L}^{k_R} \tau(k) dk$$

$$\rho = \int_{k_L}^{k_R} \frac{dk}{2\pi} = \frac{k_R - k_L}{2\pi} \quad v = \frac{k_R + k_L}{2}$$

Momentum density:

$$j = \int_{k_L}^{k_R} \frac{dk}{2\pi} k \tau(k) = \frac{k_R^2 - k_L^2}{4\pi} \quad j = \rho v = \int_{k_L}^{k_R} \frac{dk}{2\pi} k = \frac{k_R^2 - k_L^2}{4\pi}$$

Energy density:

$$\frac{E}{L} = \int_{k_L}^{k_R} \frac{dk}{2\pi} \frac{k^2}{2} \tau(k) = \frac{k_R^3 - k_L^3}{12\pi} \quad \frac{E}{L} = \int_{k_L}^{k_R} \frac{dk}{2\pi} \frac{k^2}{2} \tau(k) = \frac{k_R^3 - k_L^3}{12\pi}$$



# Hydrodynamics construction

Free Fermions

$$\tau(k) = \frac{1}{2\pi}$$

We let the parameters have a slow space-dependence!

Particle density:  $\rho(x) = \frac{k_R(x) - k_L(x)}{2\pi}$


Momentum density:  $j(x) = \frac{k_R^2(x) - k_L^2(x)}{4\pi} = \rho(x)v(x)$

Energy density:  $\frac{E}{L} = \frac{\rho(x) v^2(x)}{2} + \frac{\pi^2}{6} \rho^3(x) = \mathcal{H}(x)$



- In general, dynamics induces change in  $\tau(k)$   
→ corrections due to non-equilibrium

# Recapping...

Equilibrium  
Free Fermions



Non-Equilibrium  
Free Fermions



Space-  
Dependent  
Configuration

# Free Fermions Hydrodynamics

$$H = \int dx \mathcal{H}(x) = \int dx \left[ \frac{\rho(x) v^2(x)}{2} + \frac{\hbar^2 \pi^2}{6} \rho^3(x) \right]$$

Dynamics from commutation relations:  
(from microscopical analysis)

$$[\rho(x), v(y)] = -i\hbar\delta'(x - y)$$

Continuity  
Equation:

$$\dot{\rho} = [H, \rho] = -\partial_x (\rho v)$$

Euler  
Equation:

$$\dot{v} = [H, v] = -\partial_x \left( \frac{v^2}{2} + \frac{\hbar^2 \pi^2}{2} \rho^2 \right)$$

# Free Fermions Hydrodynamics

$$H = \int dx \left[ \frac{\rho(x) v^2(x)}{2} + \frac{\hbar^2 \pi^2}{6} \rho^3(x) \right] = \int dx \hbar^2 \frac{k_R^3(x) - k_L^3(x)}{12\pi}$$

Note that:  $[\rho(x), v(y)] = -i\hbar\delta'(x - y)$




$$[k_L(x), k_L(y)] = -[k_R(x), k_R(y)] = 2\pi i\delta'(x - y)$$

Riemann-Hopf

Equation:  $\dot{k}_{R,L} + \hbar k_{R,L} \partial_x k_{R,L} = 0$

Left and Right Fermi points  
evolve independently!



# Free Fermions Hydrodynamics

Two remarks:

1. In principle, gradient corrections from interaction.

For Free Fermions, this is **exact!**

$$H = \int dx \left[ \frac{\rho v^2}{2} + \frac{\hbar^2 \pi^2}{6} \rho^3 \right] = \frac{\hbar^2}{12\pi} \int dx [k_R^3 - k_L^3]$$

2. Same result can be derived by **conventional bosonization without** linearization:

$$\Psi_{L,R}(x) =: e^{i\sqrt{4\pi}\phi_{L,R}(x)} :$$

$$H = \frac{\hbar^2}{12\pi} \int dx \left[ (\partial_x \phi_R)^3 - (\partial_x \phi_L)^3 \right]$$

# Free Fermions with Spin

- Just add the theory for each species:

$$H = \int dx \left\{ \frac{1}{2} \rho_{\uparrow} v_{\uparrow}^2 + \frac{1}{2} \rho_{\downarrow} v_{\downarrow}^2 + \frac{\pi^2 \hbar^2}{6} (\rho_{\uparrow}^3 + \rho_{\downarrow}^3) \right\}$$

- Expanding around ( $\rho_0 = k_F/\pi$ ,  $v_0 = 0$ ):

$$\begin{aligned} H &\approx \frac{\rho_0}{2} \int dx (v_{\uparrow}^2 + \pi^2 \hbar^2 \delta \rho_{\uparrow}^2 + v_{\downarrow}^2 + \pi^2 \hbar^2 \delta \rho_{\downarrow}^2) \\ &\approx \frac{\rho_0}{4} \hbar^2 \sum_{\alpha=\uparrow,\downarrow} \int dx [(\partial_x \phi_{R,\alpha})^2 + (\partial_x \phi_{L,\alpha})^2] \end{aligned}$$

$\Rightarrow$  traditional **bosonization!**

# Bosonization of spinful Free Fermions

$$H \approx \frac{\rho_0}{4} \hbar^2 \sum_{\alpha=c,s} \int dx [(\partial_x \phi_{R,\alpha})^2 + (\partial_x \phi_{L,\alpha})^2]$$

- No true spin-charge separation (all excitations have same velocity)
- Peculiarity of FF (and of Calogero-Sutherland systems)

- Riemann-Hopf equation
 
$$\dot{k}_{\alpha,\chi} + \hbar k_{\alpha,\chi} \partial_x k_{\alpha,\chi} = 0$$

$$\downarrow k_{\alpha,\chi} \rightarrow \pi \rho_0 + k_{\alpha,\chi}$$

$$\dot{k}_{\alpha,\chi} + \hbar \pi \rho_0 \partial_x k_{\alpha,\chi} = 0$$

becomes wave equation

(  $\alpha=c,s; \chi = R,L$  )

# Semi-Classical Limit

$$H = \int dx \left\{ \frac{1}{2} \rho_{\uparrow} v_{\uparrow}^2 + \frac{1}{2} \rho_{\downarrow} v_{\downarrow}^2 + \frac{\pi^2 \hbar^2}{6} (\rho_{\uparrow}^3 + \rho_{\downarrow}^3) \right\}$$

- In the classical limit, only velocity terms survive
- Semiclassical limit:  $\rho \sim v/\hbar$   
 $\Rightarrow t \rightarrow t/\hbar$  and  $v \rightarrow \hbar v$

$$H = \int dx \left\{ \frac{1}{2} \rho_{\uparrow} v_{\uparrow}^2 + \frac{1}{2} \rho_{\downarrow} v_{\downarrow}^2 + \frac{\pi^2}{6} (\rho_{\uparrow}^3 + \rho_{\downarrow}^3) \right\}$$

- Commutation relations  $\rightarrow$  Poisson Brackets

$$\{\rho_{\alpha}(x), v_{\beta}(y)\} = \delta_{\alpha\beta} \delta'(x - y)$$



# Non-linear Hydrodynamic

- Cannot apply bosonization formula directly  
(un-renormalizable divergencies)
- We look at integrable models: can sum up interactions
- Collective field theory description:  $\rho_{c,s}(x,t)$  ,  $v_{c,s}(x,t)$   
 $\Rightarrow$  Hydrodynamics approach
- **Calogero-Moser Model (sCM)**-kind of interaction

# Calogero-Sutherland model

$$H \equiv -\frac{\hbar^2}{2} \sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \frac{\hbar^2}{2} \sum_{j \neq l} \frac{\lambda(\lambda - 1)}{(x_j - x_l)^2}$$

- Hydrodynamic description well understood

(Abanov, Wiegmann, Andric, Bardek...)

- Semi-classical limit

→ non-linear integrable classical equation:

**deformed Benjamin-Ono** equation (on the double)

$$\partial_t u + \partial \left[ \frac{u^2}{2} + i\alpha_0 \partial (u_+ - u_-) \right] = 0$$

$$2\alpha_0 = \lambda^{1/2} - \lambda^{-1/2}$$

# Spin Calogero-Sutherland model

$$H \equiv -\frac{\hbar^2}{2} \sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \frac{\hbar^2}{2} \sum_{j \neq l} \frac{\lambda(\lambda \pm \mathbf{P}_{jl})}{(x_j - x_l)^2}$$

- $\mathbf{P}_{jl}$  particle-exchange operator
- SU(2) version of the traditional CS model ( $\mathbf{P}_{jl} = \pm 1$  for a ferromagnetic state)
- $\lambda \rightarrow \infty$  : Haldane-Shastry spin chain

Bosons	$\longrightarrow$	$\left\{ \begin{array}{l} + \Rightarrow \text{Anti-ferromagnetic,} \\ - \Rightarrow \text{Ferromagnetic,} \end{array} \right.$
Fermions	$\longrightarrow$	$\left\{ \begin{array}{l} + \Rightarrow \text{Ferromagnetic,} \\ - \Rightarrow \text{Anti-ferromagnetic.} \end{array} \right.$

# Anti-Ferromagnetic Fermions

$$H = -\frac{\hbar^2}{2} \sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \frac{\hbar^2}{2} \left(\frac{\pi}{L}\right)^2 \sum_{j \neq l} \frac{\lambda(\lambda - \mathbf{P}_{jl})}{\sin^2 \frac{\pi}{L} (x_j - x_l)}$$

- AF Ground state
- We chose a fermionic Hilbert space:

Periodic boundary conditions

$$\psi_{GS} = \prod_{j < l} \left| \sin \frac{\pi}{L} (x_j - x_l) \right|^\lambda \prod_{j < l} \left[ \sin \frac{\pi}{L} (x_j - x_l) \right]^{\delta(\sigma_j, \sigma_l)} e^{i \frac{\pi}{2} \text{sgn}(\sigma_j - \sigma_l)}$$

"anyonic" (Laughlin-type) wave-function

# Hydrodynamic Description

- Collective field theory description:  $\rho_{c,s}(\mathbf{x},t)$  ,  $v_{c,s}(\mathbf{x},t)$
- Our is a **heuristic construction** based on **Bethe Ansatz** Solution ( $\rightarrow$  valid for small gradients!)
- Calogero-Sutherland model:  $\theta(k) = \pi\lambda \operatorname{sgn}(k)$   
 $\rightarrow$  **dynamical** phase like a **statistical** phase
- Distribution  $\pi(k)$  of quasi-momenta **piece-wise constant** (peculiar to Calogero interaction)

Sutherland & Shastry (1993)

Kato & Kuramoto (1995)

# (Asymptotic) Bethe Ansatz Solution

- Constant scattering phase:  $\theta(k) = \pi\lambda \operatorname{sgn}(k)$
- States defined by set of integer numbers  $\mathbf{\kappa}_{\uparrow,\downarrow}$   
for spin up/down particles
- **Hydrodynamic** distribution  $v(\mathbf{\kappa})$ : 
$$\begin{cases} \kappa_{\uparrow} = \kappa_{L\uparrow}, \dots, \kappa_{R\uparrow} \\ \kappa_{\downarrow} = \kappa_{L\downarrow}, \dots, \kappa_{R\downarrow} \end{cases}$$
- Proceeding as for FF, we make the identification:

$$v_{\alpha} \pm \pi\rho_{\alpha} \equiv \frac{2\pi}{L} \kappa_{(R,L);\alpha}$$


# The 3 regimes

$$\epsilon = \sum_{\kappa=-\infty}^{+\infty} \kappa^2 \nu(\kappa) + \frac{\lambda}{2} \sum_{\kappa, \kappa'} |\kappa - \kappa'| \nu(\kappa) \nu(\kappa')$$

Sutherland & Shastry (1993)

Kato & Kuramoto (1995)

- Because of the absolute value, we identify 3 regimes:




$$v_{\alpha} \pm \pi \rho_{\alpha} \equiv \frac{2\pi}{L} \kappa_{(R,L); \alpha}$$

Partial Overlap (PO)

No Overlap (NO)

x2: Exchanging  $\uparrow \leftrightarrow \downarrow$

# The 3 regimes



Complete Overlap	→	$ v_s  < \frac{\pi}{2}  \rho_s $
Partial Overlap	→	$\frac{\pi}{2}  \rho_s  <  v_s  < \frac{\pi}{2} \rho_c$
No Overlap	→	$\frac{\pi}{2} \rho_c <  v_s $

- CO & PO: small deviation from **AFM Ground state**
- Here, I'll concentrate **only** on the **CO regime**



# The CO regime

$$H_{\text{CO}} = \int dx \left\{ \frac{1}{2} \rho_c v_c^2 + \frac{\pi^2}{6} \left( \lambda + \frac{1}{2} \right)^2 \rho_c^3 + \rho_s v_c v_s \right. \\ \left. + \left[ \left( \lambda + \frac{1}{2} \right) \rho_c - \lambda \rho_s \right] v_s^2 + \frac{\pi^2}{4} \left( \lambda + \frac{1}{2} \right) \rho_c \rho_s^2 - \frac{\pi^2}{12} \lambda \rho_s^3 \right\}$$

$$\{\rho_\alpha(x), v_\beta(y)\} = \delta_{\alpha\beta} \delta'(x - y)$$

- Non-linear dynamics **couples spin & charge!**

# The CO regime

- Introduce the following linear combination of fields:

$$k_{R\uparrow,L\uparrow} = (v_{\uparrow} \pm \pi\rho_{\uparrow}) \pm \pi\lambda\rho_c,$$

$$k_{R\downarrow,L\downarrow} = (v_{\downarrow} \pm \pi\rho_{\downarrow}) \pm \pi\lambda\rho_c - \lambda(2v_s \pm \pi\rho_s)$$


- This transformation **decouples** the dynamics into 4 Riemann-Hopf equations:

$$H = \sum_{\alpha=\uparrow,\downarrow} \sum_{\chi=L,R} s_{\chi;\alpha} \int dx k_{\chi;\alpha}^3$$

$$u_t + uu_x = 0, \quad u = k_{R,L;\uparrow,\downarrow}$$


- These  $k$ 's are the BA **dressed "Fermi" momenta**

# The CO regime





$$u_t + uu_x = 0, \quad u = k_{R,L;\uparrow,\downarrow}$$

# Phase-Space picture for the 3 regimes



$$u_t + uu_x = 0, \quad u = k_{R,L;\uparrow,\downarrow}$$




Dressed "Fermi" momenta picture works for every regime

# A few words on the Riemann-Hopf Eq.

$$u_t + uu_x = 0$$

- Simplest non-linear equation
- Given an initial condition  $u(x,0) = u_0(x)$ , its solution is implicitly given by  $u = u_0(x-ut)$  (easy to handle **numerically**)
- Ill defined for long times (gradient catastrophe):



- Should be corrected by gradient terms like:

$$u_t + uu_x + u_{xxx} = 0,$$

$$u_t + uu_x + u_{xx}^H = 0 \dots$$

# And a few words on our solutions

$$u_t + uu_x = 0, \quad u = k_{R,L;\uparrow,\downarrow}$$

- We neglected gradient corrections from the start
- Decoupling of Fermi momenta into RH equations probably broken by gradient correction
- Our hydrodynamics valid for “small” times ( $\ll$  than gradient catastrophe)
- We require all gradients to be small compared to interparticle distance

# Spin singlet dynamics

- Consider initial condition:  $\rho_s, v_s = 0$
- Any configuration of charge sector will not perturb this spin singlet state:

$$\dot{\rho}_c = -\partial_x(\rho_c v_c)$$


$$\dot{\rho}_s = 0$$

$$\dot{v}_c = -\partial_x \left\{ \frac{v_c^2}{2} + \frac{\pi^2 \left(\lambda + \frac{1}{2}\right)^2 \rho_c^2}{2} \right\}$$

$$\dot{v}_s = 0$$

(Spinless Calogero-Sutherland with  $\lambda+1 \rightarrow \lambda + 1/2$ )


# Dynamics of a polarized center: FF



Initial condition at  $t = 0$ :

$$\rho_c = 1, v_c = 0,$$


$$v_s = 0, \rho_s = \frac{h}{1 + (x/a)^2}$$



- N.B.  $\lambda=0$ : Free Fermions!
- Essential non-linear dynamics



# Dynamics of a polarized center: sCM




Initial condition at  $t = 0$ :

$$\rho_c = 1, v_c = 0,$$


$$v_s = 0, \rho_s = \frac{h}{1 + (x/a)^2}$$

- Qualitatively similar behaviors for rescaled quantities ( $\tau = (\lambda + 1/2)t$ )
- Charge freezing



# The Haldane-Shastry model

$$H = -\frac{\hbar^2}{2} \sum_{j=1}^N \frac{\partial^2}{\partial x_j^2} + \frac{\hbar^2}{2} \left(\frac{\pi}{L}\right)^2 \sum_{j \neq l} \frac{\lambda(\lambda - \mathbf{P}_{jl})}{\sin^2 \frac{\pi}{L} (x_j - x_l)}$$

$\lambda \rightarrow \infty$   "freezing trick"  
(Polychronakos, 1993)

$$H_{\text{HSM}} = \frac{1}{2} \left(\frac{\pi}{N}\right)^2 \sum_{j < l} \frac{\mathbf{K}_{jl}}{\sin^2 \frac{\pi}{N} (j - l)} \quad \begin{array}{l} \text{(Haldane, 1988;} \\ \text{Shastry, 1988)} \end{array}$$

- Spectrum of HS equal to spinless CS with  $\lambda=2$ , but with high degeneracy (Yangian symmetry) (Haldane & Ha, 1992;  
Ha & Haldane, 1993)

# Connection to Haldane-Shastry model

Hydrodynamics of sCM from its Bethe Ansatz solution

$$H \simeq \int dx \left\{ \frac{\pi^2}{6} \mu^2 \rho_c^3 + \mu \left[ \rho_c v_s^2 - \rho_s v_s^2 + \frac{\pi^2 \rho_c \rho_s^2}{4} - \frac{\pi^2 \rho_s^3}{12} \right] + O(\mu^0) \right\}$$

$$\mu = \lambda + 1/2$$

$$H_{\text{HSM}} = \int dx \left\{ \rho_0 v_s^2 - \rho_s v_s^2 + \frac{\pi^2 \rho_0 \rho_s^2}{4} - \frac{\pi^2 \rho_s^3}{12} \right\}$$

Hydrodynamics of HSM from its Bethe Ansatz solution

$$\rho = \rho_{\downarrow} = \frac{\rho_0 - \rho_s}{2}, \quad v = -2v_s, \quad \rho_0 = 1$$

- Higher orders give corrections to freezing...

# Correlation functions

- So far: non-linear dynamics couples **spin & charge**
- Asymptotics of correlation functions easy from **field theory**
- 2-point correlation functions: **Luttinger Liquid** is sufficient
- For **extended** objects non-linear theory is **needed**
- To leading order, gradient-less theory is enough

# Emptiness Formation Probability

- It measures the probability  $P(\mathbf{R})$  that there are no particles for  $-\mathbf{R} < x < \mathbf{R}$
- Simplest correlator in **integrable models** (QISM)
- For sCM different EFPs:  $P_{\alpha}(\mathbf{R})$ ,  $\alpha = \uparrow, \downarrow, c, s \dots$
- Easy to calculate in instanton formalism
- Non-local correlation function
  - ⇒ linear bosonization **not sufficient**: full hydrodynamics

# Instanton Approach to EFP

- EFP as probability of rare fluctuation in imaginary time

$$P(R) \simeq e^{-\mathcal{S}[\phi_{\text{EFP}}]}$$

- Instanton: solution of equation of motion with b.c.'s

$$\rho_{\alpha}(\tau = 0, -R < x < R) = \bar{\rho}_{\alpha}, \quad \alpha = \uparrow, \downarrow$$

$$\rho_{\alpha}(x, \tau \rightarrow \infty) \rightarrow \rho_{0\alpha}, \quad v_{\alpha}(x, \tau \rightarrow \infty) \rightarrow 0$$

- $\bar{\rho}_{\alpha} = 0$ : Emptiness, otherwise **Depletion Formation Probability (DFP)**
- Gradient-less theory sufficient for leading order

# EFP/DFP for sCM

- Using our hydrodynamic description for sCM:

$$P(R) \simeq \exp \left\{ -\frac{\pi^2}{2} \left[ \left( \lambda + \frac{1}{2} \right) (\rho_{0c} - \bar{\rho}_c)^2 + \frac{1}{2} (\rho_{0s} - \bar{\rho}_s)^2 \right] R^2 \right\}$$

- Result factorizes: **effective spin-charge separation** in non-linear dynamics!
- Spin and charge sectors as **independent spin-less Calogero** fluids with couplings  $\lambda' = \lambda + 1/2$  and  $\lambda' = 2$
- Spin-charge separation true (at **Gaussian Order**) for other extended correlators: why?

# Conclusions

- Derived **gradient-less hydrodynamics** for **spin Calogero-Sutherland & Haldane Shastry** model
- Showed **freezing limit** and corrections
- Captured essentially **non-linear** spin-charge coupling
- EFP restores effective **spin-charge separation**

# Outlook

- Ferromagnetic Fermions, Bosons
- Gradient terms

**Thank you!**