


Determination of α_s from τ 's

Matthias Jamin



S. Bethke

For 0.6 % precision at M_Z need “only” ≈ 2 % at M_τ .

Consider the physical quantity R_τ : (Braaten, Narison, Pich 1992)

$$R_\tau \equiv \frac{\Gamma(\tau^- \rightarrow \text{hadrons} \nu_\tau(\gamma))}{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau(\gamma))} = 3.6280(94) . \text{ (HFAG 2012)}$$

R_τ is related to the QCD correlators $\Pi^{(1,0)}(z)$: ($z \equiv s/M_\tau^2$)

$$R_\tau = 12\pi \int_0^1 dz (1-z)^2 \left[(1+2z) \text{Im} \Pi^{(1)}(z) + \text{Im} \Pi^{(0)}(z) \right],$$

with the appropriate combinations

$$\Pi^{(J)}(z) = |V_{ud}|^2 \left[\Pi_{ud}^{V,J} + \Pi_{ud}^{A,J} \right] + |V_{us}|^2 \left[\Pi_{us}^{V,J} + \Pi_{us}^{A,J} \right].$$



Additional **exp** information can be inferred from the moments

$$R_{\tau}^w \equiv \int_0^1 dz w(z) \frac{dR_{\tau}}{dz} = R_{\tau,V}^w + R_{\tau,A}^w + R_{\tau,S}^w.$$

Theoretically, R_{τ}^w can be expressed as:

$$R_{\tau}^w = N_c S_{EW} \left\{ (|V_{ud}|^2 + |V_{us}|^2) [1 + \delta^{w(0)}] \right. \\ \left. + \sum_{D \geq 2} [|V_{ud}|^2 \delta_{ud}^{w(D)} + |V_{us}|^2 \delta_{us}^{w(D)}] \right\}.$$

$\delta_{ud}^{w(D)}$ and $\delta_{us}^{w(D)}$ are corrections in the Operator Product Expansion, the most important ones being $\sim m_s^2$ and $m_s \langle \bar{q}q \rangle$.



OPAL data can be updated with new branching fractions.

ALEPH data currently miss correlations from unfolding.

The **purely** perturbative contribution $\delta^{(0)}$ is plagued by differences for different RG-resummations. (FOPT vs CIPT.)

Using $\alpha_s(M_\tau) = 0.3186$, the numerical analysis results in:

$$a^1 \quad a^2 \quad a^3 \quad a^4 \quad a^5$$

$$\delta_{\text{FO}}^{(0)} = 0.101 + 0.054 + 0.027 + 0.013 (+0.006) = 0.196 \text{ (0.202)}$$

$$\delta_{\text{CI}}^{(0)} = 0.137 + 0.026 + 0.010 + 0.007 (+0.003) = 0.181 \text{ (0.185)}$$

Contour improved PT appears to be better convergent.

The difference between both approaches is 0.015 (0.017) !

This problematic entails a $\approx 6\%$ difference for $\alpha_s(M_\tau)$.


To proceed, realistic model $B[\widehat{D}](u)$: (Beneke, MJ 2008)

$$B[\widehat{D}](u) = B[\widehat{D}_1^{\text{UV}}](u) + B[\widehat{D}_2^{\text{IR}}](u) + B[\widehat{D}_3^{\text{IR}}](u) + d_0^{\text{PO}} + d_1^{\text{PO}} u,$$

where

$$B[\widehat{D}_p](u) = \frac{d_p}{(p \pm u)^{1+\gamma}} [1 + b_1(p \pm u) + b_2(p \pm u)^2].$$

- ☞ Our main model incorporates the leading UV pole ($u = -1$), as well as the two leading IR renormalons ($u = 2, 3$).
- ☞ It should reproduce the exactly known $c_{n,1}$, $n \leq 4$.
- ☞ For both UV and IR, the residues d_p are free while γ , $b_{1,2}$ depend on anomalous dimensions and β -coefficients.



$$c_{5,1} = 283, \quad \alpha_s(M_\tau) = 0.3186. \quad (\text{Beneke, MJ 2008})$$

In the OPE, close to the Minkowskian axis ($s > 0$), so-called Duality Violations (DV's) can appear.

They can be studied on the basis of a toy-model:

(Shifman et al. 1995-2000)

(Catà, Golterman, Peris 2005/2008)

$$\Pi_V(s) = -\psi\left(\frac{M_V^2 + u(s)}{\Lambda^2}\right) + \text{const.} .$$

where

$$u(s) = \Lambda^2 \left(\frac{-s}{\Lambda^2}\right)^\zeta \quad \text{and} \quad \zeta = 1 - \frac{a}{\pi N_c} .$$

The model is based on large- N_c QCD and Regge-theory.

$$M_V = 770 \text{ MeV}, \quad \Lambda = 1.2 \text{ GeV}, \quad a = 0.4 .$$


The OPE corresponds to the asymptotic expansion of the ψ -function for large s (large u).

$$\psi(z) \sim \ln z - \frac{1}{2z} - \sum_{n=1}^{\infty} \frac{B_{2n}}{2n z^{2n}}, \quad \operatorname{Re} z > 0.$$


In the Minkowskian region, an additional term arises:

$$- \pi [\cot(\pi z) \pm i], \quad \operatorname{Re} z < 0, \quad \operatorname{Im} z \gtrless 0.$$


Formally, this term is exponentially suppressed, but it is enhanced by the poles of the ψ -function.



$$z = 1.5 \cdot \exp(i\varphi) \quad (\text{MJ 2011})$$



ψ -function moment for $w(z) = 1$. (MJ 2011)



ψ -function moment for $w(z) = (1 - z)^2$.

In fits to experimental data, a model for DV's should be included.

The ψ -function model suggests an oscillating, decaying exponential, which can be chosen of the form:


$$\rho_{V/A}^{\text{DV}}(s) = \kappa_{V/A} e^{-\gamma_{V/A}s} \sin(\alpha_{V/A} + \beta_{V/A}s).$$

The fit quantities are the w -moments of the exp spectra.


$$R_{\tau,V/A}^w(s_0) \equiv \int_0^{s_0} ds w(s) \rho_{V/A}(s).$$

The cleanest moment turns out to be $w(s) = 1$.

Fitting combinations of several moments is complicated by very strong correlations.



(Boito, Golterman, MJ, Mahdavi, Maltman, Osborne, Peris 2012)



(Boito, Golterman, MJ, Mahdavi, Maltman, Osborne, Peris 2012)

- ☞ Presently, the most reliable value of α_s from τ 's including DV's comes from the trivial moment $w(s) = 1$.

$$\Rightarrow \alpha_s(M_\tau) = 0.325 \pm 0.016 \pm 0.007 \quad (\text{FOPT})$$

$$\Rightarrow \alpha_s(M_\tau) = 0.347 \pm 0.024 \pm 0.005 \quad (\text{CIPT})$$

- ☞ These values should be compared to the World Average (Bethke 2009): $\alpha_s(M_\tau) = 0.3186 \pm 0.0058$.
- ☞ Better data on exclusive and inclusive τ decay spectra would be very helpful to resolve theoretical issues.

- ☞ Presently, the most reliable value of α_s from τ 's including DV's comes from the trivial moment $w(s) = 1$.

$$\Rightarrow \quad \alpha_s(M_\tau) = 0.325 \pm 0.016 \pm 0.007 \quad (\text{FOPT})$$

$$\Rightarrow \quad \alpha_s(M_\tau) = 0.347 \pm 0.024 \pm 0.005 \quad (\text{CIPT})$$

- ☞ These values should be compared to the World Average (Bethke 2009): $\alpha_s(M_\tau) = 0.3186 \pm 0.0058$.
- ☞ Better data on exclusive and inclusive τ decay spectra would be very helpful to resolve theoretical issues.

Thank You for Your attention !