


Hunting RNA motifs

Paul Gardner

July 23, 2012

Classification problems

- ▶ What is this?
 - ▶ The second most abundant *M. tuberculosis* transcript
 - ▶ No hits to Rfam, no homologs have been identified, no known function




Rank	Reads	Gene
1	5,741K	SSU rRNA
2	689K	sRNA/RUF?
3	236K	RNaseP RNA
4	197K	LSU rRNA
5	119K	ATP synthase
6	119K	HSPX
7	115K	tRNA
8	103K	Secreted protein

Arnvig et al.
(2011) PLoS
Pathog.

Classification problems


- ▶ What about this?

- ▶ The seventh most abundant *N. gonorrhoeae* transcript
- ▶ No hits to Rfam, no homologs have been identified, no known function



Classification problems


- ▶ And this?
 - ▶ The ninth most abundant *C. difficile* transcript
 - ▶ No hits to Rfam, no homologs have been identified, no known function



Classification problems

- ▶ And this?

- ▶ The eleventh most abundant *C. difficile* transcript
- ▶ No hits to Rfam, no homologs have been identified, no known function



Rank	Reads	Gene
1	12K	RNaseP RNA
2	12K	6S RNA
3	12K	SRP RNA
4	11K	ldhA
5	10K	tmRNA
6	6K	spoVG
7	6K	Gly reductase
8	5K	slpA
9	4K	sRNA/RUF?
10	4K	hypothetical prot.
11	4K	sRNA/RUF?

Deakin, Lawley
et al. (2012)

Unpublished.

What is our next big challenge?

- ▶ Past:
 - ▶ How many non-coding RNA genes are there?
- ▶ Future:
 - ▶ Can we determine the functions, if any, for large sets of given RNAs?




Will a “periodic table of RNA” be useful?

- ▶ My aim is to build an analog to the Periodic Table for classifying RNA families and motifs, enabling researchers to predict function.

base	basepair
ANYA	GNRA
sarric1	sarric2
TRIT	IRE
SECIS	mir-TAR
mir-30	mir-9
lin-4	mir-5
mir-8	mir-8
mir-1	mir-1
mir-2	mir-6
let-7	Y_RNA
UAA_GAN	Csr
Cloop	domV
ktrn1	ktrn2
term1	term2
tRNA	RNaseP
SAM_v	symR
CPeB3	FinP
sroB	msr
SAM_u	HH_3
Vmtnn3	Vmtnn3
l1vK	DsrA
CAESAR	CAESAR
isrK	sroD
isrB	isrB
6C	6C
rspL	subB
dimB	dimB
suCA	SraD
	*
sky	*
RNAl	RNAl
Purine	SAM-Chl
cdGMP_2	Anti-Q
GadY	RnkJdr
PrfA	PrfA
OmrA-B	RyeB
traJ2	traJ2
SraH	23Smeth
23Smeth	DS-pep
Ps-Rho	rnk-ps
	**
Mgsns	Mgsns
tRNAs	Qrr
isrC	HH_1
SNR_24	TrpJdr
greA	greA
preQ12	HAR1F
TermLeu	TermLeu
MicC	C4
RsmY	RsmY
Ribosome	Ribosome

What is a motif?





- ▶ Wiktionary definitions for **motif**:
 - ▶ A recurring or dominant element; a theme.
- ▶ For the purposes of this work:
 - ▶ an RNA motif is a recurring RNA sequence and/or secondary structure found within larger structures that can be modelled by either a covariance model or a profile HMM.

Escher, MC (1938) Sky and Water 1.

What is a motif?



- ▶ The kink-turn RNA motif
 - ▶ Found in rRNA, riboswitches, RNase P, snoRNAs, ...
 - ▶ An asymmetric internal loop
 - ▶ Causes a sharp kink between two flanking helical regions



Cruz & Westhof (2011) Sequence-based identification of 3D structural modules in RNA with RMDetect. *Nature Methods*.




Can we build a seed alignment and CM resource of these motifs?

- ▶ For the hairpin motifs?
 - ▶ YES
 - ▶ For the internal loop motifs?
 - ▶ MAYBE
 - ▶ For the sequence motifs?
 - ▶ Not certain yet



Doshi et al. (2004) Evaluation of the suitability of free-energy minimization using nearest-neighbor energy parameters for RNA secondary structure prediction. *BMC Bioinformatics*.

The RNA Motifs: RMfam



5' - ● ● ● ● ● ● ● ● ● ● R R A R G G R G R R ● ● ● ● ● Y A U G R ● ● R ●

5' - ● ● ● ● ● ● ● ● ● ● R R G G R R ● ● ● R Y ● A U G A R ● R ●

5' - ● ● ● ● ● ● ● ● ● ● A A G G R R ● ● ● ● A U G ● ● R ●

Breaking Rules

- ▶ Prefer motifs found in multiple families &/or multiple locations in the same family
 - ▶ Aligning **analogous** structures
 - ▶ Motifs are allowed to overlap
 - ▶ Models are non-specific
 - ▶ High false-positive rates
 - ▶ Sequences are not edited, spliced or otherwise interfered with
-
- ▶ Sequence sources are diverse (PDB, EMBL & publications)




An example entry

```
# STOCKHOLM 1.0

#=GF ID      twist_up
#=GF AU      Gardner PP
#=GF SE      Gardner PP
#=GF SS      Published; PMID:21976732
#=GF GA      19.00
#=GF DR      URL;http://rnabrabase.cs.put.poznan.pl/?act=pdbdetails&id=1S72
#=GF RN      [1]
#=GF RM      21976732
#=GF RT      Clustering RNA structural motifs in ribosomal RNAs using
#=GF RT      secondary structural alignment.
#=GF RA      Zhong C, Zhang S
#=GF RL      Nucleic Acids Res. 2012;40:1307-17.
```

2zkr_Y/27-46 CCGAUCUCGUC-UGAUCUCGG
#=GR_2zkr_Y/27-46 SS <<<<---<----->--->>>
2gv3_A/3-20 CCAGACUC---CCGAAUCUGG
#=GR_2gv3_A/3-20 SS (((((. (.----....))))))
3iz9_C/29-48 CGGAUCCCGUC-AGAACUCCG
#=GR_3iz9_C/29-48 SS (((((...(. ...-.)...))))))
3bbo_B/31-50 CCAA-UCAUCCCGAACUUGG
#=GR_3bbo_B/31-50 SS .(.....<.....>.....).
1nkw_9/33-53 CCCACCCCAUGCCGAACUGGG
#=GR_1nkw_9/33-53 SS (((.....(.....))))
1c2x_C/31-51 CUGACCCCAUGCCGAACUCAG
#=GR_1c2x_C/31-51 SS ((((...<.....>....))))
1giy_B/33-53 CCGUUCCCAUCCGAACACGG
1S72_9/29-50 CCGUACCACAUCCGAACACGG
#=GR_1S72_9/29-50 SS (((((...(.)....))))
#=GR_1S72_9/29-50 MT ...XXXXXX.....XXXXXX...
#=GC SS_cons
//




An example entry

```
# STOCKHOLM 1.0

#=GF ID      twist_up
#=GF AU      Gardner PP
#=GF SE      Gardner PP
#=GF SS      Published; PMID:21976732
#=GF GA      19.00
#=GF DR      URL;http://rnabrabase.cs.put.poznan.pl/?act=pdbdetails&id=1S72
#=GF RN      [1]
#=GF RM      21976732
#=GF RT      Clustering RNA structural motifs in ribosomal RNAs using
#=GF RT      secondary structural alignment.
#=GF RA      Zhong C, Zhang S
#=GF RL      Nucleic Acids Res. 2012;40:1307-17.
```

2zkr_Y/27-46 CCGAUCUCGUC-UGAUCUCGG
#=GR_2zkr_Y/27-46 SS <<<<---< >--->>>
2gv3_A/3-20 CCAGACUC---CCGAAUCUGG
#=GR_2gv3_A/3-20 SS (((((.(-.-.----))))))
3iz9_C/29-48 CGGAUCCCGUC-AAACUCCG
#=GR_3iz9_C/29-48 SS ((((...(.--.----))))))
3bbo_B/31-50 CCAA-UCCAUCCCGAACUUGG
#=GR_3bbo_B/31-50 SS .((....<.....>....)).
1nkw_9/33-53 CCCACCCCCAUUGCCTAACUGGG
#=GR_1nkw_9/33-53 SS ((((.....-.....-.....))))
1c2x_C/31-51 CUGACCCCCAUGCCAACUCAG
#=GR_1c2x_C/31-51 SS ((((...<.....>....))))
1giy_B/33-53 CCGUUCCCCAUUCCGAACACCGG
#=GR_1giy_B/33-53 SS CCGUUACCCAUUCCGAACACCGG
1S72_9/29-50 ((((...(.--.----))))))
#=GR_1S72_9/29-50 SS ...XXXXXX.....XXXXXX...
#=GR_1S72_9/29-50 MT ((((...(.--.----))))))
#=GC SS_cons
//



An example annotated Rfam alignment

```


# STOCKHOLM 1.0
#=GF ID      5S rRNA
#=GF AC      RF00001
#...
#=GF WK      5S_ribosomal_RNA
#=GF SQ      712
#=GF MT.G    GNRA
#=GF MT.s    sarcin-ricin-2
#=GF MT.t    twist up

```




An example annotated Rfam alignment

```
# STOCKHOLM 1.0
#=GF ID      5S_rRNA
#=GF AC      RF00001
#...
#=GF WK      5S_ribosomal_RNA
#=GF SQ      712
#=GF MT.G    GNRA
#=GF MT.s    sarcin-ricin-2
#=GF MT.t    twist up
```




Uses for a motif DB (I): Improving a RNA alignments and structure prediction constraints




Uses for a motif DB (II): RNA structural motifs

► The Lysine Riboswitch




Uses for a motif DB (II): RNA structural motifs

► The Lysine Riboswitch




Uses for a motif DB (III): Functionally characterising novel RNAs



- ▶ Two AYGGAY sRNA: 216 homologues found in Human gut metagenome sequences, γ -Proteobacteria and Clostridiales species
- ▶ Weinberg Z et al. (2010)
"Comparative genomics reveals 104 candidate structured RNAs from bacteria, archaea and their metagenomes". *Genome Biol.*


Csr-Rsm background




Toledo-Arana *et al.* (2007) Small noncoding RNAs controlling pathogenesis. *Current Opinion in Microbiology*.

Uses for a motif DB (IV): Identifying and ranking ncRNAs of interest in transcriptome results

Distribution of motif scores on *S. typhi* ncRNAs




CDF of motif scores on *S. typhi* ncRNAs




Uses for a motif DB (IV): Identifying and ranking ncRNAs of interest in transcriptome results


Distribution of motif scores on *S. typhi* ncRNAs alignments




CDF of motif scores on *S. typhi* ncRNAs alignments



80 highest scoring matches between RMfam & Rfam



What about the highly expressed ncRNAs?




RNA	RNA code	RNAz	Motifs	complexity	Low Seq.	G+C
M.tub. RUF3	No	RNA (prob=1.00)	Terminator,k-turn	No		59% (66%)
N.gon. RUF7	No	RNA (prob=0.94)	Terminator	No		43% (52%)
C.dif. RUF9	? (P=0.044)	RNA (prob=1.00)	Terminator	No		30% (29%)
C.dif. RUF11	? (P=0.079)	RNA (prob=1.00)	Terminator	Yes (23/ 170nts)		26% (29%)




What about the highly expressed ncRNAs?

- Are they antisense to any 5' UTRs

RNAup RNA–RNA energies between sRNAs and 5' UTRs




Difference between native and shuffled CDFs




What about the highly expressed ncRNAs?

- Are they antisense to any 3' UTRs

RNAup RNA–RNA energies between sRNAs and 3' UTRs



Difference between native and shuffled CDFs





Discussion points

- ▶ A useful curation tool for building alignments and structures
- ▶ Can provide testable function hypotheses, sometimes
- ▶ Limited use outside of alignments based on the *Salmonella Typhi* results
- ▶ Even in alignments, false-positives & negatives are common
- ▶ More motifs (Terminators, Shine-Dalgarno, ...)
- ▶ Limit trivial motifs and prevent rediscovery of old motifs
- ▶ Snapshots of the data are available from
<https://github.com/ppgardne/RMfam>
- ▶ Can anyone here do better on MTB3, NGON7, CDIF9 & CDIF11?

Thanks!





- ▶ Lars Barquist, Alex Bateman & Rob Knight

RNAbiology




PPG is supported by a Rutherford Discovery Fellowship from Government funding, administered by the Royal Society of New Zealand.

RNA motifs



RNA classification



Backofen et al. (2007) RNAs everywhere: genome-wide annotation of structured RNAs. *Journal of Experimental Zoology*.