

Gravitational waves from phase transitions

3. Generation of gravitational waves

Mark Hindmarsh^{1,2}

¹Department of Physics & Astronomy
University of Sussex

²Department of Physics and Helsinki Institute of Physics
Helsinki University

Benasque GW School
2. kesäkuuta 2017

Outline

Recap

Outline

Gravitational waves and shear stresses

Gravitational wave power spectrum discussion

Summary and outlook

Yesterday ...

- Bubble nucleation rate/volume

$$\Gamma(T) = \Gamma_0(T) e^{-S(T)}$$

- Transition rate parameter β

$$\Gamma(t) = \Gamma_f e^{\beta(t-t_f)}$$

with $\frac{4\pi}{3} v_w^3 (3!/\beta^4) \Gamma_f = 1$


- Wall speed v_w

- Transition strength $\alpha_+ = \frac{4\epsilon}{3w_+}$

- Junction conditions ($r = w_+/w_-$)

$$v_+ v_- = \frac{1 - (1 - 3\alpha_+)r}{3 - 3(1 + \alpha_+)r}, \quad \frac{v_+}{v_-} = \frac{3 + (1 - 3\alpha_+)r}{1 + 3(1 + \alpha_+)r}$$

- Similarity solution for bubble growth: detonation, deflagration, hybrid
- Conversion efficiency κ



Gravitational waves from shear stresses


Metric perturbations h_{ij} from transverse-traceless part of EM tensor Π_{ij} :

$$\ddot{h}_{ij} - \nabla^2 h_{ij} = 16\pi G \Pi_{ij}$$

$$\Pi_{ij}^{\text{f}} = \left[(e + p) \gamma^2 v_i v_j + p \delta_{ij} \right]^{TT}$$


$$\Pi_{ij}^{\phi} = \left[\partial_i \phi \partial_j \phi - \frac{1}{2} (\partial \phi)^2 \delta_{ij} \right]^{TT}$$

GW power spectrum: $\frac{d\rho_{\text{GW}}(k)}{d \ln k} = \frac{k^3}{32\pi G} \int \frac{d\Omega}{(2\pi)^3} \dot{h}_{ij}(t, \mathbf{k}) \dot{h}_{ij}^*(t, \mathbf{k})$




Detection prospects

Space-based GW detector LISA (launch by 2034):



- ▶ e.g. strong EW transition with eLISA configurations: best (r), worst (g) (Caprini et al. 2015)
- ▶ $\alpha = 0.5$, $\beta/H = 100$, $v_w = 0.95$
- ▶ Total (k) sound (g) turbulence (r)



Gravitational wave equation (and an auxiliary equation)

- ▶ Assume processes happen much faster than Hubble rate ($\beta/H \gg 1$)
- ▶ Assume metric perturbations are small
- ▶ Linearised GR, neglect expansion⁽¹⁾

$$ds^2 = -dt^2 + (\delta_{ij} + h_{ij})dx^i dx^j$$

- ▶ Linearised Einstein eqn for transverse-traceless part:

$$\ddot{h}_{ij} - \nabla^2 h_{ij} = 16\pi G \Pi_{ij}$$

- ▶ Convenient to avoid TT for evolution⁽²⁾

$$\ddot{u}_{ij} - \nabla^2 u_{ij} = 16\pi G \tau_{ij}$$

where $h_{ij} = u_{ij}^{TT}$ and $\Pi_{ij} = \tau_{ij}^{TT}$.

- ▶ Can take

$$\tau_{ij}^f = (\mathbf{e} + \mathbf{p})\gamma^2 v_i v_j, \quad \tau_{ij}^\phi = \partial_i \phi \partial_j \phi$$

⁽¹⁾ Can put expansion in, not much changes

⁽²⁾ Garcia-Bellido, Figueroa, Sartre 2008

Gravitational wave spectral density

- ▶ GW energy density (average over many wavelengths, periods)

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \overline{\dot{h}_{ij}(x) \dot{h}_{ij}(x)}$$

- ▶ Fourier transform: $\dot{h}_{ij}(\mathbf{k}, t) = \int d^3x \dot{h}_{ij}(\mathbf{x}, t) e^{-i\mathbf{k} \cdot \mathbf{x}}$
- ▶ Define **spectral density** P_h through

$$\langle \dot{h}_{ij}(\mathbf{k}, t) \dot{h}_{ij}^*(\mathbf{k}', t) \rangle = P_h(\mathbf{k}) \delta^3(\mathbf{k} - \mathbf{k}')$$

- ▶ Assume gravitational waves are generated by a process which is
 - ▶ isotropic: $P_h(\mathbf{k}) \rightarrow P_h(k)$
 - ▶ homogeneous (hence $\delta^3(\mathbf{k} - \mathbf{k}')$)
 - ▶ Random, Gaussian
- ▶ GW energy density becomes

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \int d^3k P_h(k) = \frac{1}{32\pi G} \frac{4\pi}{(2\pi)^3} \int dk k^2 P_h(k)$$

Gravitational wave spectral density

- ▶ GW energy density (average over many wavelengths, periods)

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \overline{\dot{h}_{ij}(x)\dot{h}_{ij}(x)}$$

- ▶ Fourier transform: $\dot{h}_{ij}(\mathbf{k}, t) = \int d^3x \dot{h}_{ij}(\mathbf{x}, t) e^{-i\mathbf{k}\cdot\mathbf{x}}$
- ▶ Define **spectral density** P_h through

$$\langle \dot{h}_{ij}(\mathbf{k}, t) \dot{h}_{ij}^*(\mathbf{k}', t) \rangle = P_h(\mathbf{k}) \delta^3(\mathbf{k} - \mathbf{k}')$$

- ▶ Assume gravitational waves are generated by a process which is
 - ▶ isotropic: $P_h(\mathbf{k}) \rightarrow P_h(k)$
 - ▶ homogeneous (hence $\delta^3(\mathbf{k} - \mathbf{k}')$)
 - ▶ Random, Gaussian
- ▶ GW energy density becomes

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \int d^3k P_h(k) = \frac{1}{32\pi G} \frac{1}{2\pi^2} \int dk k^2 P_h(k)$$

Gravitational wave power spectrum

- ▶ Recall GW energy density from spectral density

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \frac{1}{2\pi^2} \int dk k^2 P_h(k)$$

- ▶ Convenient to introduce **power spectrum** $\mathcal{P}_h = \frac{k^3}{2\pi^2} P_h(k)$

$$\rho_{\text{gw}} = \frac{1}{32\pi G} \int \frac{dk}{k} \mathcal{P}_h(k)$$

- ▶ Cosmology: characterisation better in terms of $\Omega_{\text{gw}} = \rho_{\text{gw}}/\bar{\epsilon}$.
- ▶ Define **gravitational wave power spectrum**

$$\mathcal{P}_{\text{gw}}(k) \equiv \frac{d\Omega_{\text{gw}}}{d\ln(k)} = \frac{1}{\bar{\epsilon}} \frac{1}{32\pi G} \mathcal{P}_h(k) = \frac{1}{12H^2} \mathcal{P}_h(k)$$

GW from stochastic sources

- Equation for auxiliary tensor $u_{ij}(\mathbf{x}, t)$

$$(\partial_t^2 - \nabla^2)u_{ij}(\mathbf{x}, t) = (16\pi G)\tau_{ij}(\mathbf{x}, t)$$

- Solution with oscillator Green's function⁽³⁾

$$u_{ij}(\mathbf{k}, t) = (16\pi G) \int_0^t dt' \frac{\sin[k(t - t')]}{k} \tau_{ij}(\mathbf{k}, t')$$

- Gravitational wave from TT projector $\dot{h}_{ij}(\mathbf{k}, t) = \lambda_{ij,kl}(\mathbf{k}) \dot{u}_{kl}(\mathbf{k}, t)$
- Projector: $\lambda_{ij,kl}(\mathbf{k}) = P_{ik}(\mathbf{k})P_{jl}(\mathbf{k}) - \frac{1}{2}P_{ij}(\mathbf{k})P_{kl}(\mathbf{k})$ with $P_{ij}(\mathbf{k}) = \delta_{ij} - \hat{k}_i\hat{k}_j$.
- GW power spectrum obtained from

$$\begin{aligned} \langle \dot{h}_{\mathbf{k}}^{ij}(t) \dot{h}_{\mathbf{k}'}^{ij}(t) \rangle = \\ (16\pi G)^2 \int_0^t dt_1 dt_2 \cos[k(t - t_1)] \cos[k(t - t_2)] \lambda_{ij,kl}(\mathbf{k}) \langle \tau_{\mathbf{k}}^{ij}(\mathbf{k}, t_1) \tau_{\mathbf{k}'}^{kl}(\mathbf{k}', t_2) \rangle. \end{aligned}$$

⁽³⁾Boundary condition: $u_{ij}(\mathbf{k}, t) \rightarrow 0$ as $t \rightarrow 0$

GW from stochastic sources

- Equation for auxiliary tensor $u_{ij}(\mathbf{k}, t)$

$$(\partial_t^2 + \mathbf{k}^2)u_{ij}(\mathbf{k}, t) = (16\pi G)\tau_{ij}(\mathbf{k}, t)$$

- Solution with oscillator Green's function⁽³⁾

$$u_{ij}(\mathbf{k}, t) = (16\pi G) \int_0^t dt' \frac{\sin[k(t - t')]}{k} \tau_{ij}(\mathbf{k}, t')$$

- Gravitational wave from TT projector $\dot{h}_{ij}(\mathbf{k}, t) = \lambda_{ij,kl}(\mathbf{k}) \dot{u}_{kl}(\mathbf{k}, t)$
- Projector: $\lambda_{ij,kl}(\mathbf{k}) = P_{ik}(\mathbf{k})P_{jl}(\mathbf{k}) - \frac{1}{2}P_{ij}(\mathbf{k})P_{kl}(\mathbf{k})$ with $P_{ij}(\mathbf{k}) = \delta_{ij} - \hat{k}_i\hat{k}_j$.
- GW power spectrum obtained from

$$\begin{aligned} \langle \dot{h}_{\mathbf{k}}^{ij}(t) \dot{h}_{\mathbf{k}'}^{ij}(t) \rangle = \\ (16\pi G)^2 \int_0^t dt_1 dt_2 \cos[k(t - t_1)] \cos[k(t - t_2)] \lambda_{ij,kl}(\mathbf{k}) \langle \tau_{\mathbf{k}}^{ij}(\mathbf{k}, t_1) \tau_{\mathbf{k}'}^{kl}(\mathbf{k}', t_2) \rangle. \end{aligned}$$

⁽³⁾Boundary condition: $u_{ij}(\mathbf{k}, t) \rightarrow 0$ as $t \rightarrow 0$


Unequal time correlator (UETC) for shear stress

- Define shear stress UETC Π^2 by

$$\lambda_{ij,kl}(\mathbf{k}) \langle \tau^{ij}(\mathbf{k}, t_1) \tau^{kl}(\mathbf{k}', t_2) \rangle = \Pi^2(k, t_1, t_2) \delta^3(\mathbf{k} + \mathbf{k}')$$

- Form of UETC

- source is “on” for a time τ_Π
- auto-correlated for a time τ_{ac}
- peak at wavenumber $k \sim L_\Pi$



GW spectral density from shear stress UETC

- ▶ Averaging over a many periods of the wave⁽⁴⁾

$$P_{\dot{h}}(k, t) = (16\pi G)^2 \frac{1}{2} \int_0^t dt_1 dt_2 \cos[k(t_1 - t_2)] \Pi^2(k, t_1, t_2).$$

- ▶ Assumed form of UETC (write $t_+ = (t_1 + t_2)/2$, $t_- = t_1 - t_2$)
 $\Pi^2(k, t_-/\tau_{ac}) \theta(\tau_{\Pi} - t_+)$
- ▶ Results in spectral density for \dot{h}

$$P_{\dot{h}}(k, t) = (16\pi G)^2 \tau_{\Pi} \tau_{ac} \mathcal{C}_{\Pi}(k)$$

where $\mathcal{C}_{\Pi}(k) = \frac{1}{2} \int \frac{dt_-}{\tau_{ac}} \cos(kt_-) \Pi^2(k, t_-/\tau_{ac})$

- ▶ Available scales for auto-correlation time τ_{ac} :
 - ▶ $\tau_{ac} \sim k^{-1}$
 - ▶ $\tau_{ac} \sim L_{\Pi}$
 - ▶ $\tau_{ac} \sim \tau_{\Pi}$

⁽⁴⁾So that $\cos[k(t - t_1)] \cos[k(t - t_2)] \rightarrow \frac{1}{2} \cos[k(t_1 - t_2)]$

Sources of shear stress: fluid vs. scalar field

- ▶ Estimate size of shear stress correlator: $\Pi^2 \sim \langle \tau \tau \rangle$
 - ▶ Fluid source tensor $\tau_f^{ij} = w \gamma^2 v^i v^j$
 - ▶ Field source tensor $\tau_\phi^{ij} = \partial^i \phi \partial^j \phi$
- ▶ Kinetic energies $K_f = \int d^3x \tau_{ff}^f$, $K_\phi = \int d^3x \tau_{ff}^\phi$
 - ▶ Fluid: $K_f = \int d^3x w \gamma^2 v^2 = \frac{4\pi}{3} R^3 \bar{w} \frac{3}{4} \alpha \kappa$
 - ▶ Field: $K_\phi = \int d^3x (\nabla \phi)^2 = 4\pi R^2 \sigma$
- ▶ Ratio $K_f/K_\phi \sim R \bar{w}/\sigma \sim R/\ell \gg 1$
 - ▶ Bubble size R grows to Hubble length, ℓ is a microscopic scale (wall width).
- ▶ Scalar coupled to fluid, similarity solution: fluid shear stress dominant
 - ▶ Fluid shear stresses come from compression/rarefaction: **sound waves**
- ▶ Runaway: field shear stress also grows as R^3 (not considered here)

Sound waves

Consider EM tensor for perturbations with z dependence only

$$T^{tt} = w\gamma^2 - p, \quad T^{tz} = w\gamma^2 v^z, \quad T^{zz} = w\gamma^2 (v^z)^2 + p$$

Perturbations: $\delta e = e - \bar{e}$, $\delta p = p - \bar{p}$, v^z all $\ll 1$

$$\partial_t T^{tt} + \partial_z T^{zt} = 0 \implies \partial_t(\delta e) + \bar{w}\partial_z v^z = 0 \quad (1)$$

$$\partial_t T^{tz} + \partial_z T^{zz} = 0 \implies \bar{w}\partial_t v^z + \partial_z(\delta p) = 0 \quad (2)$$

Note that δp and δe both depends temperature T : $\delta p = \left(\frac{\partial p}{\partial T} / \frac{\partial e}{\partial T} \right) \delta e = c_s^2 \delta e$

Hence equations (1) and (2) can be combined

$$\left(\partial_t^2 - c_s^2 \partial_z^2 \right) v^z = 0, \quad \left(\partial_t^2 - c_s^2 \partial_z^2 \right) \delta T = 0$$

Sound wave is a collective mode of fluid velocity v^i and temperature T .
It is longitudinal: v^i is in direction of travel of wave.

Shear stress UETC from sound waves 1

- Recall shear stress UETC Π^2 :

$$\lambda_{ij,kl}(\mathbf{k}_1) \langle \tau^{ij}(\mathbf{k}_1, t_1) \tau^{kl}(\mathbf{k}_2, t_2) \rangle = \Pi^2(k_1, t_1, t_2) \delta^3(\mathbf{k}_1 + \mathbf{k}_2)$$

- Source tensor dominated by fluid: $\tau^{ij} = \tau_f^{ij} = w \gamma^2 v^i v^j$
- Non-relativistic fluid velocities: $\tau_f^{ij} \simeq \bar{w} v^i v^j$
- Fourier transform of velocity field $\tilde{v}^i(\mathbf{q}, t) = \int d^3x v^i(\mathbf{x}, t) e^{-i\mathbf{q}\cdot\mathbf{x}}$
- Hence

$$\tau_f^{ij}(\mathbf{k}, t) = \bar{w} \int d^3q \tilde{v}^i(\mathbf{q}, t) \tilde{v}^j(\tilde{\mathbf{q}}, t), \quad \tilde{\mathbf{q}} = \mathbf{q} - \mathbf{k}$$

- Assume velocity field is Gaussian: $\langle \tau \tau \rangle \sim \langle v v v v \rangle = \sum \langle v v \rangle \langle v v \rangle$
- Velocity unequal time correlator:

$$\langle \tilde{v}_{\mathbf{q}_1}^i(t_1) \tilde{v}_{\mathbf{q}_2}^{*j}(t_2) \rangle = \left[P_{ij}(q) F(q, t_1, t_2) + \hat{q}^i \hat{q}^j G(q, t_1, t_2) \right] \delta^3(\mathbf{q}_1 - \mathbf{q}_2).$$

- Transverse projector $P_{ij}(q) = \delta_{ij} - \hat{q}^i \hat{q}^j$
- Sound waves contribute only to longitudinal part G

Shear stress UETC from sound waves 2

- Recall shear stress UETC Π^2 :

$$\lambda_{ij,kl}(\mathbf{k}_1) \langle \tau^{ij}(\mathbf{k}_1, t_1) \tau^{kl}(\mathbf{k}_2, t_2) \rangle = \Pi^2(k_1, t_1, t_2) \delta^3(\mathbf{k}_1 + \mathbf{k}_2)$$

- With $\tau_f^{ij}(\mathbf{k}, t) = \bar{w} \int \bar{d}^3 q \tilde{v}^i(\mathbf{q}, t) \tilde{v}^j(\tilde{\mathbf{q}}, t)$, $\tilde{\mathbf{q}} = \mathbf{q} - \mathbf{k}$
- And the velocity unequal time correlator from sound waves:
 $\langle \tilde{v}_{\mathbf{q}_1}^i(t_1) \tilde{v}_{\mathbf{q}_2}^{*j}(t_2) \rangle = \hat{q}^i \hat{q}^j G(q, t_1, t_2) \delta^3(\mathbf{q}_1 - \mathbf{q}_2)$.
- A long calculation gives:⁽⁵⁾

$$\Pi^2(k, t_1, t_2) = \bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 G(q, t_1, t_2) G(\tilde{q}, t_1, t_2)$$

$$\text{where } \mu = \hat{\mathbf{k}} \cdot \hat{\mathbf{q}}, \tilde{q}^2 = q^2 - 2qk\mu + k^2.$$

⁽⁵⁾Caprini, Durrer, Servant 2007, 2009

Velocity UETC from sound waves

- ▶ Following completion of transition at t_i , sound waves propagate freely
- ▶ General sound wave solution:

$$v^i(\mathbf{x}, t) = \int \bar{d}^3 q \left(v_{\mathbf{q}}^i e^{-i\omega t + i\mathbf{q} \cdot \mathbf{x}} + v_{\mathbf{q}}^{*i} e^{i\omega t - i\mathbf{q} \cdot \mathbf{x}} \right), \quad \omega = c_s q$$

- ▶ Spectral density of velocity plane wave amplitudes

$$\langle v_{\mathbf{q}}^i v_{\mathbf{q}'}^j \rangle = \hat{q}^i \hat{q}^j P_v(q) \delta^3(\mathbf{q} + \mathbf{q}')$$

- ▶ Recall velocity UETC $\langle \tilde{v}_{\mathbf{q}}^i(t_1) \tilde{v}_{\mathbf{q}'}^{*j}(t_2) \rangle = \hat{q}^i \hat{q}^j G(q, t_1, t_2) \delta^3(\mathbf{q} - \mathbf{q}')$.
- ▶ Plane wave amplitudes related to Fourier transform $\tilde{v}_{\mathbf{q}}^i(t) = 2v_{\mathbf{q}}^i \cos(\omega t)$
- ▶ Hence (recall $t_+ = (t_1 + t_2)/2$, $t_- = t_1 - t_2$)

$$G(q, t_1, t_2) = 4P_v(q) \cos(\omega t_1) \cos(\omega t_2) = 2P_v(q) [\cos(\omega t_-) + \cos(2\omega t_+)]$$

Shear stress UETC from sound waves 3

- Recall expressions for shear stress UETC and velocity UETC

$$\Pi^2(k, t_1, t_2) = \bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 G(q, t_1, t_2) G(\tilde{q}, t_1, t_2)$$

$$G(q, t_1, t_2) = 2P_v(q) [\cos(\omega t_-) + \cos(2\omega t_+)]$$

- Argument:

- Convolution to get Π^2 has integral over $q = \omega/c_s$
- For large times $t_+ \gg \omega^{-1}$ and $\cos(2\omega t_+)$ is highly oscillatory
- $\Rightarrow \Pi^2$ dominated by $\cos^2(\omega t_-)$ terms

$$\Pi^2(k, t_1, t_2) = \bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 P_v(q) P_v(\tilde{q}) \cos(\omega t_-) \cos(\tilde{\omega} t_-)$$

- Conclusions:

- $\Pi^2(k, t_1, t_2)$ depends mostly on $t_- = t_1 - t_2$ (“stationary”)
- Autocorrelation time of mode with wavenumber k is $\tau_{ac} \sim k^{-1}$

GW from UETC

- ▶ Recall in spectral density for \dot{h}

$$P_{\dot{h}}(k, t) = (16\pi G)^2 \tau_{\Pi} \tau_{\text{ac}} C_{\Pi}(k)$$

where $C_{\Pi}(k) = \frac{1}{2} \int \frac{dt_-}{\tau_{\text{ac}}} \cos(kt_-) \Pi^2(k, t_- / \tau_{\text{ac}})$

- ▶ We justified dropping the dependence of Π^2 on t_+
- ▶ We argued that $\tau_{\text{ac}} = k^{-1}$, so

$$C_{\Pi}(k) = \frac{1}{2} k \int dt_- \cos(kt_-) \Pi^2(k, kt_-)$$
- ▶ Hence

$$C_{\Pi}(k) = 4\bar{w}^2 \int \bar{d}^3q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 P_{\nu}(q) P_{\nu}(\tilde{q}) \Delta(k, \omega, \tilde{\omega})$$

where $\Delta(k, \omega, \tilde{\omega}) = \frac{1}{2} \int dt_- \cos(kt_-) \cos(\omega t_-) \cos(\tilde{\omega} t_-)$

Kinematics of GW production from sound waves

- ▶ Spectral density of metric perturbation $P_h(k, t) = (16\pi G)^2 \pi k^{-1} C_\Pi(k)$

$$C_\Pi(k) = 4\bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 P_v(q) P_v(\tilde{q}) \Delta(k, \omega, \tilde{\omega})$$

$$\Delta(k, \omega, \tilde{\omega}) = \frac{1}{2} \int dt_- \cos(kt_-) \cos(\omega t_-) \cos(\tilde{\omega} t_-)$$

- ▶ For large time differences:

$$\Delta(k, \omega, \tilde{\omega}) \rightarrow \pi \delta(k \pm \omega \pm \tilde{\omega})$$

- ▶ Recall $\omega = c_s q$,


$$\tilde{\omega} = c_s \tilde{q} = c_s (q^2 + k^2 - 2kq\mu)$$

- ▶ Kinematics: only $k - \omega - \tilde{\omega}$ can vanish

- ▶ Conservation of energy for production of GWs

- ▶ Hence

$$C_\Pi(k) = 4\pi \bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 P_v(q) P_v(\tilde{q}) \delta(k - \omega - \tilde{\omega})$$



Kinematics of GW production from sound waves

- ▶ Spectral density of metric perturbation $P_h(k, t) = (16\pi G)^2 \tau_{\Pi} k^{-1} C_{\Pi}(k)$

$$C_{\Pi}(k) = 4\pi \bar{w}^2 \int \bar{d}^3 q \frac{q^2}{\tilde{q}^2} (1 - \mu^2)^2 P_v(q) P_v(\tilde{q}) \delta(k - \omega - \tilde{\omega})$$

- ▶ Use δ function to perform integral over $\mu = \hat{\mathbf{k}} \cdot \hat{\mathbf{q}}$
 - ▶ Solution: $\mu = \mu_* = \frac{1}{c_s} (1 - \frac{1}{2} (1 - c_s^2) \frac{k}{q})$
 - ▶ Giving $\tilde{q} = \tilde{q}_* = \frac{1}{c_s} k - q$
- ▶ Only $q_- < q < q_+$ can produce GW with frequency k : with $q_{\pm} = k \frac{1 \pm c_s}{2c_s}$

$$C_{\Pi}(k) = \frac{\bar{w}^2}{2\pi c_s} \int_{q_-}^{q_+} dq q^2 \frac{q}{\tilde{q}} (1 - \mu_*^2)^2 P_v(q) P_v(\tilde{q}_*)$$

Kinematics of GW production from sound waves

- ▶ Assume that sound waves have length scale L_f
- ▶ Scale out mean square velocity \overline{U}_f^2
- ▶ Hence $P_v(q) = \overline{U}_f^2 L_f^3 \tilde{P}_v(qL_f)$, where \tilde{P}_v is dimensionless.
- ▶ Define $z = qL_f$ and $z_{\pm} = (kL_f)^{\frac{1 \pm c_s}{2c_s}}$

$$C_{\Pi}(k) = \frac{\bar{w}^2 \overline{U}_f^4}{2\pi c_s} L_f^3 \int_{z_-}^{z_+} \frac{dz}{z} \frac{(z - z_+)^2 (z - z_-)^2}{z_+ + z_- - z} \tilde{P}_v(z) \tilde{P}_v(z_+ + z_- - z)$$

- ▶ Define dimensionless function

$$\tilde{C}_{\Pi}(kL_f) = \int_{z_-}^{z_+} \frac{dz}{z} \frac{(z - z_+)^2 (z - z_-)^2}{z_+ + z_- - z} \tilde{P}_v(z) \tilde{P}_v(z_+ + z_- - z)$$

- ▶ is dimensionless and a function of $y = kL_f$
- ▶ peaks at $y \sim 1$ (definition of L_f)
- ▶ Magnitude $O(1)$

Power spectrum of GWs from sound waves

- ▶ The final pieces are

$$\mathcal{P}_{\text{gw}}(k) \equiv \frac{d\Omega_{\text{gw}}}{d\ln(k)} = \frac{1}{12H^2} \frac{k^3}{2\pi^2} P_h(k)$$

$$P_h(k, t) = (16\pi G)^2 \tau_{\Pi} k^{-1} C_{\Pi}(k)$$

$$C_{\Pi}(k) = \frac{\bar{w}^2 \bar{U}_f^4}{2\pi c_s} L_f^3 \tilde{C}_{\Pi}(kL_f)$$

$$\tilde{C}_{\Pi}(kL_f) = \int_{z_-}^{z_+} \frac{dz}{z} \frac{(z - z_+)^2 (z - z_-)^2}{z_+ + z_- - z} \tilde{P}_v(z) \tilde{P}_v(z_+ + z_- - z)$$

- ▶ The gravitational wave power spectrum is

$$\mathcal{P}_{\text{gw}}(k) = 3\Gamma^2 \bar{U}_f^4 (H\tau_{\Pi})(HL_f) \frac{(kL_f)^3}{2\pi^2} \frac{\tilde{C}_{\Pi}(kL_f)}{2\pi c_s kL_f}.$$

where $\Gamma = \bar{w}/\bar{e}$ (adiabatic index, $\Gamma \simeq 4/3$)

Power spectrum of GWs from sound waves: amplitude

- ▶ From last slide

$$\mathcal{P}_{\text{gw}}(k) = 3\Gamma^2 \overline{U}_f^4 (H\tau_{\Pi})(HL_f) \tilde{\mathcal{P}}_{\text{gw}}(kL_f)$$

$$\tilde{\mathcal{P}}_{\text{gw}}(kL_f) = \frac{(kL_f)^3}{2\pi^2} \frac{\tilde{\mathcal{C}}_{\Pi}(kL_f)}{2\pi c_s kL_f}$$

$$\tilde{\mathcal{C}}_{\Pi}(kL_f) = \int_{z_-}^{z_+} \frac{dz}{z} \frac{(z - z_+)^2 (z - z_-)^2}{z_+ + z_- - z} \tilde{\mathcal{P}}_v(z) \tilde{\mathcal{P}}_v(z_+ + z_- - z)$$

where $z_{\pm} = (kL_f)^{\frac{1 \pm c_s}{2c_s}}$

- ▶ Power proportional to square of kinetic energy: $\Gamma \overline{U}_f^2 \simeq \kappa \alpha$
 - ▶ Phase transition strength parameter α
 - ▶ Vacuum to kinetic energy conversion efficiency κ
- ▶ Power proportional to length scale L_f
 - ▶ Length scale set by bubble separation: $L_f \sim R_* = (8\pi)^{\frac{1}{3}} v_w / \beta$
- ▶ Power proportional to time sound waves last τ_{Π} , which is the shorter of
 - ▶ Hubble time $\tau_H = H^{-1}$
 - ▶ Eddy turn-over time $\tau_e \sim L_f / \overline{U}_f$ (time for turbulence to develop)

Power spectrum of GWs from sound waves: shape

- ▶ From last-but-one slide

$$\mathcal{P}_{\text{gw}}(k) = 3\Gamma^2 \overline{U}_f^4 (H\tau_{\Pi})(HL_f) \tilde{\mathcal{P}}_{\text{gw}}(kL_f)$$

$$\tilde{\mathcal{P}}_{\text{gw}}(kL_f) = \frac{(kL_f)^3}{2\pi^2} \frac{\tilde{\mathcal{C}}_{\Pi}(kL_f)}{2\pi c_s kL_f}$$

$$\tilde{\mathcal{C}}_{\Pi}(kL_f) = \int_{z_-}^{z_+} \frac{dz}{z} \frac{(z - z_+)^2 (z - z_-)^2}{z_+ + z_- - z} \tilde{\mathcal{P}}_v(z) \tilde{\mathcal{P}}_v(z_+ + z_- - z)$$

where $z_{\pm} = (kL_f)^{\frac{1 \pm c_s}{2c_s}}$

- ▶ If velocity power spectrum $\mathcal{P}_v \sim q^n$, then $\tilde{\mathcal{P}}_v(z) \sim z^{n-3}$
- ▶ $\implies \tilde{\mathcal{C}}_{\Pi}(y) \sim y^3 (y^{n-3})^2 \sim y^{2n-3}$
- ▶ $\implies \tilde{\mathcal{P}}_{\text{gw}}(y) \sim y^3 y^{2n-4} \sim y^{2n-1} \propto k^{2n-1}$
- ▶ Prediction: $\mathcal{P}_{\text{gw}}(k) \sim k^{2n-1}$ from $\mathcal{P}_v(q) \sim q^n$

Modelling GWs from sound waves

- Phenomenological model for GW power spectrum from **linear** sound waves (v_w not near sound speed c_s)
 - Bubble nucleation temperature T_n
 - Hubble rate H_n
 - Mean bubble separation R_* , peak of power spectrum at $z_p = k_p R_* \simeq 10$

$$\frac{d\Omega_{\text{gw},0}}{d\ln(f)} = 0.68 F_{\text{gw},0} \Gamma^2 \bar{U}_f^4 (H_n R_*) \tilde{\Omega}_{\text{gw}} C \left(\frac{f}{f_{p,0}} \right).$$

where

$$C(s) = s^3 \left(\frac{7}{4 + 3s^2} \right)^{7/2}.$$

dilution of GWs since matter-domination

$$F_{\text{gw},0} = (3.57 \pm 0.05) \times 10^{-5} \left(\frac{100}{h_*} \right)^{\frac{1}{3}}.$$

Peak frequency

$$f_{p,0} \simeq 26 \left(\frac{1}{H_n R_*} \right) \left(\frac{z_p}{10} \right) \left(\frac{T_n}{10^2 \text{ GeV}} \right) \left(\frac{h_*}{100} \right)^{\frac{1}{6}} \mu\text{Hz},$$

Summary and outlook

- ▶ Gravitational wave production from 1st order phase transition
 - ▶ Promising sensitivity to EW transition from [LISA](#) (Taiji ...)
- ▶ Areas for further work:
 - ▶ GWs from turbulent velocity field
 - ▶ GWs from runaway bubble (scalar field dominant)
 - ▶ Parameter extraction from GW power spectrum α, β, v_W, T_C
 - ▶ Improve accuracy of calculations of α, β, v_W, T_C from fundamental theory
 - ▶ Collider signals and phase transition parameters
- ▶ Future:
 - ▶ information about fundamental physics from the GW background at LISA