

# BAO analysis from the DR14 QSO sample

Héctor Gil-Marín (on behalf of the eBOSS QC WG)

Laboratoire de Physique Nucleaire et de Hautes Energies (LPNHE)  
Institut Lagrange de Paris (ILP)

Understanding Cosmological Observations @ Benasque  
1st Aug 2017



## Public DR14 data (yesterday!) arXiv:1707.09322

THE FOURTEENTH DATA RELEASE OF THE SLOAN DIGITAL SKY SURVEY: FIRST SPECTROSCOPIC DATA FROM THE EXTENDED BARYON OSCILLATION SKY SURVEY AND FROM THE SECOND PHASE OF THE APACHE POINT OBSERVATORY GALACTIC EVOLUTION EXPERIMENT


BELA ABOLFATH<sup>1</sup>, D. S. AGUADO<sup>2</sup>, GABRIELA AGUILAR<sup>3</sup>, CARLOS ALLENDE PRIETO<sup>2,4</sup>, ANDRES ALMEIDA<sup>5</sup>, TONIMA TASNIM ANANNA<sup>6</sup>, FRIEDRICH ANDERS<sup>7</sup>, SCOTT F. ANDERSON<sup>8</sup>, BRETT H. ANDREWS<sup>9</sup>, BORJA ANGUIANO<sup>10</sup>, ALFONSO ARAGÓN-SALAMANCA<sup>11</sup>, MARIA ARGUDO-FERNÁNDEZ<sup>12</sup>, ERIC ARMENGAUD<sup>13</sup>, METIN ATA<sup>1</sup>, ERIC AUBOURG<sup>14</sup>, VLADIMIR AVILA-REESE<sup>5</sup>, CARLES BADENES<sup>5</sup>, STEPHEN BAILEY<sup>7,15</sup>, KATHLEEN A. BARGER<sup>16</sup>, JORGE BARRERA-BALLESTEROS<sup>17</sup>, CURTIS BARTOSZ<sup>2</sup>, DOMINIC BATES<sup>18</sup>, FALK BAUMGARTEN<sup>19</sup>, JULIAN BAUTISTA<sup>20</sup>, RACHAEL BEATON<sup>21</sup>, TIMOTHY C. BEERS<sup>22</sup>, FRANCESCO BELFIORE<sup>23,24,25</sup>, CHAD F. BENDER<sup>26</sup>, MARIANGELA BERNARDI<sup>27</sup>, MATTHEW A. BERSHADY<sup>28</sup>, FLORIAN BEUTLER<sup>29</sup>, JONATHAN C. BIRD<sup>30</sup>, DMITRY BIZYAEV<sup>31,32,33</sup>, GUILLERMO A. BLANG<sup>31</sup>, MICHAEL R. BLANTON<sup>34</sup>, MICHAEL BLOMQUIST<sup>35</sup>, ADAM S. BOLTON<sup>36</sup>, MÉDÉRIC BOQUIEN<sup>12</sup>, JURA BORISSOVA<sup>37,38</sup>, JO BOVY<sup>39,40,41</sup>, CHRISTIAN ANDRES BRADNA DIAZ<sup>42</sup>, WILLIAM NIELSEN BRANDT<sup>43,44,45</sup>, JONATHAN BRINKMANN<sup>31</sup>, JOEL R. BROWNSTEIN<sup>20</sup>, KEVIN BUNDY<sup>25</sup>, ADAM J. BURGASSER<sup>46</sup>, ETIENNE BURTON<sup>13</sup>, NICOLÁS G. BUSCA<sup>14</sup>, CALEB I. CAÑAS<sup>43</sup>, MARIANA CANO-DÍAZ<sup>17</sup>, MICHELE CAPPELLARI<sup>48</sup>, RICARDO CARRERA<sup>2,4</sup>, ANDREW R. CASEY<sup>49</sup>, YANPING CHEN<sup>50</sup>, BRIAN CHERINKA<sup>51</sup>, CRISTINA CHIAPPINI<sup>7</sup>, PETER DOOHYUN CHOI<sup>52</sup>, DREW CHOJNOWSKI<sup>52</sup>, CHIA-HSUN CHUANG<sup>1</sup>, HAEUN CHUNG<sup>53</sup>, NICOLAS CLERC<sup>54,55,56</sup>, ROGER E. COHEY<sup>57,58</sup>, JOHAN COMPARAT<sup>54</sup>, JANAINA CORREA DO NASCIMENTO<sup>59,60</sup>, LUIZ DA COSTA<sup>60,61</sup>, MARIE-CLAUDE COUSINOU<sup>62</sup>, KEVIN COVEY<sup>63</sup>, JEFFREY D. CRANE<sup>21</sup>, IRENE CRUZ-GONZALEZ<sup>2</sup>, KATIA CUNHA<sup>61,26</sup>, GUILLERMO J. DAMKE<sup>16,64,65</sup>, JEREMY DARLING<sup>66</sup>, JAMES W. DAVIDSON JR.<sup>10</sup>, KYLE DAWSON<sup>20</sup>, ANNA BÁRBARA DE ANDRADE QUEIROZ<sup>59,60</sup>, MIGUEL ANGEL C. DE ICAZA LIZAOLA<sup>3</sup>, AXEL DE LA MACORRA<sup>67</sup>, SYLVAIN DE LA TORRE<sup>35</sup>, NATHAN DE LEE<sup>68,30</sup>, VICTORIA DE SAINTE AGATHE<sup>69</sup>, ALICE DECONTO MACHADO<sup>70,60</sup>, FLAVIA DELL'AGLI<sup>2,4</sup>, TIMOTHÉE DELUBAC<sup>71</sup>, ALEXANDAR M. DIAMOND-STANIC<sup>42</sup>, JOHN DONOR<sup>16</sup>, JUAN JOSÉ DOWNES<sup>72</sup>, NIV DROTH<sup>73</sup>, HÉLION DU MAS DES BOURBOUX<sup>13</sup>, CHRISTOPHER J. DUCKWORTH<sup>18</sup>, TOM DWELLY<sup>54</sup>, JAMIE DYER<sup>20</sup>, GARRETT EBELKE<sup>10</sup>, ARTHUR DAVIS EIGENBROT<sup>28</sup>, DANIEL J. EISENSTEIN<sup>74</sup>, YVONNE P. ELSWORTH<sup>75</sup>, ERIC ESMELLEM<sup>76</sup>, MIKE ERACLEOUS<sup>44</sup>, STEPHANIE ESCOFFIER<sup>62</sup>, XIAOHU FAN<sup>26</sup>, EMMA FERNÁNDEZ ALVAR<sup>3</sup>, J. G. FERNÁNDEZ-TRINCADO<sup>27</sup>, RAFAEL FERNANDO CIROLINI<sup>60</sup>, DIANE FEUILLET<sup>77</sup>, SCOTT W. FLEMING<sup>35</sup>, ANDREU FONT-RIBERA<sup>78</sup>, GORDON FREISCHLUD<sup>31</sup>, PETER FRINCHABOY<sup>16</sup>, YILEN GÓMEZ MAQUEO CHEW<sup>3</sup>, LLUÍS GALBANY<sup>9</sup>, ANA E. GARCÍA PÉREZ<sup>7</sup>, R. GARCIA-DIAS<sup>2,4</sup>, D. A. GARCÍA-HERNÁNDEZ<sup>2,4</sup>, PATRICK GAULME<sup>31</sup>, JOSEPH GELFAND<sup>34</sup>, HÉCTOR GIL-MARÍN<sup>79,80</sup>, BRUCE A. GILLESPIE<sup>31</sup>

1-ph.GA] 28 Jul 2017

We're very happy to report that the fourteenth data release of SDSS is now live!!! Go have a look at [www.sdss.org/dr14](http://www.sdss.org/dr14), and tell all your friends and colleagues that this awesome data set is now available for them to play with!

## Introduction: the eBOSS survey


- Apache Point Observatory (APO) 2.5-m telescope.
- SDSS-III project. 2009-2014 BOSS: Baryon Oscillation Spectroscopic Survey
- SDSS-IV project. 2014-2019 eBOSS: extended Baryon Oscillation Spectroscopic Survey
- 3 galaxy clustering programs (ELG, LRG, quasars) + Ly- $\alpha$
  
- new selection algorithms to identify redshift of galaxies
- BOSS had 99% success rate identifying redshifts of LRGs
- first eBOSS tests showed 70% of success rate on LRGs using same BOSS algorithms!



## Introduction: the eBOSS survey


eBOSS survey: **ELG**, **LRG** and **QSO** samples

- **LRG**  $0.6 < z < 1.0$ ;  $z_{\text{eff}} = 0.71$ .
- **ELG**  $0.6 < z < 1.1$ ;  $z_{\text{eff}} = 0.85$ .
- **QSO**  $0.8 < z < 2.2$ ;  $z_{\text{eff}} = 1.5$ .
- **Ly- $\alpha$**   $z_{\text{eff}} = 2.33$




[Credit : Anand Raichoor]

## Introduction: the eBOSS survey




# Introduction: the eBOSS survey




## Low Density of Quasars!

- Shot noise dominated covariance matrices
- (traditional) Reconstruction algorithms do not provide much signal gain



# Introduction: the eBOSS survey

- QSO DR14 area 2044 deg<sup>2</sup>.
- quasar range:  $0.8 < z < 2.2$   
( $z_{\text{eff}} \simeq 1.5$ )
- Number quasars: 147,000
- Low density:  $2 \times 10^{-5} [\text{Mpc}/h]^3$
- 3 disconnected areas: 2 SGC & NGC.



# Introduction: the eBOSS survey

## The clustering of the SDSS-IV extended Baryon Oscillation Spectroscopic Survey DR14 quasar sample: First measurement of Baryon Acoustic Oscillations between redshift 0.8 and 2.2

Metin Ata<sup>1</sup>, Falk Baumgarten<sup>1,2</sup>, Julian Bautista<sup>3</sup>, Florian Beutler<sup>4</sup>, Jonathan A. Blazek<sup>5</sup>, Jonathan Brinkmann<sup>6</sup>, Joel R. Brownstein<sup>7</sup>, Etienne Burtin<sup>7</sup>, Chia-Hsun Chung<sup>1</sup>, Johan Comparat<sup>8</sup>, Kyle S. Dawson<sup>3</sup>, Axel De La Macorra<sup>9</sup>, Wei Du<sup>10</sup>, Héctor Gil-Marín<sup>11,12</sup>, Julien Guy<sup>13</sup>, Nick Hand<sup>14</sup>, Shirley Ho<sup>15,16,17</sup>, Timothy A. Hutchinson<sup>3</sup>, Francisco-Shu Kitaura<sup>18,19</sup>, Jean-Paul Kneib<sup>20,21</sup>, Pierre Laurent<sup>7</sup>, Jean-Marc LeGoff<sup>7</sup>, Cameron K. McBride<sup>22</sup>, Joseph E. McEwen<sup>5</sup>, Eva-Maria Mueller<sup>7</sup>, Adam D. Myers<sup>3</sup>, Jeffrey A. Newman<sup>23</sup>, Isabelle Pâris<sup>21</sup>, Nathalie Palanque-Delabrouille<sup>7</sup>, Marcos Pellejero-Ibanez<sup>18,19</sup>, Will J. Percival<sup>4</sup>, Patrick Petitjean<sup>24</sup>, Francisco Prada<sup>25,26,27</sup>, Abhishek Prakash<sup>23</sup>, Sergio A. Rodríguez-Torres<sup>25,26,28</sup>, Ashley J. Ross<sup>5,4</sup>, Graziano Rossi<sup>29</sup>, Rossana Ruggeri<sup>4</sup>, Ariel G. Sánchez<sup>8</sup>, Siddharth Satpathy<sup>16,30</sup>, Donald P. Schneider<sup>31,32</sup>, Hee-Jong Seo<sup>33</sup>, Anže Slosar<sup>34</sup>, Alina Streblyanska<sup>18,19</sup>, Jeremy L. Tinker<sup>35</sup>, Rita Tojeiro<sup>36</sup>, Mariana Vargas Magaña<sup>9</sup>, M. Vivek<sup>3</sup>, Yuting Wang<sup>10,4</sup>, Christophe Yèche<sup>7,15</sup>, Liang Yu<sup>37</sup>, Pauline Zarrouk<sup>7</sup>, Cheng Zhao<sup>37</sup>, Gong-Bo Zhao<sup>10,4</sup>, Fangzhou Zhu<sup>38</sup>

### ABSTRACT

We present measurements of the Baryon Acoustic Oscillation (BAO) scale in redshift-space using the clustering of quasars. We consider a sample of 147,000 quasars from the extended Baryon Oscillation Spectroscopic Survey (eBOSS) distributed over 2044 square degrees with redshifts  $0.8 < z < 2.2$  and measure their spherically-averaged clustering in both configuration and Fourier space. Our observational dataset and the 1400 simulated realizations of our dataset allow us to detect a preference for BAO that is greater than  $2.5\sigma$ . We determine the spherically averaged BAO distance to  $z = 1.52$  to 4.4 per cent precision,  $D_V(z = 1.52) = 3855 \pm 170 r_{d, fid}$  Mpc. This is the first time the location of the BAO feature has been measured between redshifts 1 and 2. Our measurement is fully consistent with the prediction obtained by extrapolating the Planck flat  $\Lambda$ CDM best-fit cosmology. All of our results are consistent with basic large-scale structure (LSS) theory, confirming quasars to be a reliable tracer of LSS, and provide a starting point for numerous cosmological tests to be performed with eBOSS quasar samples.

**Key words:** cosmology: observations - (cosmology:) large-scale structure of Universe

### 1 INTRODUCTION

Using Baryon Acoustic Oscillations (BAOs) to measure the expansion of the Universe is now a mature field, with the BAO signal having been detected and measured to ever greater precision using data from a number of large galaxy surveys including: the Sloan Digital Sky Survey (SDSS) I and II (e.g., Eisenstein et al. 2005; Percival et al. 2010; Ross et al. 2015), the 2-degree Field Galaxy Redshift Survey (2dFGRS) (Percival et al. 2001; Cole et al. 2005), WiggleZ (Blake et al. 2011), the 6-degree Field Galaxy Survey (6dFGS) (Beutler et al. 2011), The Baryon Oscillation Spectroscopic Survey

ability in multi-epoch imaging from the Palomar Transient Factory. These selections are presented in Myers et al. (2015), alongside the characterisation of the final sample, as determined by the early data. The early data was observed as part of SEQUELS (The Sloan Extended QUasar, ELG and LRQ Survey, undertaken as part of SDSS-III and SDSS-IV), which acted as a pilot survey for eBOSS. SEQUELS used a broader quasar selection algorithm than that adopted for eBOSS, and a subsampled version of SEQUELS forms part of the eBOSS sample.


In this paper we present the first BAO measurements obtained

- Alphabetical paper Ata et al. 2017 submitted to the Journal arXiv:1705.06373
- First BAO measurement in  $0.8 \leq z \leq 2.2$
- $D_V = 3855 \pm 170 r_{d, fid}$  Mpc at  $z=1.52$  (4.4% precision)
- $\sim 2.5\sigma$  BAO significance
- DR14 Area: 2044  $\rightarrow$  4.4% precision
- DR16 Area: 5300  $\rightarrow$  2.7% precision
- eBOSS is working!

## Alcock-Paczynski effect

By assuming a wrong cosmological model ( $\Omega_m$ ) we change the line-of-sight clustering respect to the angular clustering, creating a measurable anisotropy: constrains  $H(z)$  and  $D_A(z)$  (or a combination of both).

$$d_{\text{comov}}(z) = \int_0^z \frac{cdz'}{H(z'; \Omega_m)}$$




The BAO scale is determined by the comoving sound horizon at recombination scale (standard ruler)

$$r_s = \frac{1}{H_0 \Omega_m^{1/2}} \int_0^{a^*} da \frac{c_s}{(a + a_{\text{eq}})^{1/2}}$$

# Baryonic Acoustic Oscillations

$$k_{\parallel} \rightarrow \alpha_{\parallel} k_{\parallel} \quad k_{\perp} \rightarrow \alpha_{\perp} k_{\perp}$$

$$\alpha_{\parallel} = \frac{H^{\text{fid}}(z)}{H(z)} \quad \alpha_{\perp} = \frac{D_A(z)}{D_A^{\text{fid}}(z)}$$



[Anderson et al. 2014,  
BOSS DR11]

- Surveys measure angles and redshifts, and these are affected by the assumed fiducial model.
- This changes the apparent/observed position of the BAO peak in the power spectrum differently in the radial and angular direction
- Isotropic Correlation Function / Power Spectrum sensitive to

$$D_V(z) = \left[ (1+z)^2 D_A^2(z) \frac{cz}{H(z)} \right]^{1/3}$$

# Methodology



- We perform two complementary and independent analyses on the same dataset using the *i)* **Power Spectrum** and *ii)* **Correlation Function**.
- Both observables should contain the same amount of information, but in practice are affected differently by noise and systematic effects.
- We use mocks (1000 EZ mocks) and (400 QPM mocks) to estimate the covariance matrices and perform systematic tests.
- We model the broadband shape of the PS/CF phenomenologically and the BAO as linear+damping ( $\Sigma_{\text{nl}}$ )

$$P(k, \alpha) = P_{\text{sm}}(k) \left\{ 1 + [\mathcal{O}_{\text{lin}}(k/\alpha) - 1] e^{-\frac{1}{2} \Sigma_{\text{nl}}^2 k^2} \right\}$$

smooth PS:  $P_{\text{sm}}(k) \equiv B^2 P_{\text{nw}}^{\text{lin}}(k) + A_1 k + A_2 + A_3/k$

## Measurement



## DR14 QSO isotropic Power spectrum and Correlation function



Actual data + diagonal errors from EZ mocks.  
Dashed lines mean of the EZ-mocks



# Measurement

## DR14 QSO isotropic Power spectrum and Correlation function



Actual data + diagonal errors from EZ mocks.  
Solid lines best-fit model

# Tests on Mocks



Correlation coefficient between PS and CF  $\rho = 0.97$ .

Data is a very typical case of mocks (both in measurement and error).

# Tests on Mocks

case	$\alpha - \alpha_{\text{exp}}$
EZ mocks:	
$\xi(s)$ :	
fiducial	$0.0023 \pm 0.0016$
$5h^{-1}\text{Mpc}$	$0.0027 \pm 0.0016$
$P(k)$ :	
fiducial	$0.0019 \pm 0.0017$
$k_{\text{max}} = 0.30 h\text{Mpc}^{-1}$	$0.0009 \pm 0.0017$
$\Sigma_{\text{nl}} = [6 \pm 3] h^{-1}\text{Mpc}$	$0.0021 \pm 0.0016$
$\Sigma_{\text{nl}} = [6 \pm 3] h^{-1}\text{Mpc} \ \& \ k_{\text{max}} = 0.30$	$0.0011 \pm 0.0016$
logk - binning	$0.0032 \pm 0.0017$
logk - binning & $k_{\text{max}} = 0.30 h\text{Mpc}^{-1}$	$0.0022 \pm 0.0016$
$A_4, A_5$ terms	$0.0037 \pm 0.0017$
QPM mocks:	
$\xi(s)$ :	
fiducial	$0.0017 \pm 0.0028$
$5h^{-1}\text{Mpc}$	$0.0027 \pm 0.0028$
QPM cov	$0.0023 \pm 0.0026$
$P(k)$ :	
fiducial	$0.0017 \pm 0.0027$
QPM cov	$0.0012 \pm 0.0026$


case (+bin shift)	$\langle \alpha \rangle$	$\langle \sigma \rangle$	$S$	$N_{\text{det}}/N_{\text{tot}}$
EZ mocks:				
<b>consensus</b> $P(k) + \xi(s)$	1.003	0.050	0.050	944/1000
$\xi(s)$ :				
combined	1.003	0.049	0.049	939/1000
fiducial	1.002	0.048	0.050	932/1000
+2	1.002	0.049	0.050	928/1000
+4	1.002	0.048	0.050	938/1000
+6	1.003	0.048	0.051	929/1000
$5h^{-1}\text{Mpc}$	1.003	0.049	0.050	937/1000
$P(k)$ :				
combined	1.002	0.052	0.050	941/1000
fiducial	1.002	0.051	0.051	929/1000
+1/4	1.001	0.052	0.050	931/1000
+2/4	1.004	0.051	0.049	935/1000
+3/4	1.001	0.052	0.050	937/1000
logk - binning	1.002	0.051	0.050	927/1000
$k_{\text{max}} = 0.30 h\text{Mpc}^{-1}$	1.002	0.051	0.051	934/1000
QPM mocks:				
$\xi(s)$ :				
fiducial	1.001	0.051	0.052	361/400
$5h^{-1}\text{Mpc}$	1.000	0.050	0.051	355/400
QPM cov	1.002	0.051	0.052	369/400
$P(k)$ :				
fiducial	0.998	0.049	0.051	354/400
QPM cov	0.999	0.049	0.049	359/400

NL effects shift the BAO peak to higher  $\alpha$ .  
 $\sim 0.1\%$  effect on measured  $\alpha$ . Negligible on data!

# Tests on data



case	$\alpha$	$\chi^2/\text{dof}$
<b>DR14 Measurement</b> $P(k) + \xi(s)$	$0.996 \pm 0.044$	-
$\xi(s)$ (combined)	$0.991 \pm 0.041$	9.7/13
$P(k)$ (combined)	$1.004 \pm 0.047$	26.5/33
<b>Robustness tests</b>		
$\xi(s)$ :		
$Z_{\text{PCA}}$ (combined)	$0.992 \pm 0.045$	15.6/13
fiducial	$0.997 \pm 0.044$	7.1/13
+2	$1.001 \pm 0.047$	12.9/13
+4	$0.978 \pm 0.035$	7.4/13
+6	$0.996 \pm 0.041$	11.1/13
NGC	$0.971 \pm 0.056$	7.7/13
SGC	$1.027 \pm 0.063$	17.4/13
QPM cov	$0.994 \pm 0.043$	6.8/13
$\Delta s = 5h^{-1}\text{Mpc}$	$0.993 \pm 0.040$	18.7/24
no $w_{\text{sys}}$	$0.998 \pm 0.047$	5.1/13
$50 < s < 150h^{-1}\text{Mpc}$	$0.998 \pm 0.048$	4.8/8
$\Sigma_{\text{nl}} = 3.0h^{-1}\text{Mpc}$	$0.994 \pm 0.043$	7.2/13
$\Sigma_{\text{nl}} = 9.0h^{-1}\text{Mpc}$	$1.001 \pm 0.048$	7.3/13
$A_n = 0$	$1.000 \pm 0.044$	7.3/16
no $B$ prior	$0.998 \pm 0.043$	6.9/13
$P(k)$ :		
$Z_{\text{PCA}}$ (combined)	$1.005 \pm 0.045$	27.6/33
fiducial	$1.002 \pm 0.046$	27.7/33
+1/4	$0.994 \pm 0.044$	24.1/33
+2/4	$0.993 \pm 0.046$	27.3/33
+3/4	$1.009 \pm 0.050$	26.9/33
NGC	$0.977 \pm 0.060$	17.5/16
SGC	$1.029 \pm 0.067$	9.9/16
QPM cov	$1.014 \pm 0.045$	27.3/33
log $k$ - binning	$1.005 \pm 0.046$	30.4/39
log $k$ - binning, $k_{\text{max}} = 0.30 h\text{Mpc}^{-1}$	$1.011 \pm 0.047$	34.9/45
no $w_{\text{sys}}$	$1.003 \pm 0.052$	27.9/33
$k_{\text{max}} = 0.30 h\text{Mpc}^{-1}$	$1.011 \pm 0.048$	44.5/47
$\Sigma_{\text{nl}} = 3 h^{-1}\text{Mpc}$	$1.001 \pm 0.041$	27.5/33
$\Sigma_{\text{nl}} = 9 h^{-1}\text{Mpc}$	$1.008 \pm 0.054$	28.2/33
$\Sigma_{\text{nl}} = [6 \pm 3] h^{-1}\text{Mpc}$	$1.002 \pm 0.046$	27.6/32

Both  $P(k)$  and  $\xi(s)$  measurements are very consistent!



# Cosmology

- Fully consistent with  $\Lambda$ CDM + Planck.
- $\text{Ly-}\alpha$  2.5 measurement dominates at high  $z$ .
- Further analyses will focus on redshift weighting schemes.



# BAO Summary

## Summary,

- We have a robust BAO isotropic measurement at  $z \simeq 1.5$ .  
with 4.4% precision:  $D_V(1.52) = 3855 \pm 170 r_d/r_{d,\text{fid}}$  Mpc.
- Very consistent with mocks and between  $P$  and  $\xi$  observables.
- Fully consistent with LCDM+Planck

# BAO Summary

## Summary,


- We have a robust BAO isotropic measurement at  $z \simeq 1.5$ .  
with 4.4% precision:  $D_V(1.52) = 3855 \pm 170 r_d/r_{d,\text{fid}}$  Mpc.
- Very consistent with mocks and between  $P$  and  $\xi$  observables.
- Fully consistent with LCDM+Planck
- First BAO science result from WG, but not last one. More complex BAO analyses will be done in the forthcoming months (weighting  $z$  evolution).
- Future quasar releases (DR16) will focus on the anisotropic quasar BAO  $\rightarrow$  Constrain  $D_A$  and  $H$ .
- Further information on  $D_V$  can be extracted from RSD analysis (several papers to be released this Fall)

The main difficulty when performing RSD analyses is controlling the observational systematic effects.

Such effects have minor effect on the position of the BAO (given the current errorbars), but they turn to be more important for RSD analyses,


- Failure rate as a function of the plate position.
- Spectroscopic errors.
- Estimate of the quasar redshift using different algorithms.

Failure rate in NGC





$\sim 10\%$  failure rate

Failure rate in NGC



$\sim 3.5\%$  failure rate

## Why do the quasars always fail at the same plate positions?



[Credit: P. Zarrouk]

Even with  $\sim 3.5\%$  failure rate, applying the nearest angular neighbour reduces the amplitude of the monopole and enhances the quadrupole signal significantly.

As a consequence,  $f\sigma_8$  is affected by  $\sim 0.7\sigma$ .

BAO not affected

	$\Delta\alpha_{\text{iso}}$	$S_\alpha$	$\Delta f\sigma_8$	$S_{f\sigma_8}$	$N_{\text{det}}$
EZ $\langle x \rangle_i$	$-1.64 \pm 0.13$	-	$-1.43 \pm 0.16$	-	-
EZ $\langle x_i \rangle$	$-1.6 \pm 4.4$	4.8	$-1.4 \pm 4.9$	5.5	979
EZ $\langle x \rangle_i w_{\text{col}}$	$-1.90 \pm 0.13$	-	$2.24 \pm 0.17$	-	-
EZ $\langle x_i \rangle w_{\text{col}}$	$-1.7 \pm 4.4$	4.6	$2.4 \pm 5.4$	5.4	959

(units of the table in  $10^{-2}$ ).

Better correction than just up-weight the nearest neighbour is needed!

## Quasar Clustering Working Group Report

Current RSD & BAO projects DR14 QSO sample

- Gil-Marín et al. *Fourier Space Multipoles RSD*
- Hou et al. *Configuration Space Wedges RSD*
- Ruggeri et al. *Fourier Space Multipoles with  $z$ -weights RSD*
- Wang et al. *Fourier Space Multipoles with  $z$ -weights BAO*
- Zarrouk et al. *Configuration Space Multipoles RSD*
- Zhao et al. *Fourier Space Multipoles with  $z$ -weights RSD*
- Zhu et al. *Configuration Space Multipoles with  $z$ -weights BAO*

**WG is currently focused on**

- Producing reliable mocks for covariance matrix
- Building appropriate weighting scheme for correcting redshift failures and close pairs.

Thank you!

