

No-go for leptoquarks from pseudo-Riemannian structure in NCG

Arkadiusz Bochniak¹

¹ Institute of Physics of the Jagiellonian University

Benasque, 24.09.2018



Bochniak A., Sitarz A., *Finite Pseudo-Riemannian spectral triples and The Standard Model*, Phys. Rev. D **97** 115029 (2018)

Author acknowledge support by NCN grant OPUS 2016/21/B/ST1/02438

What is a geometry ?

Connes' reconstruction theorem

The whole metric and spin structure of a compact, orientable, Riemannian, spin^c manifold can be encoded in the $*$ -algebra $C^\infty(M)$ of smooth functions, Hilbert space $L^2(S)$ of square-integrable spinors and the Dirac operator $\mathcal{D}_M = i\gamma^\mu (\partial_\mu + \omega_\mu)$ together with the γ_5 grading and the charge conjugation operator.

What is a geometry ?

Connes' reconstruction theorem

The whole metric and spin structure of a compact, orientable, Riemannian, spin^c manifold can be encoded in the $*$ -algebra $C^\infty(M)$ of smooth functions, Hilbert space $L^2(S)$ of square-integrable spinors and the Dirac operator $\not{D}_M = i\gamma^\mu (\partial_\mu + \omega_\mu)$ together with the γ_5 grading and the charge conjugation operator.

Spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D}, \gamma, J)$

\mathcal{A} is a $*$ -algebra represented on Hilbert space \mathcal{H} , $\gamma = \gamma^\dagger$, $\gamma^2 = 1$ is a $\mathbb{Z}/2\mathbb{Z}$ -grading commuting with \mathcal{A} , J is an antilinear isometry s.th. $[Ja^*J^{-1}, b] = 0$ for all $a, b \in \mathcal{A}$.

\mathcal{D} is essentially self-adjoint operator with compact resolvent and s.th. $[\mathcal{D}, a]$ is bounded for all $a \in \text{Dom}(\mathcal{D})$ and $\mathcal{D}\gamma = -\gamma\mathcal{D}$.

Moreover $\mathcal{D}J = \epsilon J\mathcal{D}$, $J^2 = \epsilon'\text{id}$ and $J\gamma = \epsilon''\gamma J$ with $\epsilon, \epsilon', \epsilon'' = \pm 1$ defining KO -dimension.

There are additional compatibility conditions for \mathcal{D} and for γ .

Almost-commutative geometry for the Standard Model

$$(C^\infty(M) \otimes (\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})), L^2(S) \otimes H_f, \not{D}_M \otimes 1 + \gamma_5 \otimes D_f, \gamma_5 \otimes \gamma_f, J_M \otimes J_f)$$

$$H_f = H_L \oplus H_R \oplus H_L^c \oplus H_R^c$$

$$D_f \in M_{96}(\mathbb{C})$$

γ_f - chirality operator

J_f - exchange particle with antiparticle and complex conjugates

Almost-commutative geometry for the Standard Model

$$(C^\infty(M) \otimes (\mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})), L^2(S) \otimes H_f, \not{D}_M \otimes 1 + \gamma_5 \otimes D_f, \gamma_5 \otimes \gamma_f, J_M \otimes J_f)$$

$$H_f = H_L \oplus H_R \oplus H_L^c \oplus H_R^c$$

$$D_f \in M_{96}(\mathbb{C})$$

γ_f - chirality operator

J_f - exchange particle with antiparticle and complex conjugates

Expansion of the Euclidean spectral action reproduces the effective action for the SM and allows for the expression of bosonic parameters by fermionic one.

Pseudo-Riemannian structure ?

Finite pseudo-Riemannian spectral triple of signature (p, q)

$$(\mathcal{A}, \mathcal{H}, \mathcal{D}, \gamma, J, \beta)$$

1. \mathcal{A} is a $*$ -algebra represented on an Hilbert space \mathcal{H}
2. For $p + q$ even $\gamma^* = \gamma$, $\gamma^2 = 1$ is a $\mathbb{Z}/2\mathbb{Z}$ -grading commuting with \mathcal{A}
3. J is antilinear isometry with $[Ja^*J^{-1}, b] = 0$
4. $\beta = \beta^\dagger$, $\beta^2 = 1$ commuting with \mathcal{A}
5. $\mathcal{D}^\dagger = (-1)^p \beta \mathcal{D} \beta$
6. $[\mathcal{D}, a]$ is bounded
7. $\mathcal{D}\gamma = -\gamma\mathcal{D}$
8. $\mathcal{D}J = \epsilon J\mathcal{D}$, $J^2 = \epsilon' \text{id}$, $J\gamma = \epsilon'' \gamma J$

$p - q \bmod 8$	0	1	2	3	4	5	6	7
ϵ	+	-	+	+	+	-	+	+
ϵ'	+	+	-	-	-	-	+	+
ϵ''	+		-		+		-	

Finite pseudo-Riemannian spectral triple of signature (p, q)

9. $\beta\gamma = (-1)^p\gamma\beta$, $\beta J = (-1)^{\frac{p(p-1)}{2}}\epsilon^p J\beta$
10. $[JaJ^{-1}, [\mathcal{D}, b]] = 0$
11. orientability : there exist $\mathcal{A} \ni a^i, a_0^i, \dots, a_n^i$, $i = 1, \dots, k$ s.th.

$$\sum_{i=1}^k Ja^i J^{-1} a_0^i [\mathcal{D}, a_1^i] \dots [\mathcal{D}, a_n^i] = \begin{cases} \gamma, & n \text{ even} \\ 1, & n \text{ odd} \end{cases}$$

12. time-orientation : there exist $\mathcal{A} \ni b^i, b_0^i, \dots, b_p^i$, $i = 1, \dots, k'$ s.th.

$$\beta = \sum_{i=1}^{k'} Jb^i J^{-1} b_0^i [\mathcal{D}, b_1^i] \dots [\mathcal{D}, b_p^i].$$

Clifford algebra : $\gamma_a \gamma_b + \gamma_b \gamma_a = 2\eta_{ab} 1$

- $\gamma = i^{\frac{p-q}{2}} \gamma_1 \dots \gamma_{p+q}$
- there exists unitary B s.th. $B\gamma_i = \epsilon \gamma_i^* B$ and $BB^* = \epsilon'$. Define $J\psi := B\psi^*$.
- $\mathcal{D} = -\sum_j \eta_{jj} \gamma_j \partial_j$
- $B\gamma = \epsilon'' \gamma B$
- $\beta = i^{\frac{1}{2}p(p-1)} \gamma_1 \dots \gamma_p$
- $\beta \mathcal{D} \beta = (-1)^p \mathcal{D}^\dagger$

Riemannian from pseudo-Riemannian

$$\mathcal{D}_+ = \frac{1}{2}(\mathcal{D} + \mathcal{D}^\dagger), \quad \mathcal{D}_- = \frac{i}{2}(\mathcal{D} - \mathcal{D}^\dagger)$$

We get two Riemannian spectral triples $(\mathcal{A}, \pi, \mathcal{H}, \mathcal{D}_\pm, J, \gamma)$, that differ by KO -dimensions, with additional selfadjoint grading β s.th.

$$\beta \mathcal{D}_\pm = \pm(-1)^p \mathcal{D}_\pm \beta,$$

$$\beta \gamma = (-1)^p \gamma \beta, \quad \beta J = (-1)^{\frac{1}{2}p(p-1)} \epsilon^p J \beta.$$

Riemannian from pseudo-Riemannian

$$\mathcal{D}_+ = \frac{1}{2}(\mathcal{D} + \mathcal{D}^\dagger), \quad \mathcal{D}_- = \frac{i}{2}(\mathcal{D} - \mathcal{D}^\dagger)$$

We get two Riemannian spectral triples $(\mathcal{A}, \pi, \mathcal{H}, \mathcal{D}_\pm, J, \gamma)$, that differ by KO -dimensions, with additional selfadjoint grading β s.th.

$$\beta \mathcal{D}_\pm = \pm(-1)^p \mathcal{D}_\pm \beta,$$

$$\beta \gamma = (-1)^p \gamma \beta, \quad \beta J = (-1)^{\frac{1}{2}p(p-1)} \epsilon^p J \beta.$$

$$\mathcal{D}_E = \mathcal{D}_+ + \mathcal{D}_-$$

$$J_E = J\beta, \quad \text{or} \quad J_E = J\beta\gamma$$

$(\mathcal{A}, \pi, \mathcal{H}, \mathcal{D}_E, J_E, \gamma)$ is a Riemannian spectral triple of signature $(0, -(p+q))$.

Example : Functions over 2-point space

Take $A = \mathbb{C}^2$, $H = \bigoplus_{i,j} H_{i,j}$ with $H_{i,j} = \mathbb{C}$ for $i, j = 1, 2$.

Example : Functions over 2-point space

Take $A = \mathbb{C}^2$, $H = \bigoplus_{i,j} H_{i,j}$ with $H_{i,j} = \mathbb{C}$ for $i, j = 1, 2$. We define J , γ as matrices:

$$\gamma = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \circ *.$$

Example : Functions over 2-point space

Take $A = \mathbb{C}^2$, $H = \bigoplus_{i,j} H_{i,j}$ with $H_{i,j} = \mathbb{C}$ for $i, j = 1, 2$. We define J , γ as matrices:

$$\gamma = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \circ *.$$

Now, we can easily identify a nontrivial additional symmetry ($\mathbb{Z}/2\mathbb{Z}$ -grading) β and construct a Dirac operator \mathcal{D}_+ , which is real, satisfies first-order condition and commutes with β :

$$\beta = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad \mathcal{D}_+ = \begin{pmatrix} 0 & d & d^* & 0 \\ d^* & 0 & 0 & 0 \\ d & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

It gives us a Riemannian real spectral triple of KO -dimension 0 with an additional symmetry.

It gives us a Riemannian real spectral triple of KO -dimension 0 with an additional symmetry.

The restriction due to the β -symmetry gives only one free parameter into the family of possible Dirac operators.

It gives us a Riemannian real spectral triple of KO -dimension 0 with an additional symmetry.

The restriction due to the β -symmetry gives only one free parameter into the family of possible Dirac operators.

Similarly we construct \mathcal{D}_- , which satisfies $\mathcal{D}_-\beta = -\beta\mathcal{D}_-$.

It gives us a Riemannian real spectral triple of KO -dimension 0 with an additional symmetry.

The restriction due to the β -symmetry gives only one free parameter into the family of possible Dirac operators.

Similarly we construct \mathcal{D}_- , which satisfies $\mathcal{D}_-\beta = -\beta\mathcal{D}_-$.

Both these triples could be seen as Riemannian parts of a pseudo-Riemannian spectral triple with signature $(4, 4)$ or $(0, 0)$.

It gives us a Riemannian real spectral triple of KO -dimension 0 with an additional symmetry.

The restriction due to the β -symmetry gives only one free parameter into the family of possible Dirac operators.

Similarly we construct \mathcal{D}_- , which satisfies $\mathcal{D}_-\beta = -\beta\mathcal{D}_-$.

Both these triples could be seen as Riemannian parts of a pseudo-Riemannian spectral triple with signature $(4, 4)$ or $(0, 0)$.

The full Dirac operator \mathcal{D} :

$$\mathcal{D} = \begin{pmatrix} 0 & d & d^* & 0 \\ d^* & 0 & 0 & c \\ d & 0 & 0 & c^* \\ 0 & -c^* & -c & 0 \end{pmatrix},$$

where c, d are arbitrary complex numbers.

The Standard Model

$$A_f = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}), \quad H_f = (H_l \oplus H_q) \oplus (H_{\bar{l}} \oplus H_{\bar{q}})$$

$$H_l = \langle \{\nu_R, e_R, (\nu_L, e_L)\} \rangle$$

$$H_q = \langle \{u_R, d_R, (u_L, d_L)\}_{c=1,2,3} \rangle$$

The Standard Model

$$A_f = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}), \quad H_f = (H_l \oplus H_q) \oplus (H_{\bar{l}} \oplus H_{\bar{q}})$$

$$H_l = \langle \{\nu_R, e_R, (\nu_L, e_L)\} \rangle$$

$$H_q = \langle \{u_R, d_R, (u_L, d_L)\}_{c=1,2,3} \rangle$$

$$\pi(\lambda, h, m) = \lambda \oplus \bar{\lambda} \oplus h \text{ on } H_l \text{ and } H_q$$

$$\pi(\lambda, h, m) = \bar{\lambda} \text{ on } H_{\bar{l}} \text{ and } 1_4 \otimes m \text{ on } H_{\bar{q}}$$

$$D_f = \begin{pmatrix} S & T^\dagger \\ T & \bar{S} \end{pmatrix}, \quad S = \begin{pmatrix} S_l & \\ & S_q \otimes 1_3 \end{pmatrix}$$

$$T\nu_R = Y_R\bar{\nu}_R$$

The Standard Model

$$S_l = \begin{bmatrix} & & Y_\nu^\dagger & \\ & & & Y_e^\dagger \\ Y_\nu & & & \\ & Y_e & & \end{bmatrix}, \quad S_q = \begin{bmatrix} & & Y_u^\dagger & \\ & & & Y_d^\dagger \\ Y_u & & & \\ & Y_d & & \end{bmatrix}$$

γ_f - chirality grading

J_f - real structure

The Standard Model

$$S_l = \begin{bmatrix} & & Y_\nu^\dagger & \\ & & & Y_e^\dagger \\ Y_\nu & & & \\ & Y_e & & \end{bmatrix}, \quad S_q = \begin{bmatrix} & & Y_u^\dagger & \\ & & & Y_d^\dagger \\ Y_u & & & \\ & Y_d & & \end{bmatrix}$$

γ_f - chirality grading
 J_f - real structure

- The existence of right neutrinos implies nonorientability of the geometry
- It is well known that the above Dirac operator is not unique within the model-building scheme of noncommutative geometry. Even the introduction of more constraints, like the second-order condition or Hodge-duality does not allow to exclude the terms, which would introduce the couplings between lepton and quarks and lead to the leptoquark fields

The Standard Model

There exists 0-cycle

$$\beta = \pi(1, 1, -1)J_F\pi(1, 1, -1)J_F^{-1}$$

that is a $\mathbb{Z}/2\mathbb{Z}$ -grading which distinguish between leptons and quarks.

The Standard Model

There exists 0-cycle

$$\beta = \pi(1, 1, -1)J_F\pi(1, 1, -1)J_F^{-1}$$

that is a $\mathbb{Z}/2\mathbb{Z}$ -grading which distinguish between leptons and quarks.

Moreover, this β makes the geometry to be a finite pseudo-Riemannian spectral triple of signature $(4k, 4k + 2 \pmod{8})$ with $k \in \mathbb{N}$.

The Standard Model

There exists 0-cycle

$$\beta = \pi(1, 1, -1)J_F\pi(1, 1, -1)J_F^{-1}$$

that is a $\mathbb{Z}/2\mathbb{Z}$ -grading which distinguish between leptons and quarks.

Moreover, this β makes the geometry to be a finite pseudo-Riemannian spectral triple of signature $(4k, 4k + 2 \pmod{8})$ with $k \in \mathbb{N}$.

$(A_f, H_f, D_f, \gamma_f, J_f, \beta)$ could be seen as a Riemannian restriction of a real even pseudo-Riemannian spectral triple of signature $(0, 2)$.

Possible pseudo-Riemannian structures for the Standard Model

Take as a Hilbert space $H \cong F \oplus F^*$ with

$$F \ni v = \begin{bmatrix} \nu_R & u_R^1 & u_R^2 & u_R^3 \\ e_R & d_R^1 & d_R^2 & d_R^3 \\ \nu_L & u_L^1 & u_L^2 & u_L^3 \\ e_L & d_L^1 & d_L^2 & d_L^3 \end{bmatrix} \in M_4(\mathbb{C}).$$

Vectors from H can be represented as $\begin{bmatrix} v \\ w \end{bmatrix}$, with $v, w \in M_4(\mathbb{C})$.

Possible pseudo-Riemannian structures for the Standard Model

Take as a Hilbert space $H \cong F \oplus F^*$ with

$$F \ni v = \begin{bmatrix} \nu_R & u_R^1 & u_R^2 & u_R^3 \\ e_R & d_R^1 & d_R^2 & d_R^3 \\ \nu_L & u_L^1 & u_L^2 & u_L^3 \\ e_L & d_L^1 & d_L^2 & d_L^3 \end{bmatrix} \in M_4(\mathbb{C}).$$

Vectors from H can be represented as $\begin{bmatrix} v \\ w \end{bmatrix}$, with $v, w \in M_4(\mathbb{C})$. The real structure is given by

$$J \begin{bmatrix} v \\ w \end{bmatrix} = \begin{bmatrix} w^* \\ v^* \end{bmatrix}.$$

Possible pseudo-Riemannian structures for the Standard Model

Take as a Hilbert space $H \cong F \oplus F^*$ with

$$F \ni v = \begin{bmatrix} \nu_R & u_R^1 & u_R^2 & u_R^3 \\ e_R & d_R^1 & d_R^2 & d_R^3 \\ \nu_L & u_L^1 & u_L^2 & u_L^3 \\ e_L & d_L^1 & d_L^2 & d_L^3 \end{bmatrix} \in M_4(\mathbb{C}).$$

Vectors from H can be represented as $\begin{bmatrix} v \\ w \end{bmatrix}$, with $v, w \in M_4(\mathbb{C})$. The real structure is given by

$$J \begin{bmatrix} v \\ w \end{bmatrix} = \begin{bmatrix} w^* \\ v^* \end{bmatrix}.$$

We can identify $\text{End}_{\mathbb{C}}(H)$ with $M_4(\mathbb{C}) \otimes M_2(\mathbb{C}) \otimes M_4(\mathbb{C})$ and denote by e_{ij} a matrix with the 1 in position (i, j) and zero everywhere else.

Possible pseudo-Riemannian structures for the Standard Model

Elements of the algebra $A = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ are represented by

$$\left[\begin{array}{cc|c} \lambda & \bar{\lambda} & 0 \\ \hline 0 & & q \end{array} \right] \otimes e_{11} \otimes 1 + \left[\begin{array}{cc|c} \lambda & & 0 \\ \hline 0 & & m \end{array} \right] \otimes e_{22} \otimes 1,$$

where $\lambda \in \mathbb{C}$, $q \in \mathbb{H}$ and $m \in M_3(\mathbb{C})$.

Possible pseudo-Riemannian structures for the Standard Model

Elements of the algebra $A = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ are represented by

$$\left[\begin{array}{cc|c} \lambda & \bar{\lambda} & 0 \\ \hline 0 & & q \end{array} \right] \otimes e_{11} \otimes 1 + \left[\begin{array}{cc|c} \lambda & & 0 \\ \hline 0 & & m \end{array} \right] \otimes e_{22} \otimes 1,$$

where $\lambda \in \mathbb{C}$, $q \in \mathbb{H}$ and $m \in M_3(\mathbb{C})$. The grading is of the form

$$\gamma = \begin{bmatrix} 1_2 & \\ & -1_2 \end{bmatrix} \otimes e_{11} \otimes 1 + 1 \otimes e_{22} \otimes \begin{bmatrix} -1_2 & \\ & 1_2 \end{bmatrix}.$$

Possible pseudo-Riemannian structures for the Standard Model

Elements of the algebra $A = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$ are represented by

$$\left[\begin{array}{c|c} \lambda & 0 \\ \hline \bar{\lambda} & q \end{array} \right] \otimes e_{11} \otimes 1 + \left[\begin{array}{c|c} \lambda & 0 \\ \hline 0 & m \end{array} \right] \otimes e_{22} \otimes 1,$$

where $\lambda \in \mathbb{C}$, $q \in \mathbb{H}$ and $m \in M_3(\mathbb{C})$. The grading is of the form

$$\gamma = \begin{bmatrix} 1_2 & \\ & -1_2 \end{bmatrix} \otimes e_{11} \otimes 1 + 1 \otimes e_{22} \otimes \begin{bmatrix} -1_2 & \\ & 1_2 \end{bmatrix}.$$

The Dirac operator is of the form

$$D = D_0 + D_1,$$

where $D_1 = JD_0J^{-1}$.

Possible pseudo-Riemannian structures for the Standard Model

We would like to have a spectral triple of KO -dimension 6, with a selfadjoint Dirac operator, but such that commutes with a suitable β that represents the shadow of a pseudo-Riemannian structure.

Possible pseudo-Riemannian structures for the Standard Model

We would like to have a spectral triple of KO -dimension 6, with a selfadjoint Dirac operator, but such that commutes with a suitable β that represents the shadow of a pseudo-Riemannian structure.

Let us now take the general form of a Dirac operator that satisfies an order-one condition. We have

$$D_0 = \begin{bmatrix} & M \\ M^\dagger & \end{bmatrix} \otimes e_{11} \otimes e_{11} + \begin{bmatrix} & N \\ N^\dagger & \end{bmatrix} \otimes e_{11} \otimes (1 - e_{11}) + \\ + \begin{bmatrix} A & B \\ 0 & 0 \end{bmatrix} \otimes e_{12} \otimes e_{11} + \begin{bmatrix} A^\dagger & 0 \\ B^\dagger & 0 \end{bmatrix} \otimes e_{21} \otimes e_{11},$$

where M, N, A, B are 2×2 complex matrices.

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1) J \pi(\lambda_2, q_2, m_2) J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1)J\pi(\lambda_2, q_2, m_2)J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Up to the trivial rescaling (by -1) we have three possibilities.

- $\pi(1, 1, -1)$
- $\pi(1, -1, 1)$
- $\pi(-1, 1, 1)$

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1)J\pi(\lambda_2, q_2, m_2)J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Up to the trivial rescaling (by -1) we have three possibilities.

- $\pi(1, 1, -1)$
- $\pi(1, -1, 1)$
- $\pi(-1, 1, 1)$

For the case $\beta = \pi(1, -1, 1)J\pi(1, -1, 1)J^{-1}$ the restrictions for the Dirac operator are $M = N = 0$ and no restriction for A, B .

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1)J\pi(\lambda_2, q_2, m_2)J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Up to the trivial rescaling (by -1) we have three possibilities.

- $\pi(1, 1, -1)$
- $\pi(1, -1, 1)$
- $\pi(-1, 1, 1)$

For the case $\beta = \pi(1, -1, 1)J\pi(1, -1, 1)J^{-1}$ the restrictions for the Dirac operator are $M = N = 0$ and no restriction for A, B . Furthermore, if

$\beta = \pi(-1, 1, 1)J\pi(-1, 1, 1)J^{-1}$ then again $M, N, B = 0$ and A has to satisfy $A = A \cdot \text{diag}(1, -1)$.

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1)J\pi(\lambda_2, q_2, m_2)J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Up to the trivial rescaling (by -1) we have three possibilities.

- $\pi(1, 1, -1)$
- $\pi(1, -1, 1)$
- $\pi(-1, 1, 1)$

For the case $\beta = \pi(1, -1, 1)J\pi(1, -1, 1)J^{-1}$ the restrictions for the Dirac operator are $M = N = 0$ and no restriction for A, B . Furthermore, if

$\beta = \pi(-1, 1, 1)J\pi(-1, 1, 1)J^{-1}$ then again $M, N, B = 0$ and A has to satisfy $A = A \cdot \text{diag}(1, -1)$. It is worth noting that both of these restrictions lead not only to unphysical Dirac operators that do not break the electroweak symmetry but also do not satisfy the Hodge duality.

Possible pseudo-Riemannian structures for the Standard Model

We look for a β that is a 0-cycle, i.e. a sum of elements of the form

$$\beta = \pi(\lambda_1, q_1, m_1)J\pi(\lambda_2, q_2, m_2)J^{-1},$$

with $\lambda_1, \lambda_2 \in \mathbb{C}$, $q_1, q_2 \in \mathbb{H}$, $m_1, m_2 \in M_3(\mathbb{C})$.

Up to the trivial rescaling (by -1) we have three possibilities.

- $\pi(1, 1, -1)$
- $\pi(1, -1, 1)$
- $\pi(-1, 1, 1)$

For the case $\beta = \pi(1, -1, 1)J\pi(1, -1, 1)J^{-1}$ the restrictions for the Dirac operator are $M = N = 0$ and no restriction for A, B . Furthermore, if

$\beta = \pi(-1, 1, 1)J\pi(-1, 1, 1)J^{-1}$ then again $M, N, B = 0$ and A has to satisfy $A = A \cdot \text{diag}(1, -1)$. It is worth noting that both of these restrictions lead not only to unphysical Dirac operators that do not break the electroweak symmetry but also do not satisfy the Hodge duality.

Finally, with the $\beta = \pi(1, 1, -1)J\pi(1, 1, -1)J^{-1}$ we have no restriction whatsoever for M, N while then $B = 0$ and A needs to satisfy: $A = A \cdot \text{diag}(1, -1)$. That leaves the possibility that A_{11} and A_{21} coefficients are present, providing no significant physical effects, and in particular leading only to terms involving a sterile neutrino.

- We proposed new definition of the finite pseudo-Riemannian spectral triples
- We proposed an alternative explanation of the observed quarks-leptons symmetry which prevents the $SU(3)$ -breaking, as a shadow of the pseudo-Riemannian structure
- We proposed that the consistent model-building for the physical interactions and possible extensions of the Standard Model within the noncommutative geometry framework should use possibly the pseudo-Riemannian extension of finite spectral triples. We demonstrated that the pseudo-Riemannian framework allows for more restrictions and, in the discussed case introduces an extra symmetry grading, which we interpreted as the lepton-quark symmetry