


The Electroweak Standard Model



José Ignacio Illana



UNIVERSIDAD
DE GRANADA

Outline



Exercises

tools

Kinematics, Loops

1. Gauge Theories

- ▷ Internal symmetry and the gauge principle
- ▷ Quantization of gauge theories
- ▷ Spontaneous Symmetry Breaking

2. The Standard Model

- ▷ Gauge group and particle representations
- ▷ The SM with one family: electroweak interactions
- ▷ Electroweak SSB: Higgs sector, gauge boson and fermion masses
- ▷ Additional generations: fermion mixings

3. Phenomenology of the Electroweak Standard Model

- ▷ Complete Lagrangian and Feynman rules
- ▷ Input parameters, experiments, observables, precise predictions
- ▷ Global fits

1. Gauge Theories

Internal symmetry

free Lagrangian

- Lagrangian of a free fermion field $\psi(x)$:

$$\text{(Dirac)} \quad \boxed{\mathcal{L}_0 = \bar{\psi}(\mathrm{i}\not{d} - m)\psi} \quad \not{d} \equiv \gamma^\mu \partial_\mu, \quad \bar{\psi} = \psi^\dagger \gamma^0$$

\Rightarrow Invariant under (continuous) global U(1) phase transformations:

$$\psi(x) \mapsto \psi'(x) = e^{-iq\theta} \psi(x), \quad q, \theta \text{ (constants)} \in \mathbb{R}$$

\Rightarrow By Noether's theorem there is a conserved current:

$$j^\mu = q \bar{\psi} \gamma^\mu \psi, \quad \partial_\mu j^\mu = 0$$

and a Noether charge:

$$Q = \int d^3x j^0, \quad \partial_t Q = 0$$

Internal symmetry

free Lagrangian

- A free fermion field:

$$\psi(x) = \int \frac{d^3 p}{(2\pi)^3 \sqrt{2E_p}} \sum_{s=1,2} \left(a_{\mathbf{p},s} u^{(s)}(\mathbf{p}) e^{-ipx} + b_{\mathbf{p},s}^\dagger v^{(s)}(\mathbf{p}) e^{ipx} \right)$$

- is a **solution** of the **Dirac equation** (Euler-Lagrange):

$$(i\cancel{D} - m)\psi(x) = 0 , \quad (\not{p} - m)u(\mathbf{p}) = 0 , \quad (\not{p} + m)v(\mathbf{p}) = 0 ,$$

and after **quantization**

- is an **operator** from the **canonical quantization rules** (anticommutation):

$$\{a_{\mathbf{p},r}, a_{\mathbf{k},s}^\dagger\} = \{b_{\mathbf{p},r}, b_{\mathbf{k},s}^\dagger\} = (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{k}) \delta_{rs} , \quad \{a_{\mathbf{p},r}, a_{\mathbf{k},s}\} = \dots = 0 ,$$

- that annihilates/creates **particles/antiparticles** on the **Fock space** of fermions

Internal symmetry

free Lagrangian

- For a quantized free fermion field:

⇒ The Noether charge is an operator:^{*}

$$:Q: = q \int d^3x : \bar{\psi} \gamma^0 \psi : = q \int \frac{d^3p}{(2\pi)^3} \sum_{s=1,2} \left(a_{\mathbf{p},s}^\dagger a_{\mathbf{p},s} - b_{\mathbf{p},s}^\dagger b_{\mathbf{p},s} \right)$$

$$Q a_{\mathbf{k},s}^\dagger |0\rangle = +q a_{\mathbf{k},s}^\dagger |0\rangle \text{ (particle)} , \quad Q b_{\mathbf{k},s}^\dagger |0\rangle = -q b_{\mathbf{k},s}^\dagger |0\rangle \text{ (antiparticle)}$$

^{*} normal ordering prescription for fermionic operators has been introduced (subtract infinite zero-point energy):

$$: a_{\mathbf{p},r} a_{\mathbf{q},s}^\dagger : \equiv -a_{\mathbf{q},s}^\dagger a_{\mathbf{p},r} , \quad : b_{\mathbf{p},r} b_{\mathbf{q},s}^\dagger : \equiv -b_{\mathbf{q},s}^\dagger b_{\mathbf{p},r}$$

The gauge principle

gauge symmetry dictates interactions

- To make \mathcal{L}_0 invariant under local \equiv gauge transformations of U(1):

$$\psi(x) \mapsto \psi'(x) = e^{-iq\theta(x)} \psi(x), \quad \theta = \theta(x) \in \mathbb{R}$$

perform the minimal substitution:

$$\partial_\mu \rightarrow D_\mu = \partial_\mu + ie q A_\mu \quad (\text{covariant derivative})$$

where a gauge field $A_\mu(x)$ is introduced transforming as:

$$A_\mu(x) \mapsto A'_\mu(x) = A_\mu(x) + \frac{1}{e} \partial_\mu \theta(x) \quad \Leftarrow \quad D_\mu \psi \mapsto e^{-iq\theta(x)} D_\mu \psi \quad \bar{\psi} D \psi \text{ inv.}$$

\Rightarrow The new Lagrangian contains interactions between ψ and A_μ :

$$\boxed{\mathcal{L}_{\text{int}} = -eq \bar{\psi} \gamma^\mu \psi A_\mu} \quad \propto \begin{cases} \text{coupling} & e \\ \text{charge} & q \end{cases}$$

$$(= -e j^\mu A_\mu)$$

The gauge principle

gauge invariance dictates interactions

- Dynamics for the gauge field \Rightarrow add gauge invariant kinetic term:

(Maxwell)

$$\mathcal{L}_1 = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$$

$$\Leftarrow F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \mapsto F_{\mu\nu}$$

- The full U(1) gauge invariant Lagrangian for a fermion field $\psi(x)$ reads:

$$\mathcal{L}_{\text{sym}} = \bar{\psi}(iD - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (= \mathcal{L}_0 + \mathcal{L}_{\text{int}} + \mathcal{L}_1) \quad (\text{QED})$$

- The same applies to a complex scalar field $\phi(x)$:

$$\mathcal{L}_{\text{sym}} = (D_\mu\phi)^\dagger D^\mu\phi - m^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2 - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (\text{sQED})$$

The gauge principle

non-Abelian gauge theories

- A general gauge symmetry group G is an *compact N -dimensional Lie group*

$$g \in G , \quad g(\theta) = e^{-iT_a\theta^a} , \quad a = 1, \dots, N$$

$$\theta^a = \theta^a(x) \in \mathbb{R} , \quad T_a = \text{Hermitian generators} , \quad [T_a, T_b] = i f_{abc} T_c \quad (\text{Lie algebra})$$

$$\text{Tr}\{T_a T_b\} \equiv \frac{1}{2} \delta_{ab} \quad \begin{array}{l} \text{structure constants: } f_{abc} = 0 \text{ Abelian} \\ f_{abc} \neq 0 \text{ non-Abelian} \end{array}$$

\Rightarrow *Unitary finite-dimensional irreducible representations:*

$g(\theta)$ represented by $U(\theta)$

$d \times d$ matrices : $U(\theta)$ [given by $\{T_a\}$ algebra representation]

$$d\text{-multiplet} : \Psi(x) \mapsto \Psi'(x) = U(\theta)\Psi(x) , \quad \Psi = \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_d \end{pmatrix}$$

The gauge principle

non-Abelian gauge theories

- Examples:

| G | N | Abelian |
|-----------|-----------|---|
| U(1) | 1 | Yes |
| SU(n) | $n^2 - 1$ | No ($n \times n$ unitary matrices with $\det = 1$) |

- U(1): 1 generator (q), one-dimensional irreps only
- SU(2): 3 generators

$$f_{abc} = \epsilon_{abc} \text{ (Levi-Civita symbol)}$$

- * Fundamental irrep ($d = 2$): $T_a = \frac{1}{2}\sigma_a$ (3 Pauli matrices)
- * Adjoint irrep ($d = N = 3$): $(T_a^{\text{adj}})_{bc} = -i f_{abc}$

- SU(3): 8 generators

$$f^{123} = 1, f^{458} = f^{678} = \frac{\sqrt{3}}{2}, f^{147} = f^{156} = f^{246} = f^{247} = f^{345} = -f^{367} = \frac{1}{2}$$

- * Fundamental irrep ($d = 3$): $T_a = \frac{1}{2}\lambda_a$ (8 Gell-Mann matrices)
- * Adjoint irrep ($d = N = 8$): $(T_a^{\text{adj}})_{bc} = -i f_{abc}$

(for SU(n): f_{abc} totally antisymmetric)

The gauge principle

non-Abelian gauge theories

- To make \mathcal{L}_0 invariant under local \equiv gauge transformations of G :

$$\mathcal{L}_0 = \bar{\Psi}(\mathrm{i}\partial - m)\Psi , \quad \Psi(x) \mapsto \Psi'(x) = U(\theta)\Psi(x) , \quad \theta = \theta(x) \in \mathbb{R}$$

substitute the covariant derivative:

$$\partial_\mu \rightarrow D_\mu = \partial_\mu - \mathrm{i}g\tilde{W}_\mu , \quad \tilde{W}_\mu \equiv T_a W_\mu^a$$

where a gauge field $W_\mu^a(x)$ per generator is introduced (adjoint irrep), transforming as:

$$\tilde{W}_\mu(x) \mapsto \tilde{W}'_\mu(x) = U\tilde{W}_\mu(x)U^\dagger - \frac{\mathrm{i}}{g}(\partial_\mu U)U^\dagger \quad \Leftarrow \quad [D_\mu\Psi \mapsto UD_\mu\Psi] \quad \bar{\Psi}D\Psi \text{ inv. } \boxed{1}$$

\Rightarrow The new Lagrangian contains interactions between Ψ and W_μ^a :

$$\boxed{\mathcal{L}_{\text{int}} = g \bar{\Psi} \gamma^\mu T_a \Psi W_\mu^a} \quad \propto \begin{cases} \text{coupling} & g \\ \text{charge} & T_a \end{cases}$$

$$(= g j_a^\mu W_\mu^a)$$

The gauge principle

non-Abelian gauge theories

- Dynamics for the gauge fields \Rightarrow add gauge invariant kinetic terms:

(Yang-Mills)

$$\mathcal{L}_{\text{YM}} = -\frac{1}{2} \text{Tr} \left\{ \tilde{W}_{\mu\nu} \tilde{W}^{\mu\nu} \right\} = -\frac{1}{4} W_{\mu\nu}^a W^{a,\mu\nu}$$

$$\tilde{W}_{\mu\nu} \equiv D_\mu \tilde{W}_\nu - D_\nu \tilde{W}_\mu = \partial_\mu \tilde{W}_\nu - \partial_\nu \tilde{W}_\mu - ig [\tilde{W}_\mu, \tilde{W}_\nu] \quad \Leftrightarrow \quad \tilde{W}_{\mu\nu} \mapsto U \tilde{W}_{\mu\nu} U^\dagger$$

$$\Rightarrow W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g f_{abc} W_\mu^b W_\nu^c$$

$\Rightarrow \mathcal{L}_{\text{YM}}$ contains cubic and quartic self-interactions of the gauge fields W_μ^a :

$$\mathcal{L}_{\text{kin}} = -\frac{1}{4} (\partial_\mu W_\nu^a - \partial_\nu W_\mu^a) (\partial^\mu W^{a,\nu} - \partial^\nu W^{a,\mu})$$

$$\mathcal{L}_{\text{cubic}} = -\frac{1}{2} g f_{abc} (\partial_\mu W_\nu^a - \partial_\nu W_\mu^a) W^{b,\mu} W^{c,\nu}$$

$$\mathcal{L}_{\text{quartic}} = -\frac{1}{4} g^2 f_{abef} f_{cde} W_\mu^a W_\nu^b W^{c,\mu} W^{d,\nu}$$

Quantization of gauge theories

propagators

- The (Feynman) propagator of a scalar field:

$$D(x - y) = \langle 0 | T\{\phi(x)\phi^\dagger(y)\} | 0 \rangle = \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p^2 - m^2 + i0} e^{-ip \cdot (x-y)}$$

is a Green's function of the Klein-Gordon operator:

$$(\square_x + m^2)D(x - y) = -i\delta^4(x - y) \quad \Leftrightarrow \quad \tilde{D}(p) = \frac{i}{p^2 - m^2 + i0}$$

- The propagator of a fermion field:

$$S(x - y) = \langle 0 | T\{\psi(x)\bar{\psi}(y)\} | 0 \rangle = (i\cancel{\partial}_x + m) \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p^2 - m^2 + i0} e^{-ip \cdot (x-y)}$$

is a Green's function of the Dirac operator:

$$(i\cancel{\partial}_x - m)S(x - y) = i\delta^4(x - y) \quad \Leftrightarrow \quad \tilde{S}(p) = \frac{i}{\cancel{p} - m + i0}$$

Quantization of gauge theories

propagators

- BUT the propagator of a gauge field cannot be defined unless \mathcal{L} is modified:

(e.g. modified Maxwell)
$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\xi}(\partial^\mu A_\mu)^2$$

Euler-Lagrange:
$$\frac{\partial \mathcal{L}}{\partial A_\nu} - \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu A_\nu)} = 0 \quad \Rightarrow \quad \left[g^{\mu\nu} \square - \left(1 - \frac{1}{\xi}\right) \partial^\mu \partial^\nu \right] A_\mu = 0$$

- In momentum space the propagator is the inverse of:

$$-k^2 g^{\mu\nu} + \left(1 - \frac{1}{\xi}\right) k^\mu k^\nu \quad \Rightarrow \quad \tilde{D}_{\mu\nu}(k) = \frac{i}{k^2 + i0} \left[-g_{\mu\nu} + (1 - \xi) \frac{k_\mu k_\nu}{k^2} \right]$$

⇒ Note that $(-k^2 g^{\mu\nu} + k^\mu k^\nu)$ is singular!

⇒ One may argue that \mathcal{L} above will not lead to Maxwell equations ...
unless we fix a (Lorenz) gauge where:

$$\partial^\mu A_\mu = 0 \quad \Leftarrow \quad A_\mu \mapsto A'_\mu = A_\mu + \partial_\mu \Lambda \quad \text{with} \quad \partial^\mu \partial_\mu \Lambda \equiv -\partial^\mu A_\mu$$

- The extra term is called **Gauge Fixing**:

$$\mathcal{L}_{\text{GF}} = -\frac{1}{2\xi}(\partial^\mu A_\mu)^2$$

\Rightarrow modified \mathcal{L} equivalent to Maxwell Lagrangian just in the gauge $\partial^\mu A_\mu = 0$

\Rightarrow the ξ -dependence always cancels out in physical amplitudes

- Several choices for the gauge fixing term (simplify calculations): R_ξ gauges

('t Hooft-Feynman gauge) $\xi = 1$: $\tilde{D}_{\mu\nu}(k) = -\frac{\mathrm{i}g_{\mu\nu}}{k^2 + \mathrm{i}0}$

(Landau gauge) $\xi = 0$: $\tilde{D}_{\mu\nu}(k) = \frac{\mathrm{i}}{k^2 + \mathrm{i}0} \left[-g_{\mu\nu} + \frac{k_\mu k_\nu}{k^2} \right]$

Quantization of gauge theories

gauge fixing

(non-Abelian case)

- For a non-Abelian gauge theory, the gauge fixing terms:

$$\mathcal{L}_{\text{GF}} = - \sum_a \frac{1}{2\xi_a} (\partial^\mu W_\mu^a)^2$$

allow to define the propagators:

$$\tilde{D}_{\mu\nu}^{ab}(k) = \frac{i\delta_{ab}}{k^2 + i0} \left[-g_{\mu\nu} + (1 - \xi_a) \frac{k_\mu k_\nu}{k^2} \right]$$

BUT, unlike the Abelian case, this is not the end of the story ...

Quantization of gauge theories

Faddeev-Popov ghosts

- Add Faddeev-Popov ghost fields $c_a(x)$, $a = 1, \dots, N$:


$$\mathcal{L}_{\text{FP}} = (\partial^\mu \bar{c}_a) (D_\mu^{\text{adj}})_{ab} c_b = (\partial^\mu \bar{c}_a) (\partial_\mu c_a - g f_{abc} c_b W_\mu^c)$$

$$\Leftarrow D_\mu^{\text{adj}} = \partial_\mu - i g T_c^{\text{adj}} W_\mu^c$$

Computational trick: *anticommuting* scalar fields, just in loops as virtual particles

⇒ Faddeev-Popov ghosts needed to preserve gauge symmetry:

loops 2 3



Self Energy

$$= \Pi_{\mu\nu} = i(g_{\mu\nu}k^2 - k_\mu k_\nu)\Pi(k^2)$$

Ward identity: $k^\mu \Pi_{\mu\nu} = 0$

with

$$\tilde{D}_{ab}(k) = \frac{i\delta_{ab}}{k^2 + i0}$$

[(-1) sign for closed loops! (like fermions)]

- Then the complete **quantum** Lagrangian is

$$\mathcal{L}_{\text{sym}} + \mathcal{L}_{\text{GF}} + \mathcal{L}_{\text{FP}}$$

⇒ Note that in the case of a **massive** vector field

$$(\text{Proca}) \quad \mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}M^2A_\mu A^\mu$$

it is **not gauge invariant**

- The propagator is:

4

$$\tilde{D}_{\mu\nu}(k) = \frac{i}{k^2 - M^2 + i0} \left(-g_{\mu\nu} + \frac{k^\mu k^\nu}{M^2} \right)$$

Spontaneous Symmetry Breaking

discrete symmetry

- Consider a real scalar field $\phi(x)$ with Lagrangian:

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)(\partial^\mu \phi) - \frac{1}{2}\mu^2\phi^2 - \frac{\lambda}{4}\phi^4 \quad \text{invariant under } \phi \mapsto -\phi$$

$$\Rightarrow \mathcal{H} = \frac{1}{2}(\dot{\phi}^2 + (\nabla\phi)^2) + V(\phi)$$

$$V = \frac{1}{2}\mu^2\phi^2 + \frac{1}{4}\lambda\phi^4$$



$\mu^2, \lambda \in \mathbb{R}$ (Real/Hermitian Hamiltonian) and $\lambda > 0$ (existence of a ground state)

(a) $\mu^2 > 0$: min of $V(\phi)$ at $\phi = 0$

(b) $\mu^2 < 0$: min of $V(\phi)$ at $\phi = v \equiv \pm \sqrt{\frac{-\mu^2}{\lambda}}$, in QFT $\langle 0 | \phi | 0 \rangle = v \neq 0$ (VEV)

- A **quantum** field **must** have $v = 0$

$$a |0\rangle = 0$$

$$\Rightarrow \phi(x) \equiv v + \eta(x), \quad \langle 0 | \eta | 0 \rangle = 0$$

Spontaneous Symmetry Breaking

discrete symmetry

- At the quantum level, the **same** system is described by $\eta(x)$ with Lagrangian:

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \eta)(\partial^\mu \eta) - \lambda v^2 \eta^2 - \lambda v \eta^3 - \frac{\lambda}{4} \eta^4 \quad \text{not invariant under } \eta \mapsto -\eta$$
$$(m_\eta = \sqrt{2\lambda} v)$$

⇒ Lesson:

$\mathcal{L}(\phi)$ had the symmetry but the parameters can be such that the ground state of the Hamiltonian is not symmetric (Spontaneous Symmetry Breaking)

⇒ Note:

One may argue that $\mathcal{L}(\eta)$ exhibits an explicit breaking of the symmetry. However this is not the case since the coefficients of terms η^2 , η^3 and η^4 are determined by just two parameters, λ and v (remnant of the original symmetry)

Spontaneous Symmetry Breaking

continuous symmetry

- Consider a complex scalar field $\phi(x)$ with Lagrangian:

$$\mathcal{L} = (\partial_\mu \phi^\dagger)(\partial^\mu \phi) - \mu^2 \phi^\dagger \phi - \lambda(\phi^\dagger \phi)^2 \quad \text{invariant under U(1): } \phi \mapsto e^{-iq\theta}\phi$$


$$\lambda > 0, \mu^2 < 0 : \quad \langle 0 | \phi | 0 \rangle \equiv \frac{v}{\sqrt{2}}, \quad |v| = \sqrt{\frac{-\mu^2}{\lambda}}$$

Take $v \in \mathbb{R}^+$. In terms of quantum fields:

$$\phi(x) \equiv \frac{1}{\sqrt{2}}[v + \eta(x) + i\chi(x)], \quad \langle 0 | \eta | 0 \rangle = \langle 0 | \chi | 0 \rangle = 0$$

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \eta)(\partial^\mu \eta) + \frac{1}{2}(\partial_\mu \chi)(\partial^\mu \chi) - \lambda v^2 \eta^2 - \lambda v \eta (\eta^2 + \chi^2) - \frac{\lambda}{4}(\eta^2 + \chi^2)^2 + \frac{1}{4}\lambda v^4$$

Note: if $v e^{i\alpha}$ (complex) replace η by $(\eta \cos \alpha - \chi \sin \alpha)$ and χ by $(\eta \sin \alpha + \chi \cos \alpha)$



\Rightarrow The actual quantum Lagrangian $\mathcal{L}(\eta, \chi)$ is not invariant under U(1)

U(1) broken \Rightarrow one scalar field remains massless: $m_\eta = \sqrt{2\lambda}v, m_\chi = 0$

Spontaneous Symmetry Breaking

continuous symmetry

- Another example: consider a real scalar SU(2) triplet $\Phi(x)$

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \Phi^\top)(\partial^\mu \Phi) - \frac{1}{2}\mu^2 \Phi^\top \Phi - \frac{\lambda}{4}(\Phi^\top \Phi)^2 \quad \text{inv. under SU(2): } \Phi \mapsto e^{-iT_a \theta^a} \Phi$$

that for $\lambda > 0$, $\mu^2 < 0$ acquires a VEV $\langle 0 | \Phi^\top \Phi | 0 \rangle = v^2$ ($\mu^2 = -\lambda v^2$)

Assume $\Phi(x) = \begin{pmatrix} \varphi_1(x) \\ \varphi_2(x) \\ v + \varphi_3(x) \end{pmatrix}$ and define $\varphi \equiv \frac{1}{\sqrt{2}}(\varphi_1 + i\varphi_2)$

$$\mathcal{L} = (\partial_\mu \varphi^\dagger)(\partial^\mu \varphi) + \frac{1}{2}(\partial_\mu \varphi_3)(\partial^\mu \varphi_3) - \lambda v^2 \varphi_3^2 - \lambda v(2\varphi^\dagger \varphi + \varphi_3^2)\varphi_3 - \frac{\lambda}{4}(2\varphi^\dagger \varphi + \varphi_3^2)^2 + \frac{1}{4}\lambda v^4$$

\Rightarrow Not symmetric under SU(2) but invariant under U(1):

$$\varphi \mapsto e^{-iq\theta} \varphi \quad (q = \text{arbitrary}) \qquad \qquad \varphi_3 \mapsto \varphi_3 \quad (q = 0)$$

SU(2) broken to U(1) $\Rightarrow 3 - 1 = 2$ broken generators

\Rightarrow 2 (real) scalar fields (= 1 complex) remain massless: $m_\varphi = 0$, $m_{\varphi_3} = \sqrt{2\lambda}v$

Spontaneous Symmetry Breaking

continuous symmetry

⇒ Goldstone's theorem:

[Nambu '60; Goldstone '61]

The number of massless particles (Nambu-Goldstone bosons) is equal to the number of spontaneously broken generators of the symmetry

Hamiltonian symmetric under group $G \Rightarrow [T_a, H] = 0, a = 1, \dots, N$

By definition: $H |0\rangle = 0 \Rightarrow H(T_a |0\rangle) = T_a H |0\rangle = 0$

- If $|0\rangle$ is such that $T_a |0\rangle = 0$ for all generators
⇒ non-degenerate minimum: *the* vacuum
- If $|0\rangle$ is such that $T_{a'} |0\rangle \neq 0$ for some (broken) generators a'
⇒ degenerate minimum: chose one (*true* vacuum) and $e^{-iT_{a'}\theta^{a'}} |0\rangle \neq |0\rangle$
⇒ excitations (particles) from $|0\rangle$ to $e^{-iT_{a'}\theta^{a'}} |0\rangle$ cost no energy: massless!

Spontaneous Symmetry Breaking

gauge symmetry

- Consider a U(1) gauge invariant Lagrangian for a complex scalar field $\phi(x)$:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + (D_\mu\phi)^\dagger(D^\mu\phi) - \mu^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2, \quad D_\mu = \partial_\mu + ieqA_\mu$$

inv. under $\phi(x) \mapsto \phi'(x) = e^{-iq\theta(x)}\phi(x), \quad A_\mu(x) \mapsto A'_\mu(x) = A_\mu(x) + \frac{1}{e}\partial_\mu\theta(x)$

If $\lambda > 0, \mu^2 < 0$, the \mathcal{L} in terms of quantum fields η and χ with null VEVs:

$$\phi(x) \equiv \frac{1}{\sqrt{2}}[v + \eta(x) + i\chi(x)], \quad \mu^2 = -\lambda v^2$$

Comments:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}(\partial_\mu\eta)(\partial^\mu\eta) + \frac{1}{2}(\partial_\mu\chi)(\partial^\mu\chi) \\ & - \lambda v^2\eta^2 - \lambda v\eta(\eta^2 + \chi^2) - \frac{\lambda}{4}(\eta^2 + \chi^2)^2 + \frac{1}{4}\lambda v^4 \\ & + eqvA_\mu\partial^\mu\chi + eqA_\mu(\eta\partial^\mu\chi - \chi\partial^\mu\eta) \\ & + \frac{1}{2}(eqv)^2A_\mu A^\mu + \frac{1}{2}(eq)^2A_\mu A^\mu(\eta^2 + 2v\eta + \chi^2) \end{aligned}$$

(i) $m_\eta = \sqrt{2\lambda}v$
 $m_\chi = 0$

(ii) $M_A = |eqv|$ (!)

(iii) Term $A_\mu\partial^\mu\chi$ (?)

(iv) Add \mathcal{L}_{GF}

Spontaneous Symmetry Breaking

gauge symmetry

- Removing the cross term and the (new) gauge fixing Lagrangian:

$$\begin{aligned} \mathcal{L}_{\text{GF}} &= -\frac{1}{2\xi}(\partial_\mu A^\mu - \xi M_A \chi)^2 \\ \Rightarrow \quad \mathcal{L} + \mathcal{L}_{\text{GF}} &= -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}M_A^2 A_\mu A^\mu - \frac{1}{2\xi}(\partial_\mu A^\mu)^2 + \overbrace{M_A[A_\mu \partial^\mu \chi + \partial_\mu A^\mu \chi]}^{\text{total deriv.}} \\ &\quad + \frac{1}{2}(\partial_\mu \chi)(\partial^\mu \chi) - \frac{1}{2}\xi M_A^2 \chi^2 + \dots \end{aligned}$$

and the propagators of A_μ and χ are:

(5)

$$\begin{aligned} \tilde{D}_{\mu\nu}(k) &= \frac{i}{k^2 - M_A^2 + i0} \left[-g_{\mu\nu} + (1 - \xi) \frac{k_\mu k_\nu}{k^2 - \xi M_A^2} \right] \\ \tilde{D}(k) &= \frac{i}{k^2 - \xi M_A^2 + i0} \end{aligned}$$

$\Rightarrow \chi$ has a gauge-dependent mass: actually it is not a physical field!

Spontaneous Symmetry Breaking

gauge symmetry

- A more transparent parameterization of the quantum field ϕ is

$$\phi(x) \equiv e^{iq\zeta(x)/v} \frac{1}{\sqrt{2}} [v + \eta(x)] , \quad \langle 0 | \eta | 0 \rangle = \langle 0 | \zeta | 0 \rangle = 0$$

$$\phi(x) \mapsto e^{-iq\zeta(x)/v} \phi(x) = \frac{1}{\sqrt{2}} [v + \eta(x)] \quad \Rightarrow \quad \zeta \text{ gauged away!}$$

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} (\partial_\mu \eta)(\partial^\mu \eta)$$

$$- \lambda v^2 \eta^2 - \lambda v \eta^3 - \frac{\lambda}{4} \eta^4 + \frac{1}{4} \lambda v^4$$

$$+ \frac{1}{2} (eqv)^2 A_\mu A^\mu + \frac{1}{2} (eq)^2 A_\mu A^\mu (2v\eta + \eta^2)$$

Comments:

(i) $m_\eta = \sqrt{2\lambda} v$

(ii) $M_A = |eqv|$

(iii) No need for \mathcal{L}_{GF}

\Rightarrow This is the unitary gauge ($\xi \rightarrow \infty$): just physical fields

$$\tilde{D}_{\mu\nu}(k) \rightarrow \frac{i}{k^2 - M_A^2 + i0} \left[-g_{\mu\nu} + \frac{k_\mu k_\nu}{M_A^2} \right] \quad \text{and} \quad \tilde{D}(k) \rightarrow 0$$

Spontaneous Symmetry Breaking

gauge symmetry

⇒ Brout-Englert-Higgs mechanism:

[Anderson '62]

[Higgs '64; Englert, Brout '64; Guralnik, Hagen, Kibble '64]

The gauge bosons associated with the spontaneously broken generators become massive, the corresponding would-be Goldstone bosons are unphysical and can be absorbed, the remaining massive scalars (Higgs bosons) are physical (the smoking gun!)

- The would-be Goldstone bosons are ‘eaten up’ by the gauge bosons (‘get fat’) and disappear (gauge away) in the unitary gauge ($\xi \rightarrow \infty$)
⇒ Degrees of freedom are preserved

Before SSB: 2 (massless gauge boson) + 1 (Goldstone boson)

After SSB: 3 (massive gauge boson) + 0 (absorbed would-be Goldstone)

- For loops calculations, ’t Hooft-Feynman gauge ($\xi = 1$) is more convenient:
⇒ Gauge boson propagators are simpler, but
⇒ Goldstone bosons must be included in internal lines

Spontaneous Symmetry Breaking

gauge symmetry

- Comments:

- After SSB the FP ghost fields (unphysical) acquire a gauge-dependent mass, due to interactions with the scalar field(s):

$$\tilde{D}_{ab}(k) = \frac{i\delta_{ab}}{k^2 - \xi_a M_{W^a}^2 + i0}$$

- Gauge theories with SSB are renormalizable

['t Hooft, Veltman '72]

UV divergences appearing at loop level can be removed by renormalization of parameters and fields of the classical Lagrangian \Rightarrow predictive!

2. The Standard Model

Gauge group and particle representations

[Glashow '61; Weinberg '67; Salam '68]
 [D. Gross, F. Wilczek; D. Politzer '73]

- The Standard Model is a gauge theory based on the local symmetry group:

$$\underbrace{\text{SU}(3)_c \otimes \text{SU}(2)_L \otimes \text{U}(1)_Y}_{\text{strong} \quad \text{electroweak}} \rightarrow \text{SU}(3)_c \otimes \underbrace{\text{U}(1)_Q}_{\text{em}}$$

with the electroweak symmetry spontaneously broken to the electromagnetic $\text{U}(1)_Q$ symmetry by the Brout-Englert-Higgs mechanism

- The particle (field) content: (ingredients: 12 *flavors* + 12 gauge bosons + H)

| Fermions | | | I | II | III | Q | Bosons | | |
|--------------------|---------|------|-------------|-------------|-------------|----------------|----------|----------------|--------------------|
| spin $\frac{1}{2}$ | Quarks | f | u u u | c c c | t t t | $\frac{2}{3}$ | spin 1 | 8 gluons | strong interaction |
| | | f' | d d d | s s s | b b b | $-\frac{1}{3}$ | | W^\pm, Z | weak interaction |
| | Leptons | f | ν_e | ν_μ | ν_τ | 0 | γ | em interaction | |
| | | f' | e | μ | τ | -1 | | spin 0 | Higgs |

$$Q_f = Q_{f'} + 1$$

Gauge group and particle representations

- The fields lay in the following representations (color, weak isospin, hypercharge):

| Multiplets | $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ | I | II | III | $Q = T_3 + Y$ |
|------------|--|--|--|--|--|
| Quarks | $(3, 2, \frac{1}{6})$ | $\begin{pmatrix} u_L \\ d_L \end{pmatrix}$ | $\begin{pmatrix} c_L \\ s_L \end{pmatrix}$ | $\begin{pmatrix} t_L \\ b_L \end{pmatrix}$ | $\frac{2}{3} = \frac{1}{2} + \frac{1}{6}$ $-\frac{1}{3} = -\frac{1}{2} + \frac{1}{6}$ |
| | $(3, 1, \frac{2}{3})$ | u_R | c_R | t_R | $\frac{2}{3} = 0 + \frac{2}{3}$ |
| | $(3, 1, -\frac{1}{3})$ | d_R | s_R | b_R | $-\frac{1}{3} = 0 - \frac{1}{3}$ |
| Leptons | $(1, 2, -\frac{1}{2})$ | $\begin{pmatrix} \nu_{e_L} \\ e_L \end{pmatrix}$ | $\begin{pmatrix} \nu_{\mu_L} \\ \mu_L \end{pmatrix}$ | $\begin{pmatrix} \nu_{\tau_L} \\ \tau_L \end{pmatrix}$ | $0 = \frac{1}{2} - \frac{1}{2}$ $-1 = -\frac{1}{2} - \frac{1}{2}$ |
| | $(1, 1, -1)$ | e_R | μ_R | τ_R | $-1 = 0 - 1$ |
| | $(1, 1, 0)$ | ν_{e_R} | ν_{μ_R} | ν_{τ_R} | $0 = 0 + 0$ |
| Higgs | $(1, 2, \frac{1}{2})$ | (3 families of quarks & leptons) | | | |

\Rightarrow Electroweak (QFD): $SU(2)_L \otimes U(1)_Y$ Strong (QCD) $SU(3)_c$

The EWSM with one family (of quarks or leptons)

- Consider two massless fermion fields $f(x)$ and $f'(x)$ with electric charges $Q_f = Q_{f'} + 1$ in three irreps of $SU(2)_L \otimes U(1)_Y$:

$$\begin{aligned} \mathcal{L}_F^0 &= i\bar{f}\not{\partial}f + i\bar{f}'\not{\partial}f' & f_{R,L} &= \frac{1}{2}(1 \pm \gamma_5)f, \quad f'_{R,L} = \frac{1}{2}(1 \pm \gamma_5)f' \\ &= i\bar{\Psi}_1\not{\partial}\Psi_1 + i\bar{\psi}_2\not{\partial}\psi_2 + i\bar{\psi}_3\not{\partial}\psi_3 \quad ; \quad \Psi_1 = \underbrace{\begin{pmatrix} f_L \\ f'_L \end{pmatrix}}_{(\mathbf{2}, y_1)}, \quad \psi_2 = \underbrace{\psi_R}_{(\mathbf{1}, y_2)}, \quad \psi_3 = \underbrace{\psi'_R}_{(\mathbf{1}, y_3)} \end{aligned}$$

- To get a Langrangian invariant under gauge transformations:

$$\Psi_1(x) \mapsto U_L(x)e^{-iy_1\beta(x)}\Psi_1(x), \quad U_L(x) = e^{-iT_i\alpha^i(x)}, \quad T_i = \frac{\sigma_i}{2} \quad (\text{weak isospin gen.})$$

$$\psi_2(x) \mapsto e^{-iy_2\beta(x)}\psi_2(x)$$

$$\psi_3(x) \mapsto e^{-iy_3\beta(x)}\psi_3(x)$$

The EWSM with one family

gauge invariance

⇒ Introduce gauge fields $W_\mu^i(x)$ ($i = 1, 2, 3$) and $B_\mu(x)$ through **covariant derivatives**:

$$\left. \begin{aligned} D_\mu \Psi_1 &= (\partial_\mu - ig\tilde{W}_\mu + ig'y_1 B_\mu) \Psi_1 , & \tilde{W}_\mu &\equiv \frac{\sigma_i}{2} W_\mu^i \\ D_\mu \psi_2 &= (\partial_\mu + ig'y_2 B_\mu) \psi_2 \\ D_\mu \psi_3 &= (\partial_\mu + ig'y_3 B_\mu) \psi_3 \end{aligned} \right\} \Rightarrow \boxed{\mathcal{L}_F}$$

where two couplings g and g' have been introduced and

$$\begin{aligned} \tilde{W}_\mu(x) &\mapsto U_L(x)\tilde{W}_\mu(x)U_L^\dagger(x) - \frac{i}{g}(\partial_\mu U_L(x))U_L^\dagger(x) \\ B_\mu(x) &\mapsto B_\mu(x) + \frac{1}{g'}\partial_\mu\beta(x) \end{aligned}$$

⇒ Add **Yang-Mills**: gauge invariant kinetic terms for the gauge fields

$$\boxed{\mathcal{L}_{YM}} = -\frac{1}{4}W_{\mu\nu}^i W^{i,\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} , \quad W_{\mu\nu}^i = \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + g\epsilon_{ijk}W_\mu^j W_\nu^k$$

(include self-interactions of the SU(2) gauge fields) and $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$

The EWSM with one family

mass terms forbidden

⇒ Note that mass terms are not invariant under $SU(2)_L \otimes U(1)_Y$, since LH and RH components do not transform the same:

$$m\bar{f}f = m(\overline{f_L}f_R + \overline{f_R}f_L)$$

⇒ Mass terms for the gauge bosons are not allowed either

⇒ Next the different types of interactions are analyzed

The EWSM with one family


charged current interactions

-

$$\mathcal{L}_F \supset g \bar{\Psi}_1 \gamma^\mu \tilde{W}_\mu \Psi_1 , \quad \tilde{W}_\mu = \frac{1}{2} \begin{pmatrix} W_\mu^3 & \sqrt{2} W_\mu^+ \\ \sqrt{2} W_\mu^- & -W_\mu^3 \end{pmatrix}$$

\Rightarrow charged current interactions of LH fermions with complex vector boson field W_μ :

$$\mathcal{L}_{CC} = \frac{g}{2\sqrt{2}} \bar{f} \gamma^\mu (1 - \gamma_5) f' W_\mu^\dagger + \text{h.c.} , \quad W_\mu \equiv \frac{1}{\sqrt{2}} (W_\mu^1 + i W_\mu^2)$$



The EWSM with one family

neutral current interactions

- The diagonal part of

$$\mathcal{L}_F \supset g \bar{\Psi}_1 \gamma^\mu \tilde{W}_\mu \Psi_1 - g' B_\mu (y_1 \bar{\Psi}_1 \gamma^\mu \Psi_1 + y_2 \bar{\psi}_2 \gamma^\mu \psi_2 + y_3 \bar{\psi}_3 \gamma^\mu \psi_3)$$

\Rightarrow neutral current interactions with neutral vector boson fields W_μ^3 and B_μ

We would like to identify B_μ with the photon field A_μ but that requires:

$$y_1 = y_2 = y_3 \quad \text{and} \quad g' y_j = e Q_j \quad \Rightarrow \quad \text{impossible!}$$

\Rightarrow Since they are both neutral, try a combination:

$$\begin{pmatrix} W_\mu^3 \\ B_\mu \end{pmatrix} = \begin{pmatrix} c_W & -s_W \\ s_W & c_W \end{pmatrix} \begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix} \quad \begin{aligned} s_W &\equiv \sin \theta_W, & c_W &\equiv \cos \theta_W \\ \theta_W &= \text{weak mixing angle} \end{aligned}$$

$$\mathcal{L}_{\text{NC}} = \sum_{j=1}^3 \bar{\psi}_j \gamma^\mu \left\{ -[g T_3 s_W + g' y_j c_W] A_\mu + [g T_3 c_W - g' y_j s_W] Z_\mu \right\} \psi_j$$

with $T_3 = \frac{\sigma_3}{2}$ (0) the third weak isospin component of the doublet (singlet)

The EWSM with one family

neutral current interactions

- To make A_μ the photon field:

$$(1) \boxed{e = gs_W = g'c_W} \quad (2) \boxed{Q = T_3 + Y}$$

where the electric charge operator is: $Q_1 = \begin{pmatrix} Q_f & 0 \\ 0 & Q_{f'} \end{pmatrix}, \quad Q_2 = Q_f, \quad Q_3 = Q_{f'}$

\Rightarrow (1) **Electroweak unification**: g of SU(2) and g' of U(1) related to $e = \frac{gg'}{\sqrt{g^2 + g'^2}}$

\Rightarrow (2) The hypercharges are fixed in terms of electric charges and weak isospin:

$$y_1 = Q_f - \frac{1}{2} = Q_{f'} + \frac{1}{2}, \quad y_2 = Q_f, \quad y_3 = Q_{f'}$$

$$\mathcal{L}_{\text{QED}} = -e Q_f \bar{f} \gamma^\mu f A_\mu + (f \rightarrow f')$$

\Rightarrow RH neutrinos are sterile: $y_2 = Q_f = 0$

The EWSM with one family

neutral current interactions

- The Z_μ is the neutral weak boson field:

$$\mathcal{L}_{\text{NC}}^Z = e \bar{f} \gamma^\mu (v_f - a_f \gamma_5) f Z_\mu + (f \rightarrow f')$$

with

$$v_f = \frac{T_3^{f_L} - 2Q_f s_W^2}{2s_W c_W}, \quad a_f = \frac{T_3^{f_L}}{2s_W c_W}$$

- The complete neutral current Lagrangian reads:

$$\mathcal{L}_{\text{NC}} = \mathcal{L}_{\text{QED}} + \mathcal{L}_{\text{NC}}^Z$$



The EWSM with one family


gauge boson self-interactions

- Cubic:

$$\begin{aligned}\mathcal{L}_{\text{YM}} \supset \mathcal{L}_3 = & -\frac{iec_W}{s_W} \left\{ W^{\mu\nu} W_\mu^\dagger Z_\nu - W_{\mu\nu}^\dagger W^\mu Z^\nu - W_\mu^\dagger W_\nu Z^{\mu\nu} \right\} \\ & + ie \left\{ W^{\mu\nu} W_\mu^\dagger A_\nu - W_{\mu\nu}^\dagger W^\mu A^\nu - W_\mu^\dagger W_\nu F^{\mu\nu} \right\}\end{aligned}$$

with

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu \quad W_{\mu\nu} = \partial_\mu W_\nu - \partial_\nu W_\mu$$




The EWSM with one family

gauge boson self-interactions

- Quartic:

$$\begin{aligned}\mathcal{L}_{\text{YM}} \supset \mathcal{L}_4 = & -\frac{e^2}{2s_W^2} \left\{ (W_\mu^\dagger W^\mu)^2 - W_\mu^\dagger W^{\mu\dagger} W_\nu W^\nu \right\} \\ & - \frac{e^2 c_W^2}{s_W^2} \left\{ W_\mu^\dagger W^\mu Z_\nu Z^\nu - W_\mu^\dagger Z^\mu W_\nu Z^\nu \right\} \\ & + \frac{e^2 c_W}{s_W} \left\{ 2W_\mu^\dagger W^\mu Z_\nu A^\nu - W_\mu^\dagger Z^\mu W_\nu A^\nu - W_\mu^\dagger A^\mu W_\nu Z^\nu \right\} \\ & - e^2 \left\{ W_\mu^\dagger W^\mu A_\nu A^\nu - W_\mu^\dagger A^\mu W_\nu A^\nu \right\}\end{aligned}$$



Note: even number of W and no vertex with just γ or Z

Electroweak symmetry breaking

setup

- Out of the 4 gauge bosons of $SU(2)_L \otimes U(1)_Y$ with generators T_1, T_2, T_3, Y we need all to be broken except the combination $Q = T_3 + Y$ so that A_μ remains massless and the other three gauge bosons get massive after SSB
 \Rightarrow Introduce a complex $SU(2)$ Higgs doublet

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad \langle 0 | \Phi | 0 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}$$

with gauge invariant Lagrangian ($\mu^2 = -\lambda v^2$):

$$[\mathcal{L}_\Phi] = (D_\mu \Phi)^\dagger D^\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2, \quad D_\mu \Phi = (\partial_\mu - i g \tilde{W}_\mu + i g' y_\Phi B_\mu) \Phi$$

$$\text{take } y_\Phi = \frac{1}{2} \quad \Rightarrow \quad (T_3 + Y) |0\rangle = Q \begin{pmatrix} 0 \\ v \end{pmatrix} = 0$$

$$\{T_1, T_2, T_3 - Y\} |0\rangle \neq 0$$

Electroweak symmetry breaking

gauge boson masses

- Quantum fields in the unitary gauge:

$$\Phi(x) \equiv \exp \left\{ i \frac{\sigma_i}{2v} \theta^i(x) \right\} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix}$$

$$\Phi(x) \mapsto \exp \left\{ -i \frac{\sigma_i}{2v} \theta^i(x) \right\} \Phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H(x) \end{pmatrix} \Rightarrow \begin{array}{l} \text{1 physical Higgs field} \\ H(x) \\ \text{3 would-be Goldstones} \\ \theta^i(x) \text{ gauged away} \end{array}$$

- The 3 dof apparently lost become the longitudinal polarizations of W^\pm and Z that get massive after SSB:


$$\mathcal{L}_\Phi \supset \mathcal{L}_M = \underbrace{\frac{g^2 v^2}{4}}_{M_W^2} W_\mu^\dagger W^\mu + \underbrace{\frac{g^2 v^2}{8c_W^2}}_{\frac{1}{2} M_Z^2} Z_\mu Z^\mu \Rightarrow M_W = M_Z c_W = \frac{1}{2} g v$$

Electroweak symmetry breaking


Higgs sector

⇒ In the unitary gauge (just physical fields): $\mathcal{L}_\Phi = \mathcal{L}_H + \mathcal{L}_M + \mathcal{L}_{HV^2} + \frac{1}{4}\lambda v^4$

$$\mathcal{L}_H = \frac{1}{2}\partial_\mu H \partial^\mu H - \frac{1}{2}M_H^2 H^2 - \frac{M_H^2}{2v}H^3 - \frac{M_H^2}{8v^2}H^4, \quad M_H = \sqrt{-2\mu^2} = \sqrt{2\lambda}v$$



$$\mathcal{L}_M + \mathcal{L}_{HV^2} = M_W^2 W_\mu^\dagger W^\mu \left\{ 1 + \frac{2}{v}H + \frac{H^2}{v^2} \right\} + \frac{1}{2}M_Z^2 Z_\mu Z^\mu \left\{ 1 + \frac{2}{v}H + \frac{H^2}{v^2} \right\}$$



Electroweak symmetry breaking

Higgs sector

- Quantum fields in the R_ξ gauges:

$$\Phi(x) = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \equiv \begin{pmatrix} \phi^+(x) \\ \frac{1}{\sqrt{2}}[v + H(x) + i\chi(x)] \end{pmatrix}, \quad \phi^-(x) = [\phi^+(x)]^*$$

$$\begin{aligned} \mathcal{L}_\Phi = & \mathcal{L}_H + \mathcal{L}_M + \mathcal{L}_{HV^2} + \frac{1}{4}\lambda v^4 \\ & + (\partial_\mu \phi^+) (\partial^\mu \phi^-) + \frac{1}{2} (\partial_\mu \chi) (\partial^\mu \chi) \\ & + iM_W (W_\mu \partial^\mu \phi^+ - W_\mu^\dagger \partial^\mu \phi^-) + M_Z Z_\mu \partial^\mu \chi \\ & + \text{trilinear interactions [SSS, SSV, SVV]} \\ & + \text{quadrilinear interactions [SSSS, SSVV]} \end{aligned}$$

Electroweak symmetry breaking

gauge fixing

- To remove the cross terms $W_\mu \partial^\mu \phi^+$, $W_\mu^\dagger \partial^\mu \phi^-$, $Z_\mu \partial^\mu \chi$ and define propagators add:

$$\mathcal{L}_{\text{GF}} = -\frac{1}{2\xi_\gamma}(\partial_\mu A^\mu)^2 - \frac{1}{2\xi_Z}(\partial_\mu Z^\mu - \xi_Z M_Z \chi)^2 - \frac{1}{\xi_W}|\partial_\mu W^\mu + i\xi_W M_W \phi^-|^2$$

\Rightarrow Massive propagators for gauge and (unphysical) would-be Goldstone fields:

$$\tilde{D}_{\mu\nu}^\gamma(k) = \frac{i}{k^2 + i0} \left[-g_{\mu\nu} + (1 - \xi_\gamma) \frac{k_\mu k_\nu}{k^2} \right]$$

$$\tilde{D}_{\mu\nu}^Z(k) = \frac{i}{k^2 - M_Z^2 + i0} \left[-g_{\mu\nu} + (1 - \xi_Z) \frac{k_\mu k_\nu}{k^2 - \xi_Z M_Z^2} \right] ; \quad \tilde{D}^\chi(k) = \frac{i}{k^2 - \xi_Z M_Z^2 + i0}$$

$$\tilde{D}_{\mu\nu}^W(k) = \frac{i}{k^2 - M_W^2 + i0} \left[-g_{\mu\nu} + (1 - \xi_W) \frac{k_\mu k_\nu}{k^2 - \xi_W M_W^2} \right] ; \quad \tilde{D}^\phi(k) = \frac{i}{k^2 - \xi_W M_W^2 + i0}$$

('t Hooft-Feynman gauge: $\xi_\gamma = \xi_Z = \xi_W = 1$)

Electroweak symmetry breaking

Faddeev-Popov ghosts

- The SM is a non-Abelian theory \Rightarrow add Faddeev-Popov ghosts $c_i(x)$ ($i = 1, 2, 3$)

$$c_1 \equiv \frac{1}{\sqrt{2}}(u_+ + u_-) , \quad c_2 \equiv \frac{i}{\sqrt{2}}(u_+ - u_-) , \quad c_3 \equiv c_W u_Z - s_W u_\gamma$$

$$\mathcal{L}_{\text{FP}} = \underbrace{(\partial^\mu \bar{c}_i)(\partial_\mu c_i - g\epsilon_{ijk}c_j W_\mu^k)}_{\text{U kinetic} + [\text{UV}]} + \underbrace{\text{interactions with } \Phi}_{\text{U masses} + [\text{SUU}]}$$

\Rightarrow Massive propagators for (unphysical) FP ghost fields:

$$\tilde{D}^{u_\gamma}(k) = \frac{i}{k^2 + i0} , \quad \tilde{D}^{u_Z}(k) = \frac{i}{k^2 - \xi_Z M_Z^2 + i0} , \quad \tilde{D}^{u_\pm}(k) = \frac{i}{k^2 - \xi_W M_W^2 + i0}$$

('t Hooft-Feynman gauge: $\xi_Z = \xi_W = 1$)

Electroweak symmetry breaking

Faddeev-Popov ghosts

$$\mathcal{L}_{\text{FP}} = (\partial_\mu \bar{u}_\gamma)(\partial^\mu u_\gamma) + (\partial_\mu \bar{u}_Z)(\partial^\mu u_Z) + (\partial_\mu \bar{u}_+)(\partial^\mu u_+) + (\partial_\mu \bar{u}_-)(\partial^\mu u_-)$$

$$[\text{UV}] \left\{ \begin{array}{l} + ie[(\partial^\mu \bar{u}_+)u_+ - (\partial^\mu \bar{u}_-)u_-]A_\mu - \frac{iec_W}{s_W}[(\partial^\mu \bar{u}_+)u_+ - (\partial^\mu \bar{u}_-)u_-]Z_\mu \\ - ie[(\partial^\mu \bar{u}_+)u_\gamma - (\partial^\mu \bar{u}_\gamma)u_-]W_\mu^\dagger + \frac{iec_W}{s_W}[(\partial^\mu \bar{u}_+)u_Z - (\partial^\mu \bar{u}_Z)u_-]W_\mu^\dagger \\ + ie[(\partial^\mu \bar{u}_-)u_\gamma - (\partial^\mu \bar{u}_\gamma)u_+]W_\mu - \frac{iec_W}{s_W}[(\partial^\mu \bar{u}_-)u_Z - (\partial^\mu \bar{u}_Z)u_+]W_\mu \\ - \xi_Z M_Z^2 \bar{u}_Z u_Z - \xi_W M_W^2 \bar{u}_+ u_+ - \xi_W M_W^2 \bar{u}_- u_- \end{array} \right.$$

$$[\text{SUU}] \left\{ \begin{array}{l} - e\xi_Z M_Z \bar{u}_Z \left[\frac{1}{2s_W c_W} H u_Z - \frac{1}{2s_W} (\phi^+ u_- + \phi^- u_+) \right] \\ - e\xi_W M_W \bar{u}_+ \left[\frac{1}{2s_W} (H + i\chi) u_+ - \phi^+ \left(u_\gamma - \frac{c_W^2 - s_W^2}{2s_W c_W} u_Z \right) \right] \\ - e\xi_W M_W \bar{u}_- \left[\frac{1}{2s_W} (H - i\chi) u_- - \phi^- \left(u_\gamma - \frac{c_W^2 - s_W^2}{2s_W c_W} u_Z \right) \right] \end{array} \right.$$

Electroweak symmetry breaking

fermion masses

- We need masses for quarks and leptons without breaking gauge symmetry

⇒ Introduce Yukawa interactions:

$$\begin{aligned} \mathcal{L}_Y = & -\lambda_d \begin{pmatrix} \bar{u}_L & \bar{d}_L \end{pmatrix} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} d_R - \lambda_u \begin{pmatrix} \bar{u}_L & \bar{d}_L \end{pmatrix} \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix} u_R \\ & - \lambda_\ell \begin{pmatrix} \bar{\nu}_L & \bar{\ell}_L \end{pmatrix} \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \ell_R - \lambda_\nu \begin{pmatrix} \bar{\nu}_L & \bar{\ell}_L \end{pmatrix} \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix} \nu_R + \text{h.c.} \end{aligned}$$

where $\Phi^c \equiv i\sigma_2\Phi^* = \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix}$ transforms under SU(2) like $\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ 6

⇒ After EW SSB, fermions acquire masses ($\bar{f}f = \bar{f}_L f_R + \bar{f}_R f_L$):

$$\mathcal{L}_Y \supset -\frac{1}{\sqrt{2}}(v + H) \left\{ \lambda_d \bar{d}d + \lambda_u \bar{u}u + \lambda_\ell \bar{\ell}\ell + \lambda_\nu \bar{\nu}\nu \right\} \Rightarrow m_f = \lambda_f \frac{v}{\sqrt{2}}$$

- There are 3 generations of quarks and leptons in Nature. They are identical copies with the same properties under $SU(2)_L \otimes U(1)_Y$ differing only in their masses

\Rightarrow Take a general case of n_G generations and let $u_i^I, d_i^I, \nu_i^I, \ell_i^I$ be the members of family i ($i = 1, \dots, n_G$). Superindex I (interaction basis) was omitted so far

\Rightarrow General gauge invariant Yukawa Lagrangian:

$$\boxed{\mathcal{L}_Y} = - \sum_{ij} \left\{ \begin{pmatrix} \bar{u}_{iL}^I & \bar{d}_{iL}^I \end{pmatrix} \left[\begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \lambda_{ij}^{(d)} d_{jR}^I + \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix} \lambda_{ij}^{(u)} u_{jR}^I \right] \right. \\ \left. + \begin{pmatrix} \bar{\nu}_{iL}^I & \bar{\ell}_{iL}^I \end{pmatrix} \left[\begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} \lambda_{ij}^{(\ell)} \ell_{jR}^I + \begin{pmatrix} \phi^{0*} \\ -\phi^- \end{pmatrix} \lambda_{ij}^{(\nu)} \nu_{jR}^I \right] \right\} + \text{h.c.}$$

where $\lambda_{ij}^{(d)}, \lambda_{ij}^{(u)}, \lambda_{ij}^{(\ell)}, \lambda_{ij}^{(\nu)}$ are arbitrary Yukawa matrices

Additional generations

mass matrices

- After EW SSB, in n_G -dimensional matrix form:

$$\mathcal{L}_Y \supset -\left(1 + \frac{H}{v}\right) \left\{ \bar{\mathbf{d}}_L^I \mathbf{M}_d \mathbf{d}_R^I + \bar{\mathbf{u}}_L^I \mathbf{M}_u \mathbf{u}_R^I + \bar{\mathbf{l}}_L^I \mathbf{M}_\ell \mathbf{l}_R^I + \bar{\nu}_L^I \mathbf{M}_\nu \nu_R^I + \text{h.c.} \right\}$$

with mass matrices

$$(\mathbf{M}_d)_{ij} \equiv \lambda_{ij}^{(d)} \frac{v}{\sqrt{2}} \quad (\mathbf{M}_u)_{ij} \equiv \lambda_{ij}^{(u)} \frac{v}{\sqrt{2}} \quad (\mathbf{M}_\ell)_{ij} \equiv \lambda_{ij}^{(\ell)} \frac{v}{\sqrt{2}} \quad (\mathbf{M}_\nu)_{ij} \equiv \lambda_{ij}^{(\nu)} \frac{v}{\sqrt{2}}$$

- ⇒ Diagonalization determines mass eigenstates d_j, u_j, ℓ_j, ν_j in terms of interaction states $d_j^I, u_j^I, \ell_j^I, \nu_j^I$, respectively
- ⇒ Each \mathbf{M}_f can be written as

$$\mathbf{M}_f = \mathbf{H}_f \mathcal{U}_f = \mathbf{V}_f^\dagger \mathcal{M}_f \mathbf{V}_f \mathcal{U}_f \iff \mathbf{M}_f \mathbf{M}_f^\dagger = \mathbf{H}_f^2 = \mathbf{V}_f^\dagger \mathcal{M}_f^2 \mathbf{V}_f$$

- with $\mathbf{H}_f \equiv \sqrt{\mathbf{M}_f \mathbf{M}_f^\dagger}$ a Hermitian positive definite matrix and \mathcal{U}_f unitary
 - Every \mathbf{H}_f can be diagonalized by a unitary matrix \mathbf{V}_f
 - The resulting \mathcal{M}_f is diagonal and positive definite

- In terms of diagonal mass matrices (mass eigenstate basis):

$$\mathcal{M}_d = \text{diag}(m_d, m_s, m_b, \dots), \quad \mathcal{M}_u = \text{diag}(m_u, m_c, m_t, \dots)$$

$$\mathcal{M}_\ell = \text{diag}(m_e, m_\mu, m_\tau, \dots), \quad \mathcal{M}_\nu = \text{diag}(m_{\nu_e}, m_{\nu_\mu}, m_{\nu_\tau}, \dots)$$

$$\mathcal{L}_Y \supset - \left(1 + \frac{H}{v} \right) \left\{ \bar{\mathbf{d}} \mathcal{M}_d \mathbf{d} + \bar{\mathbf{u}} \mathcal{M}_u \mathbf{u} + \bar{\mathbf{l}} \mathcal{M}_\ell \mathbf{l} + \bar{\boldsymbol{\nu}} \mathcal{M}_\nu \boldsymbol{\nu} \right\}$$

where fermion couplings to Higgs are proportional to masses and

$$\begin{aligned} \mathbf{d}_L &\equiv \mathbf{V}_d \ \mathbf{d}_L^I & \mathbf{u}_L &\equiv \mathbf{V}_u \ \mathbf{u}_L^I & \mathbf{l}_L &\equiv \mathbf{V}_\ell \ \mathbf{l}_L^I & \boldsymbol{\nu}_L &\equiv \mathbf{V}_\nu \ \boldsymbol{\nu}_L^I \\ \mathbf{d}_R &\equiv \mathbf{V}_d \mathcal{U}_d \ \mathbf{d}_R^I & \mathbf{u}_R &\equiv \mathbf{V}_u \mathcal{U}_u \ \mathbf{u}_R^I & \mathbf{l}_R &\equiv \mathbf{V}_\ell \mathcal{U}_\ell \ \mathbf{l}_R^I & \boldsymbol{\nu}_R &\equiv \mathbf{V}_\nu \mathcal{U}_\nu \ \boldsymbol{\nu}_R^I \end{aligned}$$

$$\Rightarrow \left. \begin{array}{l} \text{Neutral Currents preserve chirality} \\ \bar{\mathbf{f}}_L^I \mathbf{f}_L^I = \bar{\mathbf{f}}_L \mathbf{f}_L \text{ and } \bar{\mathbf{f}}_R^I \mathbf{f}_R^I = \bar{\mathbf{f}}_R \mathbf{f}_R \end{array} \right\} \Rightarrow \mathcal{L}_{\text{NC}} \text{ does not change flavor}$$

\Rightarrow GIM mechanism

[Glashow, Iliopoulos, Maiani '70]

Additional generations

quark sector


- However, in Charged Currents (also chirality preserving and only LH):

$$\bar{\mathbf{u}}_L^I \mathbf{d}_L^I = \bar{\mathbf{u}}_L \mathbf{V}_u \mathbf{V}_d^\dagger \mathbf{d}_L = \bar{\mathbf{u}}_L \mathbf{V} \mathbf{d}_L$$

with $\mathbf{V} \equiv \mathbf{V}_u \mathbf{V}_d^\dagger$ the (unitary) CKM mixing matrix

[Cabibbo '63; Kobayashi, Maskawa '73]

$$\Rightarrow \quad \mathcal{L}_{CC} = \frac{g}{2\sqrt{2}} \sum_{ij} \bar{u}_i \gamma^\mu (1 - \gamma_5) \mathbf{V}_{ij} d_j W_\mu^\dagger + \text{h.c.}$$



- ⇒ If u_i or d_j had degenerate masses one could choose $\mathbf{V}_u = \mathbf{V}_d$ (field redefinition) and flavor would be conserved in the quark sector. But they are not degenerate
- ⇒ \mathbf{V}_u and \mathbf{V}_d are not observable. Just masses and CKM mixings are observable

Additional generations

quark sector

- How many physical parameters in this sector?
 - Quark masses and CKM mixings determined by mass (or Yukawa) matrices
 - A general $n_G \times n_G$ unitary matrix, like the CKM, is given by

$$n_G^2 \text{ real parameters} = n_G(n_G - 1)/2 \text{ moduli} + n_G(n_G + 1)/2 \text{ phases}$$

Some phases are unphysical since they can be absorbed by field redefinitions:

$$u_i \rightarrow e^{i\phi_i} u_i, \quad d_j \rightarrow e^{i\theta_j} d_j \quad \Rightarrow \quad V_{ij} \rightarrow V_{ij} e^{i(\theta_j - \phi_i)}$$

Therefore $2n_G - 1$ unphysical phases and the physical parameters are:

$$(n_G - 1)^2 = n_G(n_G - 1)/2 \text{ moduli} + (n_G - 1)(n_G - 2)/2 \text{ phases}$$

Additional generations

quark sector

⇒ Case of $n_G = 2$ generations: 1 parameter, the Cabibbo angle θ_C :

$$\mathbf{V} = \begin{pmatrix} \cos \theta_C & \sin \theta_C \\ -\sin \theta_C & \cos \theta_C \end{pmatrix}$$

⇒ Case of $n_G = 3$ generations: 3 angles + 1 phase. In the standard parameterization:

$$\begin{aligned} \mathbf{V} &= \begin{pmatrix} \mathbf{V}_{ud} & \mathbf{V}_{us} & \mathbf{V}_{ub} \\ \mathbf{V}_{cd} & \mathbf{V}_{cs} & \mathbf{V}_{cb} \\ \mathbf{V}_{td} & \mathbf{V}_{ts} & \mathbf{V}_{tb} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \Rightarrow \begin{matrix} \text{δ only source} \\ \text{of CP violation} \\ \text{in the SM !} \end{matrix} \end{aligned}$$

with $c_{ij} \equiv \cos \theta_{ij} \geq 0$, $s_{ij} \equiv \sin \theta_{ij} \geq 0$ ($i < j = 1, 2, 3$) and $0 \leq \delta \leq 2\pi$

- If neutrinos were massless we could redefine the (LH) fields \Rightarrow no lepton mixing
But they have (tiny) masses because there are neutrino oscillations!
- Neutrinos are special:
they *may* be their own antiparticle (Majorana) since they are neutral
- *If* they are Majorana:
 - Mass terms are different to Dirac case
(neutrino and antineutrino *may* mix)
 - Intergenerational mixings are richer (more CP phases)



lepton sector

- About Majorana fermions

- A **Dirac fermion** field is a spinor with **4** independent components: 2 LH+2 RH (left/right-handed particles and antiparticles)

$$\psi_L = P_L \psi, \quad \psi_R = P_R \psi, \quad \psi_L^c \equiv (\psi_L)^c = P_R \psi^c, \quad \psi_R^c \equiv (\psi_R)^c = P_L \psi^c$$

where $\psi^c \equiv C \bar{\psi}^\top = i\gamma^2 \psi^*$ (charge conjugate) with $C = i\gamma^2 \gamma^0$, $P_{R,L} = \frac{1}{2}(1 \pm \gamma_5)$

- A **Majorana fermion** field has just **2** independent components since $\psi^c \equiv \eta^* \psi$:

$$\psi_L = \eta \psi_R^c, \quad \psi_R = \eta \psi_L^c$$

where $\eta = -i\eta_{CP}$ (CP parity) with $|\eta|^2 = 1$. Only possible if neutral



lepton sector

- About mass terms

$$\begin{aligned} \overline{\psi_R}\psi_L &= \overline{\psi_L^c}\psi_R^c \quad , \quad \overline{\psi_L}\psi_R = \overline{\psi_R^c}\psi_L^c \quad (\Delta F = 0) \\ \overline{\psi_L^c}\psi_L &\quad , \quad \overline{\psi_L}\psi_L^c \quad \left. \right\} \quad (|\Delta F| = 2) \\ \overline{\psi_R}\psi_R^c &\quad , \quad \overline{\psi_R^c}\psi_R \end{aligned}$$
$$\Rightarrow -\mathcal{L}_m = \underbrace{m_D \overline{\psi_R}\psi_L}_{\text{Dirac term}} + \underbrace{\frac{1}{2}m_L \overline{\psi_L^c}\psi_L + \frac{1}{2}m_R \overline{\psi_R}\psi_R^c}_{\text{Majorana terms}} + \text{h.c.}$$

- A Dirac fermion can only have Dirac mass term
- A Majorana fermion can have **both** Dirac and Majorana mass terms

\Rightarrow In the SM:

- * m_D from Yukawa coupling after EW SSB $(m_D = \lambda_\nu v / \sqrt{2})$
- * m_L forbidden by gauge symmetry
- * m_R compatible with gauge symmetry!



lepton sector

- About mass terms (a more transparent parameterization)

Rewrite previous mass terms introducing an array of two Majorana fermions:

$$\chi^0 = \chi^{0c} = \chi_L^0 + \chi_L^{0c} \equiv \begin{pmatrix} \chi_1^0 \\ \chi_2^0 \end{pmatrix}, \quad \begin{aligned} \chi_1^0 &= \chi_1^{0c} = \chi_{1L}^0 + \chi_{1L}^{0c} \equiv \psi_L + \psi_L^c \\ \chi_2^0 &= \chi_2^{0c} = \chi_{2L}^0 + \chi_{2L}^{0c} \equiv \psi_R + \psi_R^c \end{aligned}$$

$$\Rightarrow -\mathcal{L}_m = \frac{1}{2} \overline{\chi_L^{0c}} \mathbf{M} \chi_L^0 + \text{h.c.} \quad \text{with} \quad \mathbf{M} = \begin{pmatrix} m_L & m_D \\ m_D & m_R \end{pmatrix}$$

\mathbf{M} is a square symmetric matrix \Rightarrow diagonalizable by a unitary matrix $\tilde{\mathcal{U}}$:

$$\tilde{\mathcal{U}}^\top \mathbf{M} \tilde{\mathcal{U}} = \mathcal{M} = \text{diag}(m'_1, m'_2), \quad \chi_L^0 = \tilde{\mathcal{U}} \chi_L \quad (\chi_L^{0c} = \tilde{\mathcal{U}}^* \chi_L^c)$$

To get real and positive eigenvalues $m_i = \eta_i m'_i$ (physical masses) take $\chi_L^0 = \mathcal{U} \xi_L$:

$$\mathcal{U} = \tilde{\mathcal{U}} \text{diag}(\sqrt{\eta_1}, \sqrt{\eta_2}), \quad \begin{aligned} \xi_1 &= \chi_{1L} + \eta_1 \chi_{1L}^c && \text{(physical fields)} \quad \eta_i = \text{CP parities} \\ \xi_2 &= \chi_{2L} + \eta_2 \chi_{2L}^c \end{aligned}$$



lepton sector

- About mass terms (a more transparent parameterization)
 - Case of **only Dirac term** ($m_L = m_R = 0$)

$$\mathbf{M} = \begin{pmatrix} 0 & m_D \\ m_D & 0 \end{pmatrix} \Rightarrow \tilde{\mathcal{U}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}, \quad m'_1 = -m_D, \quad m'_2 = m_D$$

Eigenstates

\Rightarrow Physical states

$$\chi_{1L} = \frac{1}{\sqrt{2}}(\chi_{1L}^0 - \chi_{2L}^0) = \frac{1}{\sqrt{2}}(\psi_L - \psi_R^c)$$

$$\chi_{2L} = \frac{1}{\sqrt{2}}(\chi_{1L}^0 + \chi_{2L}^0) = \frac{1}{\sqrt{2}}(\psi_L + \psi_R^c)$$

$$\xi_1 = \chi_{1L} + \eta_1 \chi_{1L}^c \quad [\eta_1 = -1]$$

$$\xi_2 = \chi_{2L} + \eta_2 \chi_{2L}^c \quad [\eta_2 = +1]$$

with masses $m_1 = m_2 = m_D$

$$\Rightarrow -\mathcal{L}_m = \frac{1}{2}m_D(-\bar{\chi}_1\chi_1 + \bar{\chi}_2\chi_2) = \frac{1}{2}m_D(\bar{\xi}_1\xi_1 + \bar{\xi}_2\xi_2) = m_D(\bar{\psi}_L\psi_R + \bar{\psi}_R\psi_L)$$

One Dirac fermion = two Majorana of equal mass and opposite CP parities




lepton sector

- About mass terms (a more transparent parameterization)
 - Case of **seesaw** (type I) [Yanagida '79; Gell-Mann, Ramond, Slansky '79; Mohapatra, Senjanovic '80]
 $(m_L = 0, m_D \ll m_R)$

$$\mathbf{M} = \begin{pmatrix} 0 & m_D \\ m_D & m_R \end{pmatrix} \Rightarrow \tilde{\mathcal{U}} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}, \quad \theta \simeq \frac{m_D}{m_R} \simeq \sqrt{\frac{m_\nu}{m_N}} \text{ (negligible)}$$

$$m_1 \equiv m_\nu \simeq \frac{m_D^2}{m_R} \ll m_2 \equiv m_N \simeq m_R$$



$$\begin{aligned} \xi_1 \equiv \nu &= \psi_L + \eta_1 \psi_L^c \quad [\eta_1 = -1] \\ \xi_2 \equiv N &= \psi_R^c + \eta_2 \psi_R \quad [\eta_2 = +1] \end{aligned} \Rightarrow -\mathcal{L}_m = \frac{1}{2} m_\nu \overline{\nu_L^c} \nu_L + \frac{1}{2} m_N \overline{N_R^c} N_R + \text{h.c.}$$

Perhaps the observed neutrino ν_L is the LH component of a light Majorana ν (then $\bar{\nu} = \text{RH}$) and light because of a very heavy Majorana neutrino N

$$\text{e.g. } m_D \sim v \simeq 246 \text{ GeV}, \quad m_R \sim m_N \sim 10^{15} \text{ GeV} \quad \Rightarrow \quad m_\nu \sim 0.1 \text{ eV} \quad \checkmark$$

Additional generations

lepton sector

- Lepton mixings

- From Z lineshape: there are $n_G = 3$ generations of ν_L [ν_i ($i = 1, \dots, n_G$)] (but we do not know (*yet*) if neutrinos are Dirac or Majorana fermions)
- From neutrino oscillations: neutrinos are light, non degenerate and mix

$$|\nu_\alpha\rangle = \sum_i \mathbf{U}_{\alpha i} |\nu_i\rangle \iff |\nu_i\rangle = \sum_\alpha \mathbf{U}_{\alpha i}^* |\nu_\alpha\rangle$$

mass eigenstates ν_i ($i = 1, 2, 3$) / interaction states ν_α ($\alpha = e, \mu, \tau$)

- ⇒ \mathbf{U} matrix is unitary (negligible mixing with heavy neutrinos) and analogous to \mathbf{V}_u , \mathbf{V}_d , \mathbf{V}_ℓ defined for quarks and charged leptons except for:
- ν fields have both chiralities
 - *If* neutrinos are Majorana, \mathbf{U} *may* contain two additional physical (Majorana) phases (irrelevant and therefore not measurable in oscillation experiments) that cannot be absorbed since then field phases are fixed by $\nu_i = \eta_i \nu_i^c$

Additional generations

lepton sector

- Lepton mixings


The so called PMNS matrix \mathbf{U}

[Pontecorvo '57; Maki, Nakagawa, Sakata '62; Pontecorvo '68]

- does not change Neutral Currents (unitarity), but
- introduces intergenerational mixings in Charged Currents:

$$\mathcal{L}_{\text{CC}} = \frac{g}{2\sqrt{2}} \sum_{\alpha i} \bar{\ell}_\alpha \gamma^\mu (1 - \gamma_5) \mathbf{U}_{\alpha i} \nu_i W_\mu + \text{h.c.}$$

(basis where charged leptons are diagonal)



Additional generations

lepton sector

⇒ Standard parameterization of the PMNS matrix:

$$\mathbf{U} = \begin{pmatrix} \mathbf{U}_{e1} & \mathbf{U}_{e2} & \mathbf{U}_{e3} \\ \mathbf{U}_{\mu 1} & \mathbf{U}_{\mu 2} & \mathbf{U}_{\mu 3} \\ \mathbf{U}_{\tau 1} & \mathbf{U}_{\tau 2} & \mathbf{U}_{\tau 3} \end{pmatrix}$$

$$= \begin{pmatrix} c_{12} c_{13} & s_{12} c_{13} & s_{13} e^{-i\delta} \\ -s_{12} c_{23} - c_{12} s_{23} s_{13} e^{i\delta} & c_{12} c_{23} - s_{12} s_{23} s_{13} e^{i\delta} & s_{23} c_{13} \\ s_{12} s_{23} - c_{12} c_{23} s_{13} e^{i\delta} & -c_{12} s_{23} - s_{12} c_{23} s_{13} e^{i\delta} & c_{23} c_{13} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha_{21}/2} & 0 \\ 0 & 0 & e^{i\alpha_{31}/2} \end{pmatrix}$$

(different values than in CKM)

(Majorana phases)

$[\theta_{13} \equiv \theta_\odot, \quad \theta_{23} \equiv \theta_{\text{atm}}, \quad \theta_{13} \quad \text{and} \quad (\text{almost}) \delta \quad \text{measured in oscillations}]$

3. Phenomenology

$$\mathcal{L} = \mathcal{L}_F + \mathcal{L}_{\text{YM}} + \mathcal{L}_\Phi + \mathcal{L}_Y + \mathcal{L}_{\text{GF}} + \mathcal{L}_{\text{FP}}$$

- Fields: [F] fermions [S] scalars (Higgs and unphysical Goldstones)
[V] vector bosons [U] unphysical ghosts
- Interactions: [FFV] [FFS] [SSV] [SVV] [SSVV]
[VVV] [VVVV] [SSS] [SSSS]
[SUU] [UUUV]

- Feynman rules for generic couplings normalized to e (all momenta incoming):

(i \mathcal{L})

$$[\text{FFV}_\mu] \quad ie\gamma^\mu(g_V - g_A\gamma_5) = ie\gamma^\mu(g_L P_L + g_R P_R)$$

$$[\text{FFS}] \quad ie(g_S - g_P\gamma_5) = ie(c_L P_L + c_R P_R)$$

$$[\text{SV}_\mu \text{V}_\nu] \quad ieKg_{\mu\nu}$$

$$[S(p_1)S(p_2)\text{V}_\mu] \quad ieG(p_1 - p_2)_\mu$$

$$[\text{V}_\mu(k_1)\text{V}_\nu(k_2)\text{V}_\rho(k_3)] \quad ieJ [g_{\mu\nu}(k_2 - k_1)_\rho + g_{\nu\rho}(k_3 - k_2)_\mu + g_{\mu\rho}(k_1 - k_3)_\nu]$$

$$[\text{V}_\mu(k_1)\text{V}_\nu(k_2)\text{V}_\rho(k_3)\text{V}_\sigma(k_4)] \quad ie^2 C_1 [2g_{\mu\nu}g_{\rho\sigma} - g_{\mu\rho}g_{\nu\sigma} - g_{\mu\sigma}g_{\nu\rho}]$$

$$[\text{SSV}_\mu \text{V}_\nu] \quad ie^2 C_2 g_{\mu\nu} \quad \text{also } [\text{UUVV}]$$

$$[\text{SSS}] \quad ieC_3 \quad \text{also } [\text{SUU}]$$

$$[\text{SSSS}] \quad ie^2 C_4$$

Note: $g_{L,R} = g_V \pm g_A$ $\partial_\mu \rightarrow -ip_\mu$

Attention to symmetry factors!

 $c_{L,R} = g_S \pm g_P$ e.g. $2 \times HZZ$

Complete SM Lagrangian

Feynman rules

('t Hooft-Feynman gauge)

| FFV | $\bar{f}_i f_j \gamma$ | $\bar{f}_i f_j Z$ | $\bar{u}_i d_j W^+$ | $\bar{d}_j u_i W^-$ | $\bar{\nu}_i \ell_j W^+$ | $\bar{\ell}_j \nu_i W^-$ |
|-------|------------------------|---------------------|---|---|---|---|
| g_L | $-Q_f \delta_{ij}$ | $g_+^f \delta_{ij}$ | $\frac{1}{\sqrt{2}s_W} \mathbf{V}_{ij}$ | $\frac{1}{\sqrt{2}s_W} \mathbf{V}_{ij}^*$ | $\frac{1}{\sqrt{2}s_W} \mathbf{U}_{ji}^*$ | $\frac{1}{\sqrt{2}s_W} \mathbf{U}_{ji}$ |
| g_R | $-Q_f \delta_{ij}$ | $g_-^f \delta_{ij}$ | 0 | 0 | 0 | 0 |

$$g_{\pm}^f \equiv v_f \pm a_f \quad v_f = \frac{T_3^{f_L} - 2Q_f s_W^2}{2s_W c_W} \quad a_f = \frac{T_3^{f_L}}{2s_W c_W}$$

Complete SM Lagrangian

Feynman rules

('t Hooft-Feynman gauge)

| FFS | $\bar{f}_i f_j H$ | $\bar{f}_i f_j \chi$ | $\bar{u}_i d_j \phi^+$ | $\bar{d}_j u_i \phi^-$ |
|-------|---|--|--|--|
| c_L | $-\frac{1}{2s_W} \frac{m_{f_i}}{M_W} \delta_{ij}$ | $-\frac{i}{2s_W} 2T_3^{f_L} \frac{m_{f_i}}{M_W} \delta_{ij}$ | $+\frac{1}{\sqrt{2}s_W} \frac{m_{u_i}}{M_W} \mathbf{V}_{ij}$ | $-\frac{1}{\sqrt{2}s_W} \frac{m_{d_j}}{M_W} \mathbf{V}_{ij}^*$ |
| c_R | $-\frac{1}{2s_W} \frac{m_{f_i}}{M_W} \delta_{ij}$ | $+\frac{i}{2s_W} 2T_3^{f_L} \frac{m_{f_i}}{M_W} \delta_{ij}$ | $-\frac{1}{\sqrt{2}s_W} \frac{m_{d_j}}{M_W} \mathbf{V}_{ij}$ | $+\frac{1}{\sqrt{2}s_W} \frac{m_{u_j}}{M_W} \mathbf{V}_{ij}^*$ |

$$(f = u, d, \ell)$$

| FFS | $\bar{\nu}_i \ell_j \phi^+$ | $\bar{\ell}_j \nu_i \phi^-$ |
|-------|---|---|
| c_L | $+\frac{1}{\sqrt{2}s_W} \frac{m_{\nu_i}}{M_W} \mathbf{U}_{ji}^*$ | $-\frac{1}{\sqrt{2}s_W} \frac{m_{\ell_j}}{M_W} \mathbf{U}_{ji}$ |
| c_R | $-\frac{1}{\sqrt{2}s_W} \frac{m_{\ell_j}}{M_W} \mathbf{U}_{ji}^*$ | $+\frac{1}{\sqrt{2}s_W} \frac{m_{\nu_i}}{M_W} \mathbf{U}_{ji}$ |

Complete SM Lagrangian

Feynman rules

('t Hooft-Feynman gauge)

| SVV | HZZ | HW^+W^- | $\phi^\pm W^\mp \gamma$ | $\phi^\pm W^\mp Z$ |
|-----|-------------------|-----------|-------------------------|--------------------|
| K | $M_W/(s_W c_W^2)$ | M_W/s_W | $-M_W$ | $-M_W s_W/c_W$ |

| SSV | χHZ | $\phi^\pm \phi^\mp \gamma$ | $\phi^\pm \phi^\mp Z$ | $\phi^\mp HW^\pm$ | $\phi^\mp \chi W^\pm$ |
|-----|-----------------------|----------------------------|--------------------------------------|----------------------|-----------------------|
| G | $-\frac{i}{2s_W c_W}$ | ∓ 1 | $\pm \frac{c_W^2 - s_W^2}{2s_W c_W}$ | $\mp \frac{1}{2s_W}$ | $-\frac{i}{2s_W}$ |

| VVV | γW^+W^- | ZW^+W^- |
|-----|-----------------|-----------|
| J | -1 | c_W/s_W |

Complete SM Lagrangian

Feynman rules

('t Hooft-Feynman gauge)

| VVVV | $W^+W^+W^-W^-$ | W^+W^-ZZ | $W^+W^-\gamma Z$ | $W^+W^-\gamma\gamma$ |
|-------|-------------------|------------------------|-------------------|----------------------|
| C_1 | $\frac{1}{s_W^2}$ | $-\frac{c_W^2}{s_W^2}$ | $\frac{c_W}{s_W}$ | -1 |

| SSVV | HHW^-W^+ | $HHZZ$ |
|-------|--------------------|-------------------------|
| C_2 | $\frac{1}{2s_W^2}$ | $\frac{1}{2s_W^2c_W^2}$ |

| SSS | HHH |
|-------|----------------------------|
| C_3 | $-\frac{3M_H^2}{2M_W s_W}$ |

| SSSS | $HHHH$ |
|-------|--------------------------------|
| C_4 | $-\frac{3M_H^2}{4M_W^2 s_W^2}$ |

- Would-be Goldstone bosons in [SSVV], [SSS] and [SSSS] omitted
- Faddeev-Popov ghosts in [SUU] and [UUUV] omitted
- All Feynman rules from **FeynArts** (same conventions; $\chi, \phi^\pm \rightarrow G^0, G^\pm$):

<http://www.ugr.es/local/jillana/SM/FeynmanRulesSM.pdf>

Input parameters

- Parameters:

| | | | | | | | |
|------------|----------|-------|-------|-----------|-------------|---------------------------|----------------------------|
| $17+9 =$ | 1 | 1 | 1 | 1 | $9+3$ | 4 | 6 |
| formal: | g | g' | v | λ | λ_f | | |
| practical: | α | M_W | M_Z | M_H | m_f | \mathbf{V}_{CKM} | \mathbf{U}_{PMNS} |

where $g = \frac{e}{s_W}$ $g' = \frac{e}{c_W}$ and

$$\underbrace{\alpha = \frac{e^2}{4\pi} \quad M_W = \frac{1}{2}gv \quad M_Z = \frac{M_W}{c_W}}_{g, g', v} \quad M_H = \sqrt{2\lambda}v \quad m_f = \frac{v}{\sqrt{2}}\lambda_f$$

⇒ Many (more) experiments

⇒ After Higgs discovery, for the first time *all* parameters measured!

Input parameters

- Experimental values

[Particle Data Group '18]

- Fine structure constant:

$$\alpha^{-1} = 137.035\,999\,139\,(31) \quad \text{from Harvard cyclotron } (g_e)$$

- The SM predicts $M_W < M_Z$ in agreement with measurements:

$$M_Z = (91.1876 \pm 0.0021) \text{ GeV} \quad \text{from LEP1/SLD}$$

$$M_W = (80.379 \pm 0.012) \text{ GeV} \quad \text{from LEP2/Tevatron/LHC}$$

- Top quark mass:

$$m_t = (173.0 \pm 0.4) \text{ GeV} \quad \text{from Tevatron/LHC}$$

- Higgs boson mass:

$$M_H = (125.18 \pm 0.16) \text{ GeV} \quad \text{from LHC}$$

- ...

Observables and experiments

- Low energy observables ($Q^2 \ll M_Z^2$)

– ν -nucleon (NuTeV) and νe (CERN) scattering:


asymmetries CC/NC and $\nu/\bar{\nu} \Rightarrow s_W^2$

– Atomic parity violation (SLAC, CERN, Jefferson Lab, Mainz):

LR asymmetries $e_{R,L}N \rightarrow eX$ due to Z-exchange between e and $N \Rightarrow s_W^2$

– muon decay: $\mu \rightarrow e \bar{\nu}_e \nu_\mu$ (PSI)
lifetime

$$\frac{1}{\tau_\mu} = \Gamma_\mu = \frac{G_F^2 m_\mu^5}{192\pi^3} f(m_e^2/m_\mu^2)$$



$$f(x) \equiv 1 - 8x + 8x^3 - x^4 - 12x^2 \ln x \Rightarrow G_F$$

$$= 0.99981295$$

$$i\mathcal{M} = \left(\frac{ie}{\sqrt{2}s_W} \right)^2 \bar{e} \gamma^\rho \nu_L \frac{-ig_{\rho\delta}}{q^2 - M_W^2} \bar{\nu}_L \gamma^\delta \mu \equiv \overbrace{i \frac{4G_F}{\sqrt{2}} (\bar{e} \gamma^\rho \nu_L)(\bar{\nu}_L \gamma_\rho \mu)}^{\text{Fermi theory } (-q^2 \ll M_W^2)} ; \frac{G_F}{\sqrt{2}} = \frac{\pi \alpha}{2s_W^2 M_W^2}$$

Observables and experiments

- Low energy observables

⇒ Fermi constant provides the Higgs VEV (electroweak scale):

$$v = \left(\sqrt{2} G_F \right)^{-1/2} \approx 246 \text{ GeV}$$

⇒ Consistency checks: e.g.

From muon lifetime:

$$G_F = 1.166\,378\,7(6) \times 10^{-5} \text{ GeV}^{-2}$$

If one compares with (tree level result)


$$\frac{G_F}{\sqrt{2}} = \frac{\pi \alpha}{2 s_W^2 M_W^2} = \frac{\pi \alpha}{2(1 - M_W^2/M_Z^2) M_W^2}$$

using measurements of M_W , M_Z and α there is a discrepancy that disappears when *quantum corrections* are included

Observables and experiments

- $e^+e^- \rightarrow \bar{f}f$ (PEP, PETRA, TRISTAN, ..., LEP1, SLD)

8



$$\frac{d\sigma}{d\Omega} = N_c^f \frac{\alpha^2}{4s} \beta_f \left\{ \left[1 + \cos^2 \theta + (1 - \beta_f^2) \sin^2 \theta \right] G_1(s) + 2(\beta_f^2 - 1) G_2(s) + 2\beta_f \cos \theta G_3(s) \right\}$$

$$G_1(s) = Q_e^2 Q_f^2 + 2Q_e Q_f v_e v_f \operatorname{Re} \chi_Z(s) + (v_e^2 + a_e^2)(v_f^2 + a_f^2) |\chi_Z(s)|^2$$

$$G_2(s) = (v_e^2 + a_e^2)a_f^2 |\chi_Z(s)|^2$$

$$G_3(s) = 2Q_e Q_f a_e a_f \operatorname{Re} \chi_Z(s) + 4v_e v_f a_e a_f |\chi_Z(s)|^2 \Rightarrow A_{FB}(s)$$

with $\chi_Z(s) \equiv \frac{s}{s - M_Z^2 + iM_Z\Gamma_Z}$, $N_c^f = 1$ (3) for $f = \text{lepton (quark)}$, $\beta_f = \text{velocity}$


$$\sigma(s) = N_c^f \frac{2\pi\alpha^2}{3s} \beta_f \left[(3 - \beta_f^2) G_1(s) - 3(1 - \beta_f^2) G_2(s) \right], \quad \beta_f = \sqrt{1 - 4m_f^2/s}$$

Observables and experiments

- **Z production** (LEP1/SLD)

$$M_Z, \Gamma_Z, \sigma_{\text{had}}, A_{FB}, A_{LR}, R_b, R_c, R_\ell \quad \Rightarrow \boxed{M_Z, s_W^2}$$

from $e^+e^- \rightarrow f\bar{f}$ at the Z pole ($\gamma - Z$ interference vanishes). Neglecting m_f :



$$\sigma_{\text{had}} = 12\pi \frac{\Gamma(e^+e^-)\Gamma(\text{had})}{M_Z^2 \Gamma_Z^2}$$

$$R_b = \frac{\Gamma(b\bar{b})}{\Gamma(\text{had})} \quad R_c = \frac{\Gamma(c\bar{c})}{\Gamma(\text{had})} \quad R_\ell = \frac{\Gamma(\text{had})}{\Gamma(\ell^+\ell^-)}$$

$$\left[\Gamma(Z \rightarrow f\bar{f}) \equiv \Gamma(f\bar{f}) = N_c^f \frac{\alpha M_Z}{3} (v_f^2 + a_f^2) \right]$$

$$\Rightarrow N_\nu = 2.990 \pm 0.007$$

kine 8

Forward-Backward and (if polarized e^-) Left-Right asymmetries due to Z:


$$A_{FB} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B} = \frac{3}{4} A_f \frac{A_e + P_e}{1 + P_e A_e}$$

$$A_{LR} = \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R} = A_e P_e \quad \text{with } A_f \equiv \frac{2v_f a_f}{v_f^2 + a_f^2}$$

Observables and experiments


- W-pair production (LEP2)

$$e^+ e^- \rightarrow WW \rightarrow 4f (+\gamma)$$




- W production (Tevatron/LHC)

$$pp/p\bar{p} \rightarrow W \rightarrow \ell\nu_\ell (+\gamma)$$



- Top-quark production (Tevatron/LHC)

$$pp/p\bar{p} \rightarrow t\bar{t} \rightarrow 6f$$






Observables and experiments

- Higgs (LHC)

Single and Double H production and decay to different channels $\Rightarrow M_H$

[PDG '18]



kine

9

Observables and experiments

- Higgs (LHC)

[PDG '18]



| | Signal strength $\mu = \frac{(\sigma \cdot \text{BR})_{\text{obs}}}{(\sigma \cdot \text{BR})_{\text{SM}}}$ | Run 1 | Run 2 |
|-------|--|------------------------|------------------------|
| ATLAS | | 1.17 ± 0.27 | 0.99 ± 0.14 |
| | | $0.78^{+0.26}_{-0.23}$ | $1.16^{+0.15}_{-0.14}$ |

Per channel:

$\gamma\gamma, ZZ, W^+W^-, \tau^+\tau^- > 5\sigma$

$b\bar{b} > 5\sigma$ at last!

[LHC Seminar 28 Jul 2018]




Precise determination of parameters

- Experimental precision requires accurate predictions \Rightarrow quantum corrections
(complication: loop calculations involve renormalization)
- Correction to G_F from muon lifetime:

$$\frac{G_F}{\sqrt{2}} \rightarrow \frac{G_F}{\sqrt{2}} = \frac{\pi\alpha}{2(1 - M_W^2/M_Z^2)M_W^2} [1 + \Delta r(m_t, M_H)]$$

when loop corrections are included:



Since muon lifetime is measured more precisely than M_W , it is traded for G_F :

$$\Rightarrow M_W^2(\alpha, G_F, M_Z, m_t, M_H) = \frac{M_Z^2}{2} \left(1 + \sqrt{1 - \frac{4\pi\alpha}{\sqrt{2}G_F M_Z^2} [1 + \Delta r(m_t, M_H)]} \right)$$

(correlation between M_W , m_t and M_H , given α , G_F and M_Z)

Precise determination of parameters

Indirect constraints from LEP1/SLD

$$M_H(M_W, m_t)$$


Direct measurements from LEP2/Tevatron

Allowed regions for M_H
allowed by direct searches



LHC excluded

[LEPEWWG 2013]




Precise determination of parameters

- Corrections to vector and axial couplings from Z pole observables:

$$v_f \rightarrow g_V^f = v_f + \Delta g_V^f \quad a_f \rightarrow g_A^f = a_f + \Delta g_A^f$$

$$\Rightarrow \sin^2 \theta_{\text{eff}}^f = \frac{1}{4|Q_f|} \left[1 - \text{Re}(g_V^f/g_A^f) \right] \equiv \overbrace{(1 - M_W^2/M_Z^2)}^{s_W^2} \kappa_Z^f$$

(Two) loop calculations are crucial and point to a light Higgs:



$$s_W^2 = 0.22290 \pm 0.00029 \text{ (tree)}$$

$$\sin^2 \theta_{\text{eff}}^{\text{lept}} = 0.23148 \pm 0.000017 \text{ (exp)}$$

Precise determination of parameters

- In addition, experiments and observables testing the flavor structure of the SM:
flavor conserving: dipole moments, ... flavor changing: $b \rightarrow s\gamma, \dots$
 \Rightarrow very sensitive to new physics through loop corrections

Extremely precise measurements are:

- electron magnetic moment (new physics suppressed by a factor of m_e^2/m_μ^2):

$$\left. \begin{array}{l} \text{exp: } g_e/2 = 1.001\,159\,652\,180\,76 \text{ (27)} \\ \text{theo: QED (8 loops!)} \end{array} \right\} \Rightarrow \alpha^{-1} = 137.035\,999\,139 \text{ (31)}$$

- muon anomalous magnetic moment: $a_\mu = (g_\mu - 2)/2$

| | |
|--|-------------------------------------|
| $a_\mu^{\text{exp}} = 116\,592\,089\,(63) \times 10^{-11}$ | [Brookhaven '06] |
| $a_\mu^{\text{QED}} = 116\,584\,718 \times 10^{-11}$ | [QED: 5 loops] |
| $a_\mu^{\text{EW}} = 154 \times 10^{-11}$ | [W, Z, H: 2 loops] |
| $a_\mu^{\text{had}} = 6\,930\,(48) \times 10^{-11}$ | [$e^+e^- \rightarrow \text{had}$] |
| $a_\mu^{\text{SM}} = 116\,591\,802\,(49) \times 10^{-11}$ | |


$$a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = 287\,(80) \times 10^{-11} \quad 3.6\sigma !$$

Precise determination of parameters

Recent update on $(g_\mu - 2)$

- New lattice calculations of a_μ^{had}

[PRL 121, 022002 & 022003, 12 July 2018]



- New Muon $g - 2$ Experiment at Fermilab (running from March 2018)

Global fits

- Fit input data from a list of observables (EWPO):

$$M_H, M_W, \Gamma_W, M_Z, \Gamma_Z, \sigma_{\text{had}}, A_{FB}^{b,c,\ell}, A_{b,c,\ell}, R_{b,c,\ell}, \sin^2 \theta_{\text{eff}}^{\text{lept}}, \dots$$

finding the χ^2_{min} for $n_{\text{dof}} = 13$ (14) when M_H is included (excluded):

$$\underbrace{\alpha_s(M_Z), \Delta\alpha_{\text{had}}(M_Z), G_F, M_Z, 9 \text{ fermion masses}, M_H}_{\begin{array}{c} 1 \text{ (QCD)} \\ 17-4=13 \text{ (CKM irrelevant)} \end{array}}$$

$$\alpha(M_Z) \equiv \frac{\alpha}{1 - \Delta\alpha(M_Z)}$$


[Gfitter 2018: 1803.0853] <http://gfitter.desy.de>

| n_{dof} | χ^2_{min} | $p\text{-value}$ |
|------------------|-----------------------|------------------|
| 15 | 18.6 | 0.23 |

p -value (goodnes of fit): probability, under assumption of hypothesis H, to observe data with equal or lesser compatibility with H relative to the data we got

Global fits (Comparisons)


- Compare direct measurements of these observables with fit values:



⇒ some tensions (none above 3σ): $A_\ell(\text{SLD})$, $A_{FB}^b(\text{LEP})$, R_b , ...

Global fits (Comparisons)

- Compare indirect determinations with fit values (error bars are direct measmts.):




[indirect determination means fit without using constraint from given direct measurement]

Global fits

(Conclusions)


⇒ Fits prefer a somewhat lighter Higgs:



Global fits

(Conclusions)


⇒ In general, impressive consistency of the SM, e.g.:




Summary

- The SM is a gauge theory with spontaneous symmetry breaking (renormalizable)
- Confirmed by many low and high energy experiments with remarkable accuracy, at the level of quantum corrections, with (almost) no significant deviations
- In spite of its tremendous success, it leaves fundamental questions unanswered:
why 3 generations? why the observed pattern of fermion masses and mixings?
- And there are several hints for physics beyond:
 - phenomenological:
 - * $(g_\mu - 2)$
 - * neutrino masses
 - * baryon asymmetry
 - * dark matter
 - * dark energy
 - * cosmological constant
 - conceptual:
 - * gravity is not included
 - * hierarchy problem

The SM is an Effective Theory
valid up to electroweak scale?



Thank You!




<http://lacienciaconhumor.blogspot.com/2011/08/investigacion-basica.html>



<http://lacienciaconhumor.blogspot.com/2011/08/investigacion-basica.html>

Kinematics

Cross-section




$$d\sigma(i \rightarrow f) = \frac{1}{4 \{(p_1 p_2)^2 - m_1^2 m_2^2\}^{1/2}} |\mathcal{M}|^2 (2\pi)^4 \delta^4(p_i - p_f) \prod_{j=3}^{n+2} \frac{d^3 p_j}{(2\pi)^3 2E_j}$$

- ▷ Sum over initial polarizations and/or average over final polarizations if the initial state is unpolarized and/or the final state polarization is not measured
- ▷ Divide the total cross-section by a symmetry factor $S = \prod_i k_i!$ if there are k_i identical particles of species i in the final state

Cross-section

case 2 → 2 in CM frame



$$\Rightarrow \int d\Phi_2 \equiv (2\pi)^4 \int \delta^4(p_1 + p_2 - p_3 - p_4) \frac{d^3 p_3}{(2\pi)^3 2E_3} \frac{d^3 p_4}{(2\pi)^3 2E_4} = \int \frac{|\mathbf{p}| d\Omega}{16\pi^2 E_{CM}}$$


$$\text{and if } m_1 = m_2 \Rightarrow 4 \{(p_1 p_2)^2 - m_1^2 m_2^2\}^{1/2} = 4E_{CM}|\mathbf{q}|$$

$$\boxed{\frac{d\sigma}{d\Omega}(1, 2 \rightarrow 3, 4) = \frac{1}{64\pi^2 E_{CM}^2} \frac{|\mathbf{p}|}{|\mathbf{q}|} |\mathcal{M}|^2}$$

Decay width

$$d\Gamma(i \rightarrow f) = \frac{1}{2M} |\mathcal{M}|^2 (2\pi)^4 \delta^4(P - p_f) \prod_{j=1}^n \frac{d^3 p_j}{(2\pi)^3 2E_j}$$

case 1 → 2



$$\boxed{\frac{d\Gamma}{d\Omega}(i \rightarrow 1, 2) = \frac{1}{32\pi^2} \frac{|\mathbf{p}|}{M^2} |\mathcal{M}|^2}$$

▷ Note that masses M , m_1 and m_2 fix final energies and momenta:


$$E_1 = \frac{M^2 - m_2^2 + m_1^2}{2M} \quad E_2 = \frac{M^2 - m_1^2 + m_2^2}{2M}$$

$$|\mathbf{p}| = |\mathbf{p}_1| = |\mathbf{p}_2| = \frac{\{[M^2 - (m_1 + m_2)^2][M^2 - (m_1 - m_2)^2]\}^{1/2}}{2M}$$

Loop calculations

Structure of one-loop amplitudes

- Consider the following generic one-loop diagram with N external legs:



$$k_1 = p_1, \quad k_2 = p_1 + p_2, \quad \dots \quad k_{N-1} = \sum_{i=1}^{N-1} p_i$$

- It contains general integrals of the kind:

$$\frac{i}{16\pi^2} T_{\mu_1 \dots \mu_P}^N \equiv \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{q_{\mu_1} \cdots q_{\mu_P}}{[q^2 - m_0^2][(q+k_1)^2 - m_1^2] \cdots [(q+k_{N-1})^2 - m_{N-1}^2]}$$

Structure of one-loop amplitudes

- ▷ D dimensional integration in **dimensional regularization**
- ▷ Integrals are symmetric under permutations of Lorentz indices
- ▷ Scale μ introduced to keep the proper mass dimensions
- ▷ P is the number of q 's in the numerator and determines the tensor structure of the integral (scalar if $P = 0$, vector if $P = 1$, etc.). Note that $P \leq N$
- ▷ Notation: A for T^1 , B for T^2 , etc. For example, the **scalar integrals** A_0, B_0 , etc.
- ▷ The **tensor integrals** can be decomposed as a linear combination of the Lorentz covariant tensors that can be built with $g_{\mu\nu}$ and a set of linearly independent momenta

[Passarino, Veltman '79]

- ▷ The **choice of basis** is not unique

Here we use the basis formed by $g_{\mu\nu}$ and the momenta k_i , where the the **tensor coefficients are totally symmetric in their indices**

[Denner '93]

This is the basis used by the computer package LoopTools

[www.feynarts.de/looptools]

Structure of one-loop amplitudes

- We focus here on:

$$\begin{aligned}B_\mu &= k_{1\mu} B_1 \\B_{\mu\nu} &= g_{\mu\nu} B_{00} + k_{1\mu} k_{1\nu} B_{11} \\C_\mu &= k_{1\mu} C_1 + k_{2\mu} C_2 \\C_{\mu\nu} &= g_{\mu\nu} C_{00} + \sum_{i,j=1}^2 k_{i\mu} k_{j\nu} C_{ij} \\C_{\mu\nu\rho} &= \dots\end{aligned}$$

- We will see that the scalar integrals A_0 and B_0 and the tensor integral coefficients B_1, B_{00}, B_{11} and C_{00} are divergent in $D = 4$ dimensions (ultraviolet divergence, equivalent to take cutoff $\Lambda \rightarrow \infty$ in q)
- It is possible to express every tensor coefficient in terms of scalar integrals (scalar reduction)

[Denner '93]

Explicit calculation

- Basic ingredients:
 - Euler Gamma function:

$$\Gamma(x+1) = x\Gamma(x)$$

Taylor expansion around poles at $x = 0, -1, -2, \dots$:

$$x = 0 : \quad \Gamma(x) = \frac{1}{x} - \gamma + \mathcal{O}(x)$$
$$x = -1 : \quad \Gamma(x) = -\frac{1}{(x+1)} + \gamma - 1 + \dots + \mathcal{O}(x+1)$$

where $\gamma \approx 0.5772\dots$ is Euler-Mascheroni constant

- Feynman parameters:

$$\frac{1}{a_1 a_2 \cdots a_n} = \int_0^1 dx_1 \cdots dx_n \delta \left(\sum_{i=1}^n x_i - 1 \right) \frac{(n-1)!}{[x_1 a_1 + x_2 a_2 + \cdots + x_n a_n]^n}$$

Explicit calculation


- The following integrals, with $\epsilon \rightarrow 0^+$, will be needed:

$$\begin{aligned} \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - \Delta + i0)^n} &= \frac{(-1)^n i}{(4\pi)^{D/2}} \frac{\Gamma(n - D/2)}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n-D/2} \\ \Rightarrow \int \frac{d^D q}{(2\pi)^D} \frac{q^2}{(q^2 - \Delta + i0)^n} &= \frac{(-1)^{n-1} i}{(4\pi)^{D/2}} \frac{D}{2} \frac{\Gamma(n - D/2 - 1)}{\Gamma(n)} \left(\frac{1}{\Delta}\right)^{n-D/2-1} \end{aligned}$$

▷ Let's solve the first integral in Euclidean space: $q^0 = iq_E^0$, $\mathbf{q} = \mathbf{q}_E$, $q^2 = -q_E^2$,

$$\int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - \Delta + i0)^n} = i(-1)^n \int \frac{d^D q_E}{(2\pi)^D} \frac{1}{(q_E^2 + \Delta)^n}$$

(equivalent to a **Wick rotation** of 90°). The second integral follows from this one



Explicit calculation

In D -dimensional spherical coordinates:

$$\int \frac{d^D q_E}{(2\pi)^D} \frac{1}{(q_E^2 + \Delta)^n} = \int d\Omega_D \int_0^\infty dq_E q_E^{D-1} \frac{1}{(q_E^2 + \Delta)^n} \equiv \mathcal{I}_A \times \mathcal{I}_B$$

where

$$\mathcal{I}_A = \int d\Omega_D = \frac{2\pi^{D/2}}{\Gamma(D/2)}$$

$$\begin{aligned} \text{since } (\sqrt{\pi})^D &= \left(\int_{-\infty}^\infty dx e^{-x^2} \right)^D = \int d^D x e^{-\sum_{i=1}^D x_i^2} = \int d\Omega_D \int_0^\infty dx x^{D-1} e^{-x^2} \\ &= \left(\int d\Omega_D \right) \frac{1}{2} \int_0^\infty dt t^{D/2-1} e^{-t} = \left(\int d\Omega_D \right) \frac{1}{2} \Gamma(D/2) \end{aligned}$$


and, changing variables: $t = q_E^2$, $z = \Delta/(t + \Delta)$, we have

$$\mathcal{I}_B = \frac{1}{2} \left(\frac{1}{\Delta} \right)^{n-D/2} \int_0^1 dz z^{n-D/2-1} (1-z)^{D/2-1} = \frac{1}{2} \left(\frac{1}{\Delta} \right)^{n-D/2} \frac{\Gamma(n-D/2)\Gamma(D/2)}{\Gamma(n)}$$

$$\text{where Euler Beta function was used: } B(\alpha, \beta) = \int_0^1 dz z^{\alpha-1} (1-z)^{\beta-1} = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

Explicit calculation

Two-point functions



$$\frac{i}{16\pi^2} \{B_0, B^\mu, B^{\mu\nu}\}(\text{args}) = \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q^\mu, q^\mu q^\nu\}}{(q^2 - m_0^2) [(q+p)^2 - m_1^2]}$$

- ▷ $k_1 = p$
- ▷ The integrals depend on the masses m_0, m_1 and the invariant p^2 :

$$(\text{args}) = (p^2; m_0^2, m_1^2)$$

Explicit calculation

Two-point functions

- Using Feynman parameters,

$$\frac{1}{a_1 a_2} = \int_0^1 dx \frac{1}{[a_1 x + a_2(1-x)]^2}$$

$$\Rightarrow \frac{i}{16\pi^2} \{B_0, B^\mu, B^{\mu\nu}\} = \mu^{4-D} \int_0^1 dx \int \frac{d^D q}{(2\pi)^D} \frac{\{1, -A^\mu, q^\mu q^\nu + A^\mu A^\nu\}}{(q^2 - \Delta_2)^2}$$

with

$$\Delta_2 = x^2 p^2 + x(m_1^2 - m_0^2 - p^2) + m_0^2$$

$$\begin{aligned} a_1 &= (q + p)^2 - m_1^2 \\ a_2 &= q^2 - m_0^2 \end{aligned}$$

and a loop momentum shift to obtain a perfect square in the denominator:

$$q^\mu \rightarrow q^\mu - A^\mu, \quad A^\mu = x p^\mu$$

Explicit calculation

Two-point functions

- Then, the scalar function is:

$$\begin{aligned}\frac{i}{16\pi^2}B_0 &= \mu^{4-D} \int_0^1 dx \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - \Delta_2)^2} \\ \Rightarrow B_0 &= \Delta_\epsilon - \int_0^1 dx \ln \frac{\Delta_2}{\mu^2} + \mathcal{O}(\epsilon) \quad [D = 4 - \epsilon]\end{aligned}$$

where $\Delta_\epsilon \equiv \frac{2}{\epsilon} - \gamma + \ln 4\pi$ and the Euler Gamma function was expanded around $x = 0$ for $D = 4 - \epsilon$, using $x^\epsilon = \exp\{\epsilon \ln x\} = 1 + \epsilon \ln x + \mathcal{O}(\epsilon^2)$:

$$\mu^{4-D} \frac{i\Gamma(2 - D/2)}{(4\pi)^{D/2}} \left(\frac{1}{\Delta_2}\right)^{2-D/2} = \frac{i}{16\pi^2} \left(\Delta_\epsilon - \ln \frac{\Delta_2}{\mu^2}\right) + \mathcal{O}(\epsilon)$$

- Comparing with the definitions of the tensor coefficients we have:

$$\begin{aligned}\frac{i}{16\pi^2}B^\mu &= -\mu^{4-D} \int_0^1 dx \int \frac{d^D q}{(2\pi)^D} \frac{A^\mu}{(q^2 - \Delta_2)^2} \\ \Rightarrow B_1 &= -\frac{1}{2}\Delta_\epsilon + \int_0^1 dx x \ln \frac{\Delta_2}{\mu^2} + \mathcal{O}(\epsilon) \quad [D = 4 - \epsilon]\end{aligned}$$

Explicit calculation

Two-point functions

and


$$\begin{aligned} \frac{i}{16\pi^2} B^{\mu\nu} &= \mu^{4-D} \int_0^1 dx \int \frac{d^D q}{(2\pi)^D} \frac{(q^2/D)g^{\mu\nu} + A^\mu A^\nu}{(q^2 - \Delta_2)^2} \\ \Rightarrow B_{00} &= -\frac{1}{12}(p^2 - 3m_0^2 - 3m_1^2)\Delta_\epsilon + \mathcal{O}(\epsilon) \quad [D = 4 - \epsilon] \\ B_{11} &= \frac{1}{3}\Delta_\epsilon - \int_0^1 dx x^2 \ln \frac{\Delta_2}{\mu^2} + \mathcal{O}(\epsilon) \quad [D = 4 - \epsilon] \end{aligned}$$

where $q^\mu q^\nu$ have been replaced by $(q^2/D)g^{\mu\nu}$ in the integrand and the Euler Gamma function was expanded around $x = -1$ for $D = 4 - \epsilon$:

$$-\mu^{4-D} \frac{i\Gamma(1 - D/2)}{(4\pi)^{D/2} 2\Gamma(2)} \left(\frac{1}{\Delta_2}\right)^{1-D/2} = \frac{i}{16\pi^2} \frac{1}{2} \left(\Delta_\epsilon - \ln \frac{\Delta_2}{\mu^2} + 1\right) \Delta_2 + \mathcal{O}(\epsilon)$$

Explicit calculation

Three-point functions



$$\frac{i}{16\pi^2} \{C_0, C^\mu, C^{\mu\nu}\} (\text{args}) = \mu^{4-D} \int \frac{d^D q}{(2\pi)^D} \frac{\{1, q^\mu, q^\mu q^\nu\}}{(q^2 - m_0^2) [(q + p_1)^2 - m_1^2] [(q + p_2)^2 - m_2^2]}$$

▷ It is convenient to choose the external momenta so that:

$$k_1 = p_1, \quad k_2 = p_2.$$

▷ The integrals depend on the masses m_0 , m_1 , m_2 and the invariants:

$$(\text{args}) = (p_1^2, Q^2, p_2^2; m_0^2, m_1^2, m_2^2), \quad Q^2 \equiv (p_2 - p_1)^2.$$

Explicit calculation

Three-point functions

- Using Feynman parameters,

$$\frac{1}{a_1 a_2 a_3} = 2 \int_0^1 dx \int_0^{1-x} dy \frac{1}{[a_1 x + a_2 y + a_3 (1 - x - y)]^3}$$

$$\Rightarrow \frac{i}{16\pi^2} \{C_0, C^\mu, C^{\mu\nu}\} = 2\mu^{4-D} \int_0^1 dx \int_0^{1-x} dy \int \frac{d^D q}{(2\pi)^D} \frac{\{1, -A^\mu, q^\mu q^\nu + A^\mu A^\nu\}}{(q^2 - \Delta_3)^3}$$

with

$$\Delta_3 = x^2 p_1^2 + y^2 p_2^2 + xy(p_1^2 + p_2^2 - Q^2) + x(m_1^2 - m_0^2 - p_1^2) + y(m_2^2 - m_0^2 - p_2^2) + m_0^2$$

$$\begin{aligned} a_1 &= (q + p_1)^2 - m_1^2 \\ a_2 &= (q + p_2)^2 - m_2^2 \\ a_3 &= q^2 - m_0^2 \end{aligned}$$

and a loop momentum shift to obtain a perfect square in the denominator:

$$q^\mu \rightarrow q^\mu - A^\mu, \quad A^\mu = x p_1^\mu + y p_2^\mu$$

Explicit calculation

Three-point functions

- Then the scalar function is:

$$\begin{aligned}\frac{i}{16\pi^2}C_0 &= 2\mu^{4-D} \int_0^1 dx \int_0^{1-x} dy \int \frac{d^D q}{(2\pi)^D} \frac{1}{(q^2 - \Delta_3)^3} \\ \Rightarrow C_0 &= - \int_0^1 dx \int_0^{1-x} dy \frac{1}{\Delta_3} \quad [D = 4]\end{aligned}$$

- Comparing with the definitions of the tensor coefficients we have:

$$\begin{aligned}\frac{i}{16\pi^2}C^\mu &= -2\mu^{4-D} \int_0^1 dx \int_0^{1-x} dy \int \frac{d^D q}{(2\pi)^D} \frac{A^\mu}{(q^2 - \Delta_3)^3} \\ \Rightarrow C_1 &= \int_0^1 dx \int_0^{1-x} dy \frac{x}{\Delta_3} \quad [D = 4] \\ C_2 &= \int_0^1 dx \int_0^{1-x} dy \frac{y}{\Delta_3} \quad [D = 4]\end{aligned}$$

Explicit calculation

Three-point functions

$$\begin{aligned}
\frac{i}{16\pi^2} C^{\mu\nu} &= 2\mu^{4-D} \int_0^1 dx \int_0^{1-x} dy \int \frac{d^D q}{(2\pi)^D} \frac{(q^2/D)g^{\mu\nu} + A^\mu A^\nu}{(q^2 - \Delta_3)^3} \\
\Rightarrow C_{11} &= - \int_0^1 dx \int_0^{1-x} dy \frac{x^2}{\Delta_3} \quad [D = 4] \\
C_{22} &= - \int_0^1 dx \int_0^{1-x} dy \frac{y^2}{\Delta_3} \quad [D = 4] \\
C_{12} &= - \int_0^1 dx \int_0^{1-x} dy \frac{xy}{\Delta_3} \quad [D = 4] \\
C_{00} &= \frac{1}{4}\Delta_\epsilon - \frac{1}{2} \int_0^1 dx \int_0^{1-x} dy \ln \frac{\Delta_3}{\mu^2} + \mathcal{O}(\epsilon) \quad [D = 4 - \epsilon]
\end{aligned}$$

where $\Delta_\epsilon \equiv \frac{2}{\epsilon} - \gamma + \ln 4\pi$ and $q^\mu q^\nu$ was replaced by $(q^2/D)g^{\mu\nu}$ in the integrand

In C_{00} the Euler Gamma function was expanded around $x = 0$ for $D = 4 - \epsilon$:

$$\mu^{4-D} \frac{i\Gamma(2 - D/2)}{(4\pi)^{D/2}\Gamma(3)} \left(\frac{1}{\Delta_3}\right)^{2-D/2} = \frac{i}{16\pi^2} \frac{1}{2} \left(\Delta_\epsilon - \ln \frac{\Delta_3}{\mu^2}\right) + \mathcal{O}(\epsilon)$$

Note about Diracology in D dimensions

- Attention should be paid to the traces of Dirac matrices when working in D dimensions (dimensional regularization) since

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu} \mathbf{1}_{4 \times 4}, \quad g^{\mu\nu} g_{\mu\nu} = \text{Tr}\{g^{\mu\nu}\} = D$$

Thus, the following identities involving contractions of Lorentz indices can be proven:

$$\begin{aligned}\gamma^\mu \gamma_\mu &= D \\ \gamma^\mu \gamma^\nu \gamma_\mu &= -(D - 2)\gamma^\nu \\ \gamma^\mu \gamma^\nu \gamma^\rho \gamma_\mu &= 4g^{\nu\rho} - (4 - D)\gamma^\nu \gamma^\rho \\ \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma \gamma_\mu &= -2\gamma^\sigma \gamma^\rho \gamma^\nu + (4 - D)\gamma^\nu \gamma^\rho \gamma^\sigma\end{aligned}$$