

ELECTROWEAK STRUCTURE OF THE NUCLEON

Fernando Alvarado (falvar@ific.uv.es) Luis Alvarez-Ruso Di An Stefan Leupold

January 15, 2024

Exited QCD 2024, Benasque



VNIVERSITAT
ID VALÈNCIA



ELECTROWEAK STRUCTURE OF THE NUCLEON

- Structure of protons and neutrons is encoded in form factors

F_{EM} → charge distribution

- $V^\mu = \bar{q}Q\gamma^\mu q$,

$$\langle N|V^\mu(0)|N\rangle = \bar{u}' \left[\gamma^\mu F_1(q^2) + \frac{i\sigma^{\mu\nu}q_\nu}{2m_N} F_2(q^2) \right] u$$

- $G_E = F_1 - \frac{Q^2}{4m_N^2} F_2$, $G_M = F_1 + F_2$

- slope=charge radius: $\langle r_E^2 \rangle = 6 \left. \frac{dG_E}{dq^2} \right|_0$



- $A^\mu = \bar{q}\gamma^\mu\gamma_5 q$,

$$\langle N|A^\mu|N\rangle = \bar{u}' \left\{ \gamma_\mu F_A(q^2) + \frac{q_\mu}{2m_N} G_P(q^2) \right\} \gamma_5 u$$


- F_A → spin distribution

Weak interaction is $V - A$

relevant for ν experiments



ELECTROMAGNETIC FORM FACTOR



- Experimental determinations

- $\langle r_E^2 \rangle$: proton radius puzzle [Pohl et al., Nature 466, 213 (2010)]

- Lattice QCD


- Nucleon f.f. is a benchmark for LQCD
- Uncertainties reduced for unphysical large M_π
- Technical difficulties \rightarrow recent progress
- Experimental and lattice q^2 parametrisation:

- dipole ansatz
- z-expansion
- ...

} \Rightarrow different $\langle r_1^2 \rangle$, and F_{EM} in general

- Theoretical input needed

ELECTROMAGNETIC FORM FACTOR



- **Experimental determinations**

- $\langle r_E^2 \rangle$: proton radius puzzle [Pohl et al., Nature 466, 213 (2010)]

- **Lattice QCD**


- Nucleon f.f. is a benchmark for LQCD
- Uncertainties reduced for unphysical large M_π
- Technical difficulties \rightarrow recent progress
- Experimental and lattice q^2 parametrisation:

- dipole ansatz
- z-expansion
- ...

} \Rightarrow different $\langle r_1^2 \rangle$, and F_{EM} in general

- **Theoretical input needed**

- **Chiral Perturbation Theory** (χ PT) \rightarrow parametrise M_π and q^2 dep.
- **Dispersion theory** \rightarrow enlarge q^2 range
- **Goal**: Disp+ χ PT = good q^2 and M_π description



- Lattice QCD **parametrisation issue**
- Chiral Perturbation Theory (χ PT)
 - QCD based parametrisation of q^2 and M_π dependencies \implies extrapolate lattice results to the phys. point and extract $\langle r_1^2 \rangle$ and κ from the lattice simulations
 - Relativistic baryon χ PT (EOMS)
 - Terms that break power counting are absorbed in LECs

DISPERSION THEORY

- Enlarge the q^2 range of χ PT (ρ dynamics)

1. Analyticity \Rightarrow Disp. rel. (Cauchy)

$$F(q^2) = \int_{4M_\pi^2}^{\infty} \frac{ds}{\pi} \frac{\text{Im} F(s)}{s - q^2 - i\epsilon}$$



2. Unitarity \Rightarrow $\text{Im} F = \frac{1}{2} \sum_n T \gamma^{*n} T_{n\bar{N}N}^\dagger$, $n = \pi^+ \pi^-$, ...

- $\ell = 1$, $\pi\pi$ must be iso-vector (ρ channel)

$$\Rightarrow F_i = \frac{1}{2} F_i^{(s)} + F_i^{(v)} \frac{\sigma^3}{2}, \quad F_i^{(v)} = F_i^p - F_i^n$$

3. Using full $N\bar{N}\pi\pi$ and $\gamma^*\pi\pi$ vertices with M_π dep.

$$F(q^2) = \frac{1}{12\pi} \int_{4M_\pi^2}^{\infty} \frac{ds}{\pi} \frac{T p_{\text{cm}}^3 F_\pi^{V*}}{s^{1/2}(s - q^2 - i\epsilon)},$$



DISPERSION THEORY

- Our two vertices, T and F_{π}^V , include nonperturbatively the $\pi\pi$ scattering amplitude, t , thanks to IAM (resummation)

$$\mathcal{A}(s) = \frac{8}{\pi} \sum_{\ell} (2\ell + 1) P_{\ell}(\cos \theta) t_{\ell}(s)$$



$$\ell = 1, t = \frac{\sqrt{s}}{2p_{\text{cm}}} \sin \delta e^{i\delta}$$

$$\Omega(s) = \exp \left\{ s \int_{4M_{\pi}^2}^{\infty} \frac{ds'}{\pi} \frac{\delta(s')}{s'(s'-s-i\epsilon)} \right\}$$

- We fit NLO t_{IAM} to physical δ from [Garcia-Martin PRD 83(2011)]

- We check that the M_{π} dependence is realistic

$$F_{\pi}^V(s) = [1 + \alpha_V s] \Omega(s)$$



DISPERSION THEORY

- Our two vertices, T and F_{π}^V , include nonperturbatively the $\pi\pi$ scattering amplitude, t , thanks to IAM (resummation)

- $\mathcal{A}(s) = \frac{8}{\pi} \sum_{\ell} (2\ell + 1) P_{\ell}(\cos \theta) t_{\ell}(s)$



- $\ell = 1, t = \frac{\sqrt{s}}{2p_{\text{cm}}} \sin \delta e^{i\delta}$

- $\Omega(s) = \exp \left\{ s \int_{4M_{\pi}^2}^{\infty} \frac{ds'}{\pi} \frac{\delta(s')}{s'(s'-s-i\epsilon)} \right\}$

- We fit NLO t_{IAM} to physical δ from [Garcia-Martin PRD 83(2011)]

- We check that the M_{π} dependence is realistic

- $F_{\pi}^V(s) = [1 + \alpha_V s] \Omega(s)$



$\rho \rightarrow \pi\pi, \Gamma_{\rho} \rightarrow 0$

DISPERSION THEORY

- Our two vertices, T and F_π^V


- $F_\pi^V(s) = [1 + \alpha_V s] \Omega(s)$

- $T_i(s) = K_i(s) + \Omega(s) P_i + I_i(s)$

- $I_i(s) = \Omega(s) \int_{4M_\pi^2}^{\infty} \frac{ds'}{\pi} \frac{K_i(s') \sin \delta(s')}{|\Omega(s')|(s'-s-i\epsilon)}$

- K and P from $N\bar{N} \rightarrow \pi\pi$ in χ PT
 $\mathcal{M} = A\bar{v}u - \frac{1}{2}B\bar{v}\not{k}u \longleftrightarrow K, P$


$$\Rightarrow F(q^2) = \frac{1}{12\pi} \int_{4M_\pi^2}^{\infty} \frac{ds}{\pi} \frac{T p_{\text{cm}}^3 F_\pi^{V*}}{s^{1/2}(s - q^2 - i\epsilon)},$$








· $F_{EM} = F_{EM}^{disp} + \text{diagrams w/o } 2\pi \text{ cut from } \chi PT$

· $F(q^2) = \int \frac{ds}{4M_\pi^2} \frac{d s}{\pi} \frac{\text{Im} F_{2\pi}(s)}{s - q^2} + F_{\chi PT \text{ no } 2\pi}$

· F_{EM}^{disp}




· $+\chi PT$ diagrams



- $F_{EM} = F_{EM}^{disp} + \text{diagrams w/o } 2\pi \text{ cut from } \chi PT$


- $F(q^2) = \int \frac{ds}{4M_\pi^2} \frac{\text{Im}F_{2\pi}(s)}{s-q^2} + F_{\chi PT \text{ no } 2\pi}$

- F_{EM}^{disp}



- $+\chi PT$ diagrams


- Relativistic and with explicit $\Delta(1232)$ [Bauer et al., PRC 86 (2012)]
- green: $F_1 - F_1(0)$ (the charge is trivial)
- blue: $F_2 \rightarrow$ we add the $\mathcal{O}(p^4)$ Δ terms



- $F_{EM} = F_{EM}^{disp} + \text{diagrams w/o } 2\pi \text{ cut from } \chi PT$

- $F(q^2) = \int \frac{ds}{4M_\pi^2} \frac{\pi}{\pi} \frac{\text{Im}F_{2\pi}(s)}{s-q^2} + F_{\chi PT \text{ no } 2\pi}$

- F_{EM}^{disp}




- + χPT diagrams

- Relativistic and with explicit $\Delta(1232)$ [Bauer et al., PRC 86 (2012)]
- green: $F_1 - F_1(0)$ (the charge is trivial)
- blue: $F_2 \rightarrow$ we add the $\mathcal{O}(p^4)$ Δ terms


- Disp and χPT differ in the renormalization (UV)

- At $\mathcal{O}(p^3)$ disp and χPT agree on the M_π nonanalyticities
- Example: the dispersive contribution from $T^{\text{point}} \sim \frac{1}{F^2}$ agrees with χPT
 $F_1^{\text{point}} \sim F_1^{(9)} \sim q^2 \log M_\pi$
- differences absorbed in LECs




· Dirac f.f., F_1


- $\langle N|V^\mu|N\rangle = \bar{u}' \left[\gamma^\mu F_1 + \frac{i\sigma^{\mu\nu}q_\nu}{2m_N} F_2 \right] u$
- $F_1 = 1 + \frac{q^2}{6} \left[-12d_6 + \langle r_1^{2(\log M_\pi)} \rangle \log M_\pi \right] + \mathcal{O}(p^4)$
- Comparison with LQCD data [Djukanovic PRD 103(2021)] ← controlled FV and discret. effects
- In the χ PT and disp+ χ PT F_1 , d_6 is fitted to LQCD
- Disp \rightarrow q^2 curvature




· Dirac f.f., F_1

- $\langle N|V^\mu|N\rangle = \bar{u}' \left[\gamma^\mu F_1 + \frac{i\sigma^{\mu\nu}q_\nu}{2m_N} F_2 \right] u$
- $F_1 = 1 + \frac{q^2}{6} \left[-12d_6 + \langle r_1^{2(\log M_\pi)} \rangle \log M_\pi \right] + \mathcal{O}(p^4)$
- Comparison with LQCD data [Djukanovic PRD 103(2021)] ← controlled FV and discret. effects
- In the χ PT and disp+ χ PT F_1 , d_6 is fitted to LQCD
- Disp $\rightarrow q^2$ curvature





(h) H105 $M_\pi = 0.278$ GeV




(i) N302 $M_\pi = 0.353$ GeV

- Disp+ χ PT describes well the M_π dep.

- good d_6 fit for $Q^2 < 0.6$ GeV² and $M_\pi \lesssim 350$ MeV
- outperforms the pure dispersive and plain χ PT descriptions

	Disp (prediction)	χ PT	Disp+ χ PT
$d_6(\mu = m_\rho)$ (GeV ⁻²)	-	0.074 ± 0.010	0.416 ± 0.010
$d_6(\mu = m_N)$ (GeV ⁻²)	-	-0.422 ± 0.010	0.155 ± 0.010
χ^2/dof	$108.9/47 = 2.32$	$73.9/(47 - 1) = 1.61$	$24.6/(47 - 1) = 0.53$
$\langle r_1^2 \rangle_{\text{phys}}$ (fm ²)	0.4541	0.3626 ± 0.0047	0.4838 ± 0.0047



- $F_1^{(v)} = 1 + \frac{1}{6} \langle r_1^2 \rangle^{(v)} q^2 + \mathcal{O}(q^4)$,
 $\langle r_1^2 \rangle^{\text{PDG}} = 0.577 \text{ fm}^2$
- Heavy baryon fit to LQCD from [Djukanovic PRD 103(2021)]:
 $\langle r_1^2 \rangle^{\text{HB}} = 0.554 \pm 0.035 \text{ fm}^2$

	Disp (prediction)	χ^2_{PT}	Disp+ χ^2_{PT}
$d_6(\mu = m_\rho)$ (GeV ⁻²)	-	0.074 ± 0.010	0.416 ± 0.010
$d_6(\mu = m_N)$ (GeV ⁻²)	-	-0.422 ± 0.010	0.155 ± 0.010
χ^2/dof	$108.9/47 = 2.32$	$73.9/(47 - 1) = 1.61$	$24.6/(47 - 1) = 0.53$
$\langle r_1^2 \rangle_{\text{phys}}$ (fm ²)	0.4541	0.3626 ± 0.0047	0.4838 ± 0.0047


• Pauli f.f., F_2

• $\langle N | V^\mu | N \rangle = \bar{u}' \left[\gamma^\mu F_1 + \frac{i\sigma^{\mu\nu} q_\nu}{2m_N} F_2 \right] u$, $F_2 = \frac{F_M - F_E}{(1 + Q^2/(4m^2))}$

• $F_2^{(\nu)} = \kappa^{(\nu)} \left[1 + \frac{1}{6} \langle r_2^2 \rangle^{(\nu)} q^2 + \mathcal{O}(q^4) \right]$

• $\mathcal{O}(p^3)$ χ PT is not enough (and Δ subtleties)


$F_2(0)^{\mathcal{O}(3)} = c_6 - \left(\frac{\pi m_N g_A^2}{4\pi^2 F^2} \right) M_\pi$



• \implies include $\Delta \mathcal{O}(p^4)$


• disp+ χ PT $\mathcal{O}(p^4)$: $F_2 = F_2^{\text{disp}} + F_2^{\text{tree}} + F_2^{\chi\text{PTloop}}$,

$F_2^{\text{tree}} = c_6 - 16e_{106} m_N M_\pi^2 + 2q^2(d_6 + 2e_{74} m_N)$




· F_2 fit to LQCD

- In F_2^{disp} free c_6
- In $F_2^{\chi\text{PT}}$ and $F_2^{\text{disp}+\chi\text{PT}}$, free c_6, e_{106}, e_{74}
- χPT $\mathcal{O}(p^4)$ and disp separately are good enough to describe the data




(a) H105 $M_\pi = 0.278$ GeV




(b) N302 $m_\pi = 0.353$ GeV

· F_2 fit to LQCD



- In F_2^{disp} free c_6
- In $F_2^{\chi\text{PT}}$ and $F_2^{\text{disp}+\chi\text{PT}}$, free c_6, e_{106}, e_{74}
- χPT $\mathcal{O}(p^4)$ and disp separately are good enough to describe the data



(a) H105 $M_\pi = 0.278$ GeV



(b) N302 $m_\pi = 0.353$ GeV




- $F_2^{(\nu)} = \kappa^{(\nu)} \left[1 + \frac{1}{6} \langle r_2^2 \rangle^{(\nu)} q^2 + \mathcal{O}(q^4) \right]$
- $\kappa_{\text{PDG}} = 3.706,$
 $\kappa_{\text{HB}} = 3.71 \pm 0.17,$
 $\langle r_2^2 \rangle_{\text{PDG}} = 0.7754 \text{ fm}^2,$
 $\langle r_2^2 \rangle_{\text{HB}} = 0.690 \pm 0.042 \text{ fm}^2.$

	Disp+c6	χ^{PT}	Disp+ χ^{PT}
χ^2/dof	$\frac{49.95}{47-2} = 1.11$	$\frac{44.18}{47-4} = 1.027$	$\frac{56.08}{47-4} = 1.304$
χ_0^2/dof	1.09	1.027	1.283
κ_{phys}	3.64	3.42	3.61
$\langle r_2^2 \rangle_{\text{phys}} (\text{fm}^2)$	0.673	0.619	0.668

- In general **good $F_{1,2}$ description at different M_π values**
- **Our extraction of $\langle r_{1,2}^2 \rangle$ and κ from LQCD is in line with PDG**
 - In this case HB yields also reasonable results

AXIAL FORM FACTOR

- Analogously to the F_{EM} work:
 1. calculate F_A in ChPT
 2. analyse LQCD data



AXIAL FORM FACTOR

- $F_A \rightarrow$ spin distribution

Weak interaction is $V - A$

$$\cdot A^{i\mu} = \bar{q} \gamma^\mu \gamma_5 \frac{\tau^i}{2} q, \langle N | A^\mu | N \rangle = \bar{u}' \left[\gamma^\mu F_A(q^2) + \frac{q_\mu}{2m_N} G_P(q^2) \right] \gamma_5 \frac{\tau^i}{2} u$$

- Nucleon axial isovector form factor

$$\cdot F_A(q^2) = g_A \left[1 + \frac{1}{6} \langle r_A^2 \rangle q^2 + \mathcal{O}(q^4) \right]$$

- g_A and F_A dependence in q^2 are necessary in ν oscillations experiments

- μ capture, β -decay

- χ PT calculation of F_A

\implies extract $\langle r_A^2 \rangle$ from lattice QCD without ad-hoc parametrization

- $\mathcal{O}(p^4)$ F_A in relativistic χ PT

$$\cdot F_A = \hat{g}_A + 4d_{16}M_\pi^2 + d_{22}q^2 + \text{loops}(M_\pi, q^2)$$




Figure: $\mathcal{O}(p)$ and $\mathcal{O}(p^3)$ (w. f. renormalisation not shown)




Figure: $\mathcal{O}(p^4)$

$g_A(M_\pi)$ study

[FA & Alvarez-Ruso PRD 105 \(2021\)](#)

Motivations:

1. **test** the LECs extracted from πN scattering
2. **determine** d_{16} from fit to lattice:
 - source of uncertainty in m_q dependence of nuclear properties (ground-state and binding energies)
3. **study** more deeply the **convergence** of g_A in χ PT

$g_A \equiv F_A(q^2 = 0)$ ANALYSIS

- $\pi N \rightarrow \pi\pi N \Rightarrow d_{16}, c_i$
 - $\pi N \rightarrow \pi N \Rightarrow c_i$
 - $g_A^{\text{phys}} = g_A(M_\pi^{\text{phys}}) \Rightarrow \hat{g}_A$
- combined fit Refs. [*]

⇒ **Bad Δ -less $g_A(M_\pi)$ prediction**




Table: $\mathcal{O}(p^4)$: **steep rise** (OK with [**])

	$\mathcal{O}(p^3)$	$\mathcal{O}(p^4)$
d_{16}	-2.2 ± 1.1	-1.86 ± 0.80
c_1	-	-0.89 ± 0.06
c_2	-	3.38 ± 0.15
c_3	-	-4.59 ± 0.09
c_4	-	3.31 ± 0.13



- ⇒ We include explicit $\Delta(1232)$ and fit d_{16} to lattice
- ⇒ This reconciles for g_A^{LQCD} with the πN LECs, c_i (with the caveat of a slow convergence)
- we will discuss the whole $F_A(q^2)$ directly

[*] [Siemens et al. PRC 94 \(2016\)](#), [Siemens et al. PRC 96 \(2017\)](#), (value converted to standard EOMS)

[**] [Bernard et al PLD 639 \(2006\)](#)

- F_A : Meta-analysis of large set of recent LQCD results

- Many recent works \Rightarrow substantial improvements
- RQCD^[1] + PNDME^[2] + "Mainz"^[3] + PACS^[4] + ETMC^[5]





[1] [Bali et al. JHEP 05 \(2020\)](#)

[2] [Park et al. 2103.05599](#)

[3] [Meyer et al. Modern Phys. A 34 \(2019\)](#)

[4] [Shintani et al. PRD 102 \(2020\)](#)

[5] [Alexandrou et al. PRD 103 \(2021\)](#)





- F_A fit procedure:

- Difference between orders \simeq theoretical uncertainty [Epelbaum EPJA 53 (2015)]

$$\Delta g_{A\chi}^{(4)} = \max \left\{ \left(\frac{M_\pi}{\Lambda} \right)^4 |\dot{g}_A|, \left(\frac{M_\pi}{\Lambda} \right)^2 |g_A^{(3)}|, \frac{M_\pi}{\Lambda} |g_A^{(4)}| \right\}$$

- $\Delta F_{A\chi}$ is added to LQCD errors in the χ^2
- LECs have naturalness priors
- fit range: χ^2 plateau $\implies M_\pi^{\text{cut}} \simeq 400$ MeV, $Q_{\text{cut}}^2 = 0.36$ GeV²





- F_A fit procedure:

- Difference between orders \simeq theoretical uncertainty [Epelbaum EPJA 53 (2015)]

$$\Delta g_{A\chi}^{(4)} = \max \left\{ \left(\frac{M_\pi}{\Lambda} \right)^4 |\dot{g}_A|, \left(\frac{M_\pi}{\Lambda} \right)^2 |g_A^{(3)}|, \frac{M_\pi}{\Lambda} |g_A^{(4)}| \right\}$$



- $\Delta F_{A\chi}$ is added to LQCD errors in the χ^2
- LECs have naturalness priors
- fit range: χ^2 plateau $\implies M_\pi^{\text{cut}} \simeq 400$ MeV, $Q_{\text{cut}}^2 = 0.36$ GeV²
- **Fit results: good description**
 - accurate description at the physical point
 - Δ baryon is a necessary d.o.f.
 - $\mathcal{O}(p^5)$ still needed for full convergence

$$F_A(q^2 = 0) = \boxed{g_A}$$



- Axial charge results from $F_A(q^2)$ fit
 - $g_A(M_{\pi\text{phys}}) = 1.273 \pm 0.014$ vs $g_A^{\text{exp}} = 1.2754(13)_{\text{exp}(2)\text{RC}}$
 \Rightarrow excellent agreement with exp.
 vs $g_A^{\text{FLAG}} = 1.246 \pm 0.028$
 - $g_A(M_\pi) = \mathring{g}_A + 4d_{16}M_\pi^2 + \text{loop}(M_\pi)$
 - $d_{16} = -1.46 \pm 1.00 \text{ GeV}^{-2}$
- $\rightarrow M_\pi$ dependence of long range nuclear forces
 - Can not be extracted from πN elastic scattering
 - In line with $d_{16} = -1.0 \pm 1.0 \text{ GeV}^{-2}$ from $\pi N \rightarrow \pi\pi N$
 [Siemens et al. PRC 96 (2017)] (EOMS corrected)

$$F_A = g_A \left(1 + \frac{1}{6} \langle r_A^2 \rangle q^2 \right) \quad \text{AXIAL RADIUS}$$



(Bodek)=Bodek, Eur. Phys. J. C 53, 349 (2008)

(Meyer)=Meyer, PRD 93, 113015 (2016)

(Hill)=Hill, Rept. Prog. Phys. 81 (2018)

• Our $\mathcal{O}(p^4)$ χ PT extraction:

- M_π slope driven by loops with Δ
- $d_{22} = 0.29 \pm 1.69 \text{ GeV}^{-2}$ (no assumptions on $\Delta\Delta\pi$ coupling enlarges error)
 - d_{22} compatible with $\mathcal{O}(p^3)$ π electroprod. [Guerrero et al. PRD, 102 \(2020\)](#)

$$\langle r_A^2 \rangle(M_{\text{phys}}) = 0.293 \pm 0.044 \text{ fm}^2$$

- Empirical determinations (model dependent) are in **tension** with ours and with most of LQCD extractions
- Typically the extracted $\langle r_A^2 \rangle^{\text{phys}}$ value varies depending on the parametrisation
- Our QCD-based parametrisation leads to a value in line with most individual LQCD extractions

CONCLUSIONS

- **F_{EM}**
 - **Dirac f.f., F₁**
 - The dispersive calculation supplemented with χ PT contributions outperforms the pure dispersive and plain χ PT descriptions
 - it fits well the LQCD F₁ at least for $Q^2 < 0.6 \text{ GeV}^2$ and $M_\pi \lesssim 350 \text{ MeV}$
 - $\langle r_1^2 \rangle_{\text{phys}} = 0.4838 \pm 0.0047 \text{ fm}^2$
 - value close to the LQCD HB [Djukanovic PRD 103(2021)] extraction and to the experimental one
 - **Pauli f.f., F₂**
 - Disp, $\mathcal{O}(p^4)$ χ PT and disp+ χ PT describe the data well
 - $\kappa_{\text{phys}} = 3.61$ and $\langle r_2^2 \rangle_{\text{phys}} = 0.668 \text{ fm}^2$
 - values in line with the HB and the experimental ones
- **F_A**
 - Successful description of LQCD F_A(q²) using $\mathcal{O}(p^4)$ relativistic χ PT
 - Explicit $\Delta(1232)$ necessary
 - Fit describes data in $M_\pi^{\text{cut}} \simeq 400 \text{ MeV}$, $Q_{\text{cut}}^2 = 0.36 \text{ GeV}^2$
 - There is tension between the experimental and lattice extraction of $\langle r_A^2 \rangle$
 - We extract $\langle r_A^2 \rangle^{\text{phys}} = 0.291 \pm 0.052 \text{ fm}^2$ without ad hoc parametrisations
- Useful LEC values extracted from both calculations (agreement with other analysis)

· Summary of dispersive nucleon F_{EM}

$$1. F(q^2) \approx \int_{4M_\pi^2}^{\Lambda^2} \frac{ds}{\pi} \frac{\text{Im}F_{2\pi}(s)}{s-q^2-i\epsilon} + F_{\text{ChPT no } 2\pi}$$

$$2. \text{Im}F_{2\pi}(s) = F_\pi^*(s) \frac{p_{\text{cm}}(s)^3}{12\pi\sqrt{s}} T(s), \text{ need } F_\pi(s) \text{ and } T(s) \leftrightarrow \text{given by } \pi\pi \text{ scattering}$$

3. We use t_{IAM} NLO with a Blatt-Weiskopf ff on t_2 to go to π smoothly: $\tilde{t}_2 = t_2/(1+r^2 p_{\text{cm}}^2)$.

- We fit $l_2 - 2l_1$ and r to $\delta(s)$ of [García-Martín] at M_π^{phys}

- Describe well the $m_\rho(M_\pi)$ from LQCD

$$4. F_\pi^V(s, M_\pi) = (1 + \alpha_V(M_\pi))\Omega(s, M_\pi), \alpha_V \text{ determined from } dF_\pi^V(s)/ds \text{ in ChPT and } \Omega \rightarrow \text{improves } F_\pi^V \text{ } \rho \text{ peak}$$

5. T has to respect Watson th. ($\text{Im} F_{2\pi} \in \mathbb{R}$), therefore Omnès solution

$$T(s) = K(s) + \Omega(s)P + \Omega(s) \int_{4M_\pi^2}^{\Lambda^2} \frac{ds'}{\pi} \frac{\sin \delta_1(s') K(s')}{|\Omega(s')|(s'-s-i\epsilon)}$$

- P and K determined from ChPT πN p-wave projected

6. Subtractions on $T_{1,2}$ and $F_{1,2}$ if M_π -indep

- T_1 subtracted, $P_1 = P_1^{LO}$ enough for leading 1-loop F_1 . (F_1 also trivially subtracted)

- T_2 and F_2 unsubtracted due to M_π dependences (but still the Λ dep. mild)

- EM f.f.
 - $V^\mu = \bar{q}Q\gamma^\mu q$,
 $\langle N(p')|V^\mu(0)|N(p)\rangle = \bar{u}' \left[\gamma^\mu F_1(q^2) + \frac{i\sigma^{\mu\nu}q_\nu}{2m_N} F_2(q^2) \right] u$
 - $G_E^N = F_1^N - \frac{Q^2}{4m_N^2} F_2^N$, $G_M^N = F_1^N + F_2^N$
 - $F_1 = 1 \times \left[1 + \frac{1}{6} \langle r_1^2 \rangle q^2 + \mathcal{O}(q^4) \right]$
 $F_1(0) = F_E(0) = 1$ electric charge
 - $F_2 = \kappa \left[1 + \frac{1}{6} \langle r_2^2 \rangle q^2 + \mathcal{O}(q^4) \right]$
 $\kappa = F_M(0) = \mu - 1$ anom. magn. mom.
 - $\langle r_E^2 \rangle = \langle r_1^2 \rangle + \frac{3}{2m_N^2} \kappa$ proton radius puzzle
 - $F_i = \frac{1}{2} F_i^{(s)} + F_i^{(v)} \frac{\sigma^3}{2}$
 - $F_1^{(v)} = 1 \times \left[1 + \frac{1}{6} \langle r_1^2 \rangle^{(v)} q^2 + \mathcal{O}(q^4) \right]$
 $F_1^{(v)}(0) = F_E^{(v)}(0) = 1$ electric charge
 - $F_2^{(v)} = \kappa^{(v)} \left[1 + \frac{1}{6} \langle r_2^2 \rangle^{(v)} q^2 + \mathcal{O}(q^4) \right]$
 $\kappa^{(v)} = F_M^{(v)}(0) = \mu^{(v)} - 1$ magnetic moment
 - $\langle r_E^2 \rangle^{(v)} = \langle r_1^2 \rangle^{(v)} + \frac{3}{2m_N^2} \kappa^{(v)}$ proton radius puzzle

	$\sigma(p^3) \Delta$	$\sigma(p^4) \Delta$	$\sigma(p^3) \Delta$	$\sigma(p^4) \Delta$
\tilde{g}_A (free)	1.1782 ± 0.0073		1.2041 ± 0.0074	1.274 ± 0.041
d_{16} (GeV^{-2}) (free)	-1.021 ± 0.048		0.983 ± 0.062	-1.46 ± 1.00
d_{22} (GeV^{-2}) (free)	1.275 ± 0.086		3.77 ± 1.96	0.29 ± 1.69 (free g_1)
h_A	-	-	1.35	1.35
g_1 (free)	-	-	-0.69 ± 0.69	0.66 ± 0.56
c_1 (GeV^{-1})	-	-0.89 ± 0.06	-	-1.15 ± 0.05
c_2 (GeV^{-1})	-	3.38 ± 0.15	-	1.57 ± 0.10
c_3 (GeV^{-1})	-	-4.59 ± 0.09	-	-2.54 ± 0.05
c_4 (GeV^{-1})	-	3.31 ± 0.13	-	2.61 ± 0.10
a_1 (GeV^{-1})	-	-	-	0.90
b_1 (GeV^{-2}) (free)	-	-	-	-0.27 ± 4.96
b_2 (GeV^{-2}) (free)	-	-	-	2.27 ± 2.28
\tilde{b}_4 (GeV^{-2}) (free)	-	-	-	-12.48 ± 1.28
x_1 (fm^{-2}) (free)	-8.4 ± 5.8	-	-5.6 ± 5.9	-0.25 ± 16.5 (consistent)
x_2 (fm^{-2}) (free)	-8.6 ± 2.6	-	-7.1 ± 2.6	-6.36 ± 4.20
x_3 (fm^{-1}) (free)	-0.25 ± 0.21	-	-0.08 ± 0.22	0.36 ± 0.47
y_1 ($\text{fm}^{-2} \text{GeV}^{-2}$) (free)	-100 ± 40	-	-76 ± 44	-64 ± 121
y_2 ($\text{fm}^{-2} \text{GeV}^{-2}$) (free)	-31 ± 21	-	-21 ± 22	-15 ± 46
y_3 ($\text{fm}^{-1} \text{GeV}^{-2}$) (free)	-0.63 ± 1.49	-	0.36 ± 1.63	2.54 ± 3.98
\tilde{m} (GeV)	0.874	0.874	0.855	0.855
\tilde{m}_Δ (GeV)	-	-	1.166	1.166

	$\sigma(p^3) \Delta$	$\sigma(p^4) \Delta$	$\sigma(p^3) \Delta$	$\sigma(p^4) \Delta$
\tilde{g}_A (free)	1.1782 ± 0.0073		1.2041 ± 0.0074	1.274 ± 0.041
d_{16} (GeV^{-2}) (free)	-1.021 ± 0.048		0.983 ± 0.062	-1.46 ± 1.00
d_{22} (GeV^{-2}) (free)	1.275 ± 0.086		3.77 ± 1.96	0.29 ± 1.69 (free g_1)
h_A	-	-	1.35	1.35
g_1 (free)	-	-	-0.69 ± 0.69	0.66 ± 0.56
c_1 (GeV^{-1})	-	-0.89 ± 0.06	-	-1.15 ± 0.05
c_2 (GeV^{-1})	-	3.38 ± 0.15	-	1.57 ± 0.10
c_3 (GeV^{-1})	-	-4.59 ± 0.09	-	-2.54 ± 0.05
c_4 (GeV^{-1})	-	3.31 ± 0.13	-	2.61 ± 0.10
a_1 (GeV^{-1})	-	-	-	0.90
b_1 (GeV^{-2}) (free)	-	-	-	-0.27 ± 4.96
b_2 (GeV^{-2}) (free)	-	-	-	2.27 ± 2.28
\tilde{b}_4 (GeV^{-2}) (free)	-	-	-	-12.48 ± 1.28
x_1 (fm^{-2}) (free)	-8.4 ± 5.8	-	-5.6 ± 5.9	-0.25 ± 16.5 (consistent)
x_2 (fm^{-2}) (free)	-8.6 ± 2.6	-	-7.1 ± 2.6	-6.36 ± 4.20
x_3 (fm^{-1}) (free)	-0.25 ± 0.21	-	-0.08 ± 0.22	0.36 ± 0.47
y_1 ($\text{fm}^{-2} \text{GeV}^{-2}$) (free)	-100 ± 40	-	-76 ± 44	-64 ± 121
y_2 ($\text{fm}^{-2} \text{GeV}^{-2}$) (free)	-31 ± 21	-	-21 ± 22	-15 ± 46
y_3 ($\text{fm}^{-1} \text{GeV}^{-2}$) (free)	-0.63 ± 1.49	-	0.36 ± 1.63	2.54 ± 3.98
\tilde{m} (GeV)	0.874	0.874	0.855	0.855
\tilde{m}_Δ (GeV)	-	-	1.166	1.166
χ^2/dof	$46.13/(127 - 9) = 0.391$		$39.17/(127 - 10) = 0.326$	$14.64/(127 - 13) = 0.129$ Δ trunc. overestim.
χ_0^2/dof	$857.31/(127 - 9) = 7.27$		$533.87/(127 - 10) = 4.45$	$196.58/(127 - 13) = 1.724$

	$\mathcal{O}(p^4) \Delta$
\tilde{g}_A (free par.)	1.240 ± 0.046
d_{16} (free par.)	-0.88 ± 0.88
h_A	$h^{Nc} = 1.35$
g_1	$- g_1^A = -2.29$
c_1	-1.15 ± 0.05
c_2	1.57 ± 0.10
c_3	-2.54 ± 0.05
c_4	2.61 ± 0.10
a_1	0.90
\tilde{b}_4 (free par.)	-12.3 ± 1.0
\tilde{m}	0.855
\tilde{m}_Δ	1.166
χ^2/dof	$11.14/(43 - 7) = 0.31$